

**CHILDRENS BENADRYL ALLERGY- diphenhydramine hydrochloride solution**  
**Kenvue Brands LLC**

-----  
**Children's Benadryl® ALLERGY**

***Drug Facts***

**Active ingredient (in each 5 mL)**

Diphenhydramine HCl 12.5 mg

**Purpose**

Antihistamine

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat

**Warnings**

**Do not use**

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if the child has**

- a breathing problem such as chronic bronchitis
- glaucoma
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if the child is** taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- find right dose on chart below
- mL = milliliter
- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 2 years	do not use
children 2 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL to 10 mL

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

## Other information

- **each 5 mL contains:** sodium 14 mg
- store between 20-25°C (68-77°F). Protect from light. Store in outer carton until contents used.
- **do not use if bottle wrap or foil inner seal imprinted with “Sealed For Your Safety” is broken or missing**

## Inactive ingredients

anhydrous citric acid, D&C red no. 33, FD&C red no. 40, flavors, glycerin, monoammonium glycyrrhizinate, poloxamer 407, purified water, sodium benzoate, sodium chloride, sodium citrate, sucrose

## Questions or comments?

call **1-877-717-2824** or **215-273-8755 (collect)**

## PRINCIPAL DISPLAY PANEL

NDC 50580-534-04

Children's

Benadryl®

ALLERGY

Diphenhydramine HCl/antihistamine

12.5 mg/5 mL oral solution

4-6 Hours/Dose

RELIEF OF:

- Runny Nose
- Sneezing
- Itchy, Watery Eyes
- Itchy Throat or Nose

Alcohol Free

Cherry!

Flavored

4 fl oz (118 mL)

Children's  
**Benadryl**<sup>®</sup>

**ALLERGY**  
ALERGIA

*How can we help?*

*¿Cómo podemos ayudarlo?*

**1-877-717-2824**  
**benadryl.com**

NDC 50580-534-04

Children's  
**Benadryl**<sup>®</sup>

**ALLERGY**  
ALERGIA

**Diphenhydramine HCl/antihistamine**  
**12.5 mg/5 mL oral solution**

**Difenhidramina ClH/antihistamínico**  
**12.5 mg/5 ml solución oral**

**4-6** Hours/Dose  
Horas/Dosis

**RELIEF OF:**  
ALIVIA:

 **Runny Nose**  
Secreción nasal

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
Diphenhydramine HCl 12.5 mg.....	Antihistamine

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat

**Warnings**

- Do not use**
- to make a child sleepy
  - with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if the child has**

- a breathing problem such as chronic bronchitis
- glaucoma
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if the child is taking** sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

✓ Sneezing  
Estornudos

✓ Itchy, Watery Eyes  
Lagrimo, picazón en los ojos

✓ Itchy Throat or Nose  
Picazón de garganta o nariz



Alcohol Free  
Sin alcohol



**Cherry!** **iCereza!**  
Flavored Saborizado

4 fl oz (118 mL)



LOT/EXP

### Directions

- find right dose on chart below ■ mL = milliliter
- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 2 years	do not use
children 2 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL to 10 mL

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.


### Other information

- each 5 mL contains: sodium 14 mg
- store between 20-25°C (68-77°F). Protect from light. Store in outer carton until contents used.
- do not use if bottle wrap or foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

**Inactive ingredients** anhydrous citric acid, D&C red no. 33, FD&C red no. 40, flavors, glycerin, monoammonium glycyrrhizinate, poloxamer 407, purified water, sodium benzoate, sodium chloride, sodium citrate, sucrose

### Questions or comments?

call 1-877-717-2824 or 215-273-8755 (collect)



Children's  
**Benadryl**

ALLERGY  
ALERGIA

Etiqueta de información

### Etiqueta de información (continuación)

#### Posología

- encuentre la dosis adecuada en el siguiente cuadro
- ml = mililitro
- tomar cada 4 a 6 horas, o según lo indicado por el médico
- no tomar más de 6 dosis en 24 horas

Edad (años)	Dosis (ml)
niños menores de 2 años	no usar
niños de 2 a 5 años	no usar, salvo por indicación médica
niños de 6 a 11 años	5 ml a 10 ml

## Etiqueta de información

### **Ingrediente activo Acción terapéutica (cada 5 ml)**

Difenhidramina ClH 12.5 mg.....Antihistamínico

### **Indicaciones**

- alivia temporalmente estos síntomas por rinitis alérgica u otras alergias del tracto respiratorio superior:
  - secreción nasal
  - estornudos
  - lagrimeo, picazón en los ojos
  - comezón en la nariz o la garganta

### **Advertencias**

#### **No usar**

- para dormir a un niño
- con otro producto que contenga difenhidramina, incluso si es de uso tópico

#### **Consulte a su médico antes de usar si el niño**

- tiene problemas respiratorios, como bronquitis crónica
- tiene glaucoma
- sigue una dieta baja en sodio

#### **Consulte a un médico o farmacéutico antes de usar este producto si el niño toma sedantes o tranquilizantes**

#### **Al usar este producto**

- puede sentir una marcada somnolencia
- los sedantes y tranquilizantes pueden aumentar la somnolencia
- puede causar excitabilidad, especialmente en niños

**Mantener fuera del alcance de los niños.** En caso de sobredosis, busque ayuda médica o comuníquese con un centro de toxicología de inmediato. (1-800-222-1222) ▶

niños de 6 a 11 años 1.5 ml a 10 ml

**Atención:** usar solo el dosificador que se incluye, diseñado exclusivamente para usar con este producto. No usar ningún otro dispositivo de dosificación.

### **Otra información**

- **cada 5 ml contiene:** sodio 14 mg
- almacenar a 20-25 °C (68-77 °F). Proteger de la luz. Almacenar en el envase exterior hasta haber utilizado su contenido.
- **no usar si se ha roto o falta el envoltorio de la botella o el sello interior de papel aluminio impreso con las palabras "Sealed For Your Safety"**

### **Ingredientes inactivos**

ácido cítrico anhidro, rojo n.° 33 D&C, rojo n.° 40 FD&C, saborizantes, glicerina, glicirricinato de monoamonio, poloxámero 407, agua purificada, benzoato de sodio, cloruro de sodio, citrato de sodio, sacarosa

### **¿Tiene preguntas o comentarios?**

llame al **1-877-717-2824** o al **215-273-8755** (cobro revertido)

Active ingredient made in Japan /  
Ingrediente activo hecho en Japón

Distributed by / Distribuido por:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
©J&JCI 2022  
Pat. www.jjciipats.com 30053652

**For more information:**  
Para obtener más información:

**www.benadryl.com**

## CHILDRENS BENADRYL ALLERGY

diphenhydramine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-534
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 5 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>AMMONIUM GLYCYRRHIZATE</b> (UNII: 3VRD35U26C)	
<b>POLOXAMER 407</b> (UNII: TUF21VW3M2)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SUCROSE</b> (UNII: C151H8M554)	

**Product Characteristics**

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:50580-534-04	1 in 1 CARTON	07/01/2008	
1		118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
2	NDC:50580-534-08	1 in 1 CARTON	07/01/2008	
2		236 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
3	NDC:50580-534-16	2 in 1 PACKAGE	01/28/2009	12/28/2022
3		1 in 1 CARTON		
3		236 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
4	NDC:50580-534-18	2 in 1 PACKAGE	04/12/2021	
4		1 in 1 CARTON		
4		236 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2008	

**Labeler** - Kenvue Brands LLC (118772437)

Revised: 5/2025

Kenvue Brands LLC