WALGREENS WAL-FLU COLD AND SORE THROAT LEMON FLAVOR- acetaminophen, pheniramine maleate, phenylephrine hcl granule, for solution Walgreen Co.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreens Wal-Flu Cold and Sore Throat Lemon Flavor

Uses

- temporarily relieves these symptoms due to a cold:
- minor aches and pains headache minor sore throat pain
- nasal congestion temporarily reduces fever
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose sneezing itchy nose and throat itchy, watery eyes.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take - more than 6 packets in 24 hours, which is the maximum daily amount - with other drugs containing acetaminophen - 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: - skin reddening - blisters - rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a physician promptly.

Do not use:

- In a child under 4 years of age.
- If you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription and nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease heart disease high blood pressure thyroid disease diabetes glaucoma trouble urinating due to an enlarged prostate gland cough that occurs with too much phlegm (mucus) cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin.

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks may cause drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.

Overdose Warning:

In case of overdose, get medical help or contact a poison control center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor

Age	Dose
children under 4 years of age	do not use
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

- dissolve contents of one packet in 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- If using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other Information

- each packet contains: sodium 66 mg
- store at controlled room temperature 20-25C (68-77F). Protect from heat and moisture.

Citric acid, D and C Yellow 10, Ethylcellulose, FD and C Red 40, Flavor, Microcrystalline cellulose, Povidone, Silicon dioxide, Sodium citrate, Starch, Sugar and Vitamin C.



WALGREENS WAL-FLU COLD AND SORE THROAT LEMON FLAVOR

acetaminophen, pheniramine maleate, phenylephrine hcl granule, for solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0601
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg	
PHENIRAMINE MALEATE (UNII: NYW905655B) (PHENIRAMINE - UNII:134FM9ZZ6M)	PHENIRAMINE MALEATE	20 mg	
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6 MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg	

Inactive Ingredients	
Ingredient Name	Strength

CITRIC ACID MO NO HYDRATE (UNII: 2968 PHW8 QP)	
ETHYLCELLULOSE (100 MPA.S) (UNII: 47MLB0F1MV)	
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
PO VIDO NE (UNII: FZ989 GH94E)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
SO DIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	

Product Characteristics			
Color	yellow (D and C Yellow 10), red (FD and C Red 40)	Score	
Shape		Size	
Flavor	LEMON (Nature Lemon Flavor)	Imprint Code	
Contains			

I	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0601-06	6 in 1 BOX; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/07/2013	

Labeler - Walgreen Co. (008965063)

Revised: 3/2016 Walgreen Co.