VICKS NYQUIL COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid The Procter & Gamble Manufacturing Company

VICKS ® NyQuil COLD & FLU

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg Dextromethorphan HBr 30 mg Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer Cough suppressant Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- · avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not

notice any signs or symptoms.

Directions

- · take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains:sodium 24 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

alcohol, citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label

VICKS®

NyQuil™

COLD & FLU

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

- Aches, Fever & Sore Throat
- Sneezing, Runny Nose
- Cough

Nighttime Relief

Alcohol 10%

8 FL OZ (236 ml)







Drug Facts (continued) Warnings Liver warning: This product contains acetaminophen. Severe bronchitis or emphysema • trouble urinating due to enlarged prostate gland err warming: Inis product contains accaminopient. Severe of danage may occur if you take • more than 4 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers taking the blood thinning drug warfarin When using this product - excitability may occur, especially in children - marked drowsiness may occur - avoid alcoholic drinks - be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase ight away fore throat warning: If sore throat is severe, persists for more han 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. op use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • lever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • rough comes back or occurs with rash or headache that lasts. Stop use and ask a doctor if Do not use with any other drug containing acetaminophen (precorption or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MMO) (pertain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, If you do not know if your prescription drug contains an IMOI, ask a doctor or pharmacist before taking this product. These could be signs of a serious condition. I pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children Ask a doctor before use if you have • liver disease • glaucoma • ough that occurs with too much phlegm (mucus) • a breathing problem or chronic ough that lasts or as occurs with smoking, asthma, chronic even if you do not notice any signs or symptoms.



VICKS NYQUIL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-791
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL	

Inactive Ingredients		
Ingredient Name	Strength	
SUCRALOSE (UNII: 96K6UQ3ZD4)		
SORBITOL (UNII: 506T60A25R)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
GLYCERIN (UNII: PDC6A3C0OX)		
ALCOHOL (UNII: 3K9958V90M)		
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
WATER (UNII: 059QF0KO0R)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE (UNII: 1Q73Q2JULR)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	CHERRY, MENTHOL	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423- 791-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/01/2022	
2	NDC:69423- 791-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/01/2022	
3	NDC:69423- 791-24	2 in 1 PACKAGE, COMBINATION	01/01/2022	
3	NDC:69423- 791-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:69423- 791-36	3 in 1 PACKAGE, COMBINATION	01/01/2022	
4	NDC:69423- 791-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2022	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 11/2024 The Procter & Gamble Manufacturing Company