

CHILDRENS ROBITUSSIN COUGH AND COLD CF- dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid
Haleon US Holdings LLC

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

Phenylephrine HCl, USP 5 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

1. heart disease
2. high blood pressure
3. thyroid disease
4. diabetes
5. trouble urinating due to an enlarged prostate gland
6. cough that occurs with too much phlegm (mucus)
7. cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- do not take more than 6 doses in any 24-hour period

| age | dose |
|---------------------------------------|---------------------|
| children under 4 years | do not use |
| children 4 to under 6 years | 5 mL every 4 hours |
| children 6 to under 12 years | 10 mL every 4 hours |
| adults and children 12 years and over | 20 mL every 4 hours |

Other information

- **each 10 mL contains:**sodium 7 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, artificial flavor, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

call weekdays from 9 AM to 5 PM EST at **1-800-245-1040**

Additional Information

Packaged with Tamper-Evident bottle cap. Do Not Use if breakable ring is separated or missing.

Children's

Robitussin liquid

is specially

Children's Robitussin liquid is especially formulated to provide soothing action, control your child's cough plus relieve other cold symptoms.

Should be 18 or older to purchase

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Distributed by: GSK Consumer Healthcare,

Warren, NJ 07059

For most recent product information,

visit www.robitussin.com

Trademarks owned or licensed by GSK.

©2022 GSK or licensor

Made in Canada

PRINCIPAL DISPLAY PANEL

Children's

Robitussin

FOR AGES 4 & OVER

Cough & Cold

CF

DEXTROMETHORPHAN HBr

(Cough Suppressant)

GUAIFENESIN (Expectorant)

PHENYLEPHRINE HCl (Nasal Decongestant)

Relieves:

- **Cough**
- **Chest Congestion/Mucus**
- **Stuffy Nose**

Non-Drowsy

grape

flavor

4 FL OZ

(118 mL)

PAA172104 Front Carton

Children's

Robitussin

FOR AGES 4 & OVER

Cough & Cold

CF

DEXTROMETHORPHAN HBr
(Cough Suppressant)
GUAIFENESIN (Expectorant)
PHENYLEPHRINE HCl (Nasal Decongestant)

Relieves:

- ✓ Cough
- ✓ Chest Congestion/Mucus
- ✓ Stuffy Nose

Non-Drowsy



grape
flavor



4 FL OZ
(118 mL)

CHILDRENS ROBITUSSIN COUGH AND COLD CF

dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0031-8716 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|-------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 5 mg in 5 mL |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 50 mg in 5 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 2.5 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL, (R)- (UNII: 602HN5L69H) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |

Product Characteristics

| | | | |
|-----------------|-----------------|---------------------|--|
| Color | red (clear red) | Score | |
| Shape | | Size | |
| Flavor | GRAPE | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0031-8716-12 | 1 in 1 CARTON | 09/01/2009 | |
| 1 | | 118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M001 | 09/01/2009 | |

