NITE-TIME COLD/FLU ORIGINAL- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid Salado Sales, Inc.

Nite Time 501

Drug FactsActive ingredients (in each 30 mL dose cup)

Acetaminophen 650 mg Dextromethorphan HBr 30 mg Doxylamine Succinate 12.5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Keep Out Of Reach Of Children Keep out of reach of children.

Uses

temporarily relieves cold/flu symptoms:

- sore throat
- minor aches and pain
- runny nose and sneezing
- cough due to minor sore throat and bronchial irritation
- headache
- fever

Warnings

Failure to follow these warnings could result in serious consequences

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amoun
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks
- every day while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash

If a skin reaction ccurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Ask a doctor before use if you have

- a sodium restricted diet
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask A Doctor or Pharmacist before use

- if you are taking sedatives or tranquilizers
- if you are taking the blood thinning drug warfarin

When Using This Product

- do not use more than directed
- avoid alcoholic drinks
- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop Use And Ask A Doctor If

- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash, headache that lasts

Pregnant Or Breast-Feeding

ask a health professional before use.

Overdose Warning

Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as recommended-see Overdose warnings
- Use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hours
- If taking Nite Time at night and Day Time during the day, limit total to 4 doses per 24 hours



Other Information

- each 30 mL dose cup contains:
- sodium 45 mg
- store at room temperature

Inactive ingredients

citric acid, D and C yellow No. 10, FD and C Green No. 3, FD and C Yellow No. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- to make a child sleepy

Product Label

Compare to VICKS® NYQUIL® active ingredients CVP CONSUMER VALUE PRODUCTS

Nite-Time COLD/FLU Medicine

Multi-Symptom ORIGINAL FLAVOR

AcetaminophenAches, Fever and Sore Throat

DextromethorphanCough

Doxylamine SuccinateSneezing, Runny nose

6 FL OZ (177 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

*This product is not manufactured or distributed by Proctor and Gamble, distributor of Vicks® Nyquil® LR-002

Questions? Call weekdays from 9:30am to 4:30pm EST 1-877-795-5944

Distributed by: Salado Sales, Inc. P.O. Box 6115 Temple, TX 76503 www.saladosales.com

LOT: EXP:

Peel Here DRUG FACTS CONTINUED ON BACK

ORIGINAL FLAVOR Nite-Time Multi-Symptom COLD/FLU Medicine

- Antihistamine
- Cough Suppressant
- Pain Reliever
- Fever Reducer

MULTI-SYMPTOM COLD/FLU MEDICINE

ORIGINAL FLAVOR

- ANTIHISTAMINE
- COUGH SUPPRESSANT
- · PAIN RELIEVER
- FEVER REDUCER

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, nausea, or womiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug containing acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine codase inhibitor (MADI) (certain drugs for depression, psychiatric or emotion conditions, or Parkinson's disease), or for 2 weeks after stoping the MADI (any, if you do not know if your prescription drug contains an MADI, ask a doctor or pharmacist before taking this product.
• to make a child sleepy

Ask a doctor before use if you have • a sodium restricted diet • liver disease • glaucoma • cough that occurs with too much philogin (inucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble unnating due to enlarged prostate gland.

Ask a doctor or pharmacist before use • if you are taking sedatives or tranquilizers • if you are taking ng drug wartarin

When using this product • do not use more than directed • avoid alcoholic drinks • excitability may occur, especially in children • marked drowsiness may occur • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN



*This product is not manufactured or distributed by Proctor & Gamble, distributor of Vicks® NyQuil®.

Questions? Call weekdays from 9:30am to 4:30pm EST at 1-877-798-5944

Distributed by: Salado Sales, Inc P.O. Box 5115 Temple, TX 76503 www.saladosales.com

5

EXP.

Drug Facts

ON BACK

CONTINUED

Failure to follow these warnings could result in serious consequences

Active ingredients (in each 30 mL dose cup) Purpose

- Pain reli

Uses temporarily relieves cold/flu symptoms:

sore throat

ninor aches and pain

sunny nose and sneezing

cough due to minor throat and bronchial irritation

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take * more than 4,000mg of acetaminophen in 24 hours * with other drugs containing acetaminophen * 3 or more alcoholic drinks every day white using this product. Alterny sizer Acetaminophen may cause severe skin reactions. Symptons may include: * \$\text{skin reactions. Symptons may include: * \$\text{s



ORIGINAL FLAVOR

- ACETAMINOPHEN ACHES, FEVER & SORE THROAT
- DEXTROMETHORPHAN HBr COUGH SUPPRESSANT
- · DOXYLAMINE SUCCINATE SNEEZING, RUNNY NOSE

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN



This product is not manufactured or distributed by Proctor & Gambi distributor of Vicks® NyQuil®

Questions? Call weekdays from 9:30am to 4:30pm EST at 1-877-798-5944

LR-002 Rev 01

Salado Sales, Inc. P.O. Box 6115 Temple, TX 76503 www.saladosales.com

5

Drug Facts (continued)

Stop use and ask a doctor if * nedness or swelling is present * symptoms do not get better within 7 days or are accompanied by a fever * fever gets worse or lasts more than 3 days * new symptoms occur * cough lasts more than 7 days, comes back, or occurs with fever rash, or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Keep out or reach or consumers. Overvidese warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poision Control Center right away. Quick medical alteration is critical for adults as well as for children even if you do not notice any

Directions • take only as recommended-see Overdose warning • the dose cup or tablespoon (TISSP)

do not exceed 4 doses per 24 hox

Drug Facts (continued)

adults & children 12 years and over | 30 mL (2 TBSP) every 6 hours children 4 to under 12 years ask a doctor children under 4 years do not use children under 4 years

If taking Nite Time at night and Day Time during the day, limit total to 4 doses per 24 hours.

Other information • each 30 mL dose cup contains sodium 45mg • store at room temperature

Inactive ingredients altric acid, D&C yellow # 10, FD&C Green No. 3, FD&C Yellow No. 6, flavor, glycorin, polyethylene glycol, propytere glycol, purified water, saccharin sodium, sodium benzoate, sodium cetate, success

99



res

NITE-TIME COLD/FLU ORIGINAL

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57243-501	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE -	DOYYLAMINE SUCCINATE	12.5 mg	

Inactive Ingredients			
Ingredient Name	Strength		
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
GLYCERIN (UNII: PDC6A3C0OX)			
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
SUCROSE (UNII: C151H8M554)			

Packaging				
#	# Item Code Package Description		Marketing Start Date	Marketing End Date
1	NDC:57243-501- 25	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/16/2012	

Marketing Information				
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	01/16/2012		

Labeler - Salado Sales, Inc. (009830555)

Registrant - AptaPharma Inc. (790523323)

Establishment				
Name	Address	ID/FEI	Business Operations	
AptaPharma Inc.		790523323	manufacture(57243-501)	

Revised: 12/2023 Salado Sales, Inc.