

**VICKS DAYQUIL SEVERE NYQUIL SEVERE HONEY FLAVOR COLD AND FLU-  
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl and  
doxylamine succinate  
The Procter & Gamble Manufacturing Company**

-----

**VICKS ® DayQuil™ SEVERE HONEY FLAVOR | VICKS ® NyQuil™ SEVERE HONEY  
FLAVOR  
COLD & FLU Convenience Pack**

VICKS ® NyQuil™ SEVERE HONEY FLAVOR  
COLD & FLU

## **Drug Facts**

### **Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

### **Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

## **Warnings**

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks daily while using this product

### **Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers

- taking the blood thinning drug warfarin

### **When using this product**

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

---

### **Other information**

- **each 15 mL contains:** sodium 14 mg

- Store at no greater than 25°C and do not refrigerate.

citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

## **Questions?**

1-800-362-1683

VICKS<sup>®</sup> DayQuil<sup>™</sup> SEVERE HONEY FLAVOR

COLD & FLU

## **Drug Facts**

### **Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

### **Uses**

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

## **Warnings**

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks daily while using this product

### **Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product,**

## **do not use more than directed**

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

---

### **Other information**

- **each 15 mL contains:**sodium 14 mg
- Store at no greater than 25°C and do not refrigerate

### **Inactive ingredients**

citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

### **Questions?**

1-800-362-1683

**TAMPER EVIDENT:** Do not use if printed shrinkband seal around the neck is broken or missing.

Made in Canada

DIST. BY PROCTER & GAMBLE,

CINCINNATI, OH 45202

**VICKS® DayQuil™ SEVERE HONEY FLAVOR | VICKS® NyQuil™ SEVERE HONEY FLAVOR  
COLD & FLU Convenience Pack**

MAX STRENGTH

COLD & FLU

DAY & NIGHT PACK

VICKS® DayQuil™ SEVERE HONEY FLAVOR

COLD & FLU

**Acetaminophen**, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Chest Congestion, Thins & Loosens Mucus

Nasal Congestion, Sinus Pressure

Cough

*Non-Drowsy*

VICKS® NyQuil™ SEVERE HONEY FLAVOR

COLD & FLU

**Acetaminophen**, Dextromethorphan HBr, Doxylamine succinate, Phenylephrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

*Nighttime Relief*

2 Bottles - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



# VICKS DAYQUIL SEVERE NYQUIL SEVERE HONEY FLAVOR COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl and doxylamine succinate kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-808

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-808-24	1 in 1 CARTON; Type 0: Not a Combination Product	04/28/2020	

Quantity of Parts			
Part #	Package Quantity		Total Product Quantity
Part 1	1 BOTTLE		354 mL
Part 2	1 BOTTLE		354 mL



Part 1 of 2

VICKS DAYQUIL SEVERE HONEY FLAVOR COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Item Code (Source)	NDC:69423-806
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
GLYCERIN (UNII: PDC6A3C0OX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

Product Characteristics

Color	brown	Score	
Shape		Size	
Flavor	HONEY	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/28/2020	

Part 2 of 2

VICKS NYQUIL SEVERE HONEY FLAVOR COLD AND FLU

acetaminophen, dextromethrophan, doxylamine succinate, phenylephrine hcl liquid

Product Information	
Item Code (Source)	NDC:69423-807
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients	
Ingredient Name	Strength
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SORBITOL (UNII: 506T60A25R)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	HONEY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/28/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/28/2020	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2024

The Procter & Gamble Manufacturing Company