

**VICKS DAYQUIL SEVERE NYQUIL SEVERE HONEY FLAVOR COLD AND FLU-
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl and
doxylamine succinate
The Procter & Gamble Manufacturing Company**

**VICKS[®] DayQuil[™] SEVERE HONEY FLAVOR | VICKS[®] NyQuil[™] SEVERE HONEY
FLAVOR
COLD & FLU Convenience Pack**

VICKS[®] NyQuil[™] SEVERE HONEY FLAVOR
COLD & FLU

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks daily while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers

- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

| | |
|------------------------------------|---|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | do not use unless directed by a doctor |
| children under 4 yrs | do not use |

Other information

- **each 15 mL contains:** sodium 14 mg

- Store at no greater than 25°C and do not refrigerate.

citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-800-362-1683

VICKS[®] DayQuil[™] SEVERE HONEY FLAVOR
COLD & FLU

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks daily while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product,

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

| | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- **each 15 mL contains:**sodium 14 mg
- Store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband seal around the neck is broken or missing.

Made in Canada

DIST. BY PROCTER & GAMBLE,

CINCINNATI, OH 45202

**VICKS® DayQuil™ SEVERE HONEY FLAVOR | VICKS® NyQuil™ SEVERE HONEY FLAVOR
COLD & FLU Convenience Pack**

MAX STRENGTH

COLD & FLU
DAY & NIGHT PACK

VICKS® DayQuil™ SEVERE HONEY FLAVOR

COLD & FLU

Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains
Chest Congestion, Thins & Loosens Mucus

Nasal Congestion, Sinus Pressure
Cough

Non-Drowsy

VICKS® NyQuil™ SEVERE HONEY FLAVOR

COLD & FLU

Acetaminophen, Dextromethorphan HBr, Doxylamine succinate, Phenylephrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

Nighttime Relief

2 Bottles - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



VICKS DAYQUIL SEVERE NYQUIL SEVERE HONEY FLAVOR COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl and doxylamine succinate kit

Product Information

| | | | |
|---------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69423-808 |
|---------------------|----------------|---------------------------|---------------|

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:69423-808-24 | 1 in 1 CARTON; Type 0: Not a Combination Product | 04/28/2020 | |

Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 1 BOTTLE | 354 mL |
| Part 2 | 1 BOTTLE | 354 mL |

Part 1 of 2

VICKS DAYQUIL SEVERE HONEY FLAVOR COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Item Code (Source) NDC:69423-806

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|--------------------|
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg in 15 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 15 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg in 15 mL |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg in 15 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S) | |
| SORBITOL (UNII: 506T60A25R) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |

Product Characteristics

| | | | |
|----------|-------|--------------|--|
| Color | brown | Score | |
| Shape | | Size | |
| Flavor | HONEY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|---|----------------------|--------------------|
| 1 | | 354 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 04/28/2020 | |

Part 2 of 2

VICKS NYQUIL SEVERE HONEY FLAVOR COLD AND FLU

acetaminophen, dextromethorphan, doxylamine succinate, phenylephrine hcl liquid

Product Information

| | |
|-------------------------|---------------|
| Item Code (Source) | NDC:69423-807 |
| Route of Administration | ORAL |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|---------------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg in 15 mL |
| DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 6.25 mg in 15 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 15 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg in 15 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| SORBITOL (UNII: 506T60A25R) | |
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |

| | |
|---|--|
| XANTHAN GUM (UNII: TTV12P4NEE) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S) | |

| Product Characteristics | | | |
|--------------------------------|-------|---------------------|--|
| Color | brown | Score | |
| Shape | | Size | |
| Flavor | HONEY | Imprint Code | |
| Contains | | | |

| Packaging | | | | |
|------------------|-----------|---|----------------------|--------------------|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | | 354 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

| Marketing Information | | | |
|------------------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug | M012 | 04/28/2020 | |

| Marketing Information | | | |
|------------------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug | M012 | 04/28/2020 | |

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2025

The Procter & Gamble Manufacturing Company