# MEJORALITO- acetaminophen suspension OPMX Chula Vista

-----

### Mejoralito

## Active Ingredient(in each 5 mL) Purpose

Acetaminophen 160 mg...... Pain reliever/fever reducer

Pain Reliever & Fever Reducer

#### Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- sore throat
- the common cold
- flu
- headache
- toothache

### Warnings

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if your child takes takes

- more than 5 doses in 24 hours. which is the maximum daily amount.
- with other drugs containing acetaminophen.

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommened dose (see overdose

### warning)

### Stop use and ask a doctor if

- new sympthoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition

### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- this product does not contain directions or complete warnings for adult use. Shake well before using
- mL= millilitar
- find right dose on chart below, if possible, use weight to dose; otherwise, use age
- repeat dose every 4 hours while symptoms last
- only use enclosed measuring syringe
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor

Weight (lb,)	Age (yrs.)	Dose (mL)*
Under 24	Under 2	Ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

<sup>\*</sup>always check with your doctor before administering medicine to your infant/child

#### Other information

- Each teaspoon (5 mL) contains: sodium 3 mg
- Store between 15°-30°C (59°-86°F)
- Protect from freezing. Protect from light

# Inactive ingredients

acesulfame potassium, citric acid, FD&C red no. 40, flavor, glycerine, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, polysorbate, propylene glycol, prosweet N&AK, purified water, sodium benzoate, sucralose, sorbitol,

xanthan gum.

# Questions or comments?

Call toll free 619-600-5632

Monday through Fnday 9AM - 5PM EST

## PRINCIPAL DISPLAY PANEL

NDC 69729-068-04 Temperal-ito





Tamper Evident: Do not use this product if the safety

### **Drug Facts**

Active ingredient (in each 5 mL) Purpose

**Uses** temporarily: ■ reduces fever

■ relieves minor aches and pains due to: ■ sore throat
■ the common cold ■ flu ■ headache ■ toothache

#### Warnings

Liver warning: This product contains acetaminophen.
Severe liver damage may occur if your child takes

more than 5 doses in 24 hours, which is the maximum

daily amount

with other drugs containing acetaminophen

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a depter promptly. doctor promptly.

#### Do not use

with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

# When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

new symptoms occur redness or swelling is present
pain gets worse or lasts more than 5 days
fever gets worse or lasts more than 3 days
These could be signs of a serious condition.

#### Keep out of reach of children

Neep out or reach or children

Overdose warning: Taking more than the
recommended dose (overdose) may cause liver damage.
In case of overdose, get medical help or contact a Poison
Control Center right away, Outke medical attention is
critical for adults as well as for children even if you do not notice any signs or symptoms.





4 fl oz (118 mL) 160 mg cada 5 mL

#### Drug Facts (continued)

#### Directions

- Directions

  It is product does not contain directions or complete warnings for adult use. Shake well before using

  Int. = milliliter

  Interpret does on chart below. If possible, use weight to dose; otherwise, use age.

  In eppart does every 4 hours while symptoms last

  only use enclosed measuring cup

  Id not give more than 5 times in 24 hours

  do not give more than 5 times in 24 hours

weight (ib.)	Age (yrs.)	Dose (mi)	
Under 24	Under 2	Ask a doctor	
24 - 35	2-3	5mL	
36 - 47	4-5	7.5 mL	
48 - 59	6-8	10mL	
60 - 71	9-10	12.5mL	
72-95	11	15mL	
talium a aleasti with come destante fores			

always check with your doctor before administering medicine to your infant/child

- each teaspoon (5 ml.) contains: sodium 3 mg
   Store between 15'-30' C (59'-86" F)
   Protect from Freezing. Protect from light

#### Inactive ingredients

Inactive Ingreatents
acesulfame potssium, citric acid, FD&C red no. 40, flavor,
glycerin, high fructose corn syrup, microcrystalline cellulose
and carboxymethylicalliose sodium, polysorbate, propylene
glycol, proswer N & K, furpfied water, sodium benzoate,
sucralose, sorbitol, xanthan gum

#### Questions or comments

Call toll free 619-600-5632 Monday through Friday 9AM - 5PM EST

#### VERSION EN ESPAÑOL EN EL INTERIOR DE LA CAJA

⊚РМ̈́Х



# Lot: Exp: NON VARNISH AREA

# **MEJORALITO**

acetaminophen suspension

#### **Product Information**

ı	<b>Product Type</b>	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-068
ı	Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
POLYSORBATE 20 (UNII: 7T1F30V5YH)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
SORBITOL (UNII: 506T60A25R)			
XANTHAN GUM (UNII: TTV12P4NEE)			
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)			
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			

Product Characteristics		
Color		Score
Shape		Size
Flavor	CHERRY	Imprint Code
Contains		

I	Packaging					
4	tem Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:69729-068- 04	1 in 1 CARTON	06/02/2023			
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/02/2023	

# Labeler - OPMX Chula Vista (029918743)

Establishment			
Name	Address	ID/FEI	Business Operations
Bajaj Medical, LLC		078774921	manufacture(69729-068), pack(69729-068), label(69729-068)

Revised: 4/2024 OPMX Chula Vista