

**COLD AND FLU HBP- acetaminophen, chlorpheniramine maleate tablet, film coated**  
**Kroger Company**

-----

**Kroger 44-692**

***Active ingredients (in each tablet)***

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg

***Purpose***

Pain reliever/fever reducer  
Antihistamine

***Uses***

- temporarily relieves:
  - headache
  - sneezing
  - runny nose
  - minor aches and pains
  - sore throat
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy

**Ask a doctor before use if the user has**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if the user is**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days (children 6 to under 12 years) or 10 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**

adults and children 12 years and over	2 tablets every 4 to 6 hours; do not take more than 12 tablets in 24 hours
children 6 to under 12	1 tablet every 4 to 6 hours; do not take more than 5

years	tablets in 24 hours
children under 6 years	do not use unless directed by a doctor

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, FD&C red #40 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

### ***Questions or comments?***

Call **1-800-632-6900**

### ***Principal display panel***

NDC 30142-698-09

**Kroger®**  
health

COMPARE TO THE ACTIVE INGREDIENTS  
IN CORICIDIN® HBP COLD & FLU\*

**Cold & Flu**  
**HBP**  
FOR  
PEOPLE WITH  
**HIGH BLOOD**  
**PRESSURE**

**Acetaminophen**  
Pain Reliever/Fever Reducer  
Chlorpheniramine Maleate  
Antihistamine

**RELIEVES: RUNNY NOSE,  
SNEEZING, BODY ACHES  
AND PAINS, FEVER,  
SORE THROAT  
DECONGESTANT-FREE**

ACTUAL  
SIZE



# COLD AND FLU HBP

acetaminophen, chlorpheniramine maleate tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30142-698
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C RED NO. 40 ALUMINUM LAKE</b> (UNII: 6T47AS764T)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;692
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-698-09	2 in 1 CARTON	04/14/2022	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/14/2022	

**Labeler** - Kroger Company (006999528)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(30142-698) , pack(30142-698)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(30142-698)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(30142-698)

Revised: 1/2026

Kroger Company