

**NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr,
doxylamine succinate liquid
Walgreens**

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

if a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is

accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be a signs of a serious conditions.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- **each 30 mL contains:** potassium 5 mg
- **each 30 mL contains:** sodium 24 mg
- store between 20-25°C (68-77°F). Do not refrigerate

Inactive ingredients

acesulfame potassium, alcohol, citric acid, FD&C blue1, FD&C red 40, Flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu††

NIGHTTIME

Cold & Flu

ACETAMINOPHEN 650 mg / PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr 30 mg / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE 12.5 mg / ANTIHISTAMINE

Multi-Symptom

- Relieves headache, fever, sore throat, minor aches & pains, sneezing, runny nose & cough
- For ages 12 years & over
- ALCOHOL 10%

Cherry flavor

FL OZ (mL)

††This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY:

WALGREEN CO.

200 WILMOT RD.,

DEERFIELD, IL 60015

Product Label

NDC 0363-9975-08

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 30 mL) **Purposes**
 Acetaminophen 650 mg..... Pain reliever/fever reducer
 Dextromethorphan HBr 30 mg..... Cough suppressant
 Doxylamine succinate 12.5 mg..... Antihistamine

Uses ■ temporarily relieves these common cold/flu symptoms
 ■ minor aches and pains ■ headache ■ sore throat
 ■ runny nose and sneezing ■ fever
 ■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks

← PEEL HERE

Drug Facts (continued under label)

Walgreens



Compare to the active ingredients in Vicks® NyQuil® Cold & Flu[†]

NIGHTTIME Cold & Flu

ACETAMINOPHEN 650 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 30 mg / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE 12.5 mg / ANTIHISTAMINE

Multi-Symptom

- Relieves headache, fever, sore throat, minor aches & pains, sneezing, runny nose & cough
- For ages 12 years & over
- ALCOHOL 10%



Cherry flavor

8 FL OZ (237 mL)

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY:
WALGREEN CO.
 200 WILMOT RD.,
 DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com

©2022 Walgreen Co.
 PLD-B427B LB009016

Discard Seal, Empty & Replace Cap



PLASTIC BOTTLE



PLASTIC CUP

W3ORG1022-F
 REV-0323

ITEM 620217 W11042-0423-F



3 11917 12395 0

how2recycle.info

Drug Facts (continued)
 after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 ■ liver disease ■ glaucoma
 ■ cough that occurs with too much phlegm (mucus)
 ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ trouble urinating due to enlarged prostate gland
Ask a doctor or pharmacist before use if you are
 ■ taking sedatives or tranquilizers
 ■ taking the blood thinning drug warfarin
When using this product
 ■ excitability may occur, especially in children
 ■ marked drowsiness may occur
 ■ avoid alcoholic drinks
 ■ be careful when driving a motor vehicle or operating machinery
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
Stop use and ask a doctor if
 ■ pain or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
 ■ do not take more than directed (see Overdose warning)
 ■ do not take more than 4 doses in any 24-hour period
 ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
 ■ keep dosing cup with product
 ■ adults and children 12 years and over: 30 mL every 6 hours
 ■ children under 12 years of age: do not use
 ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing
Other information
 ■ each 30 mL contains: potassium 5 mg
 ■ each 30 mL contains: sodium 24 mg
 ■ store between 20-25°C (68-77°F). Do not refrigerate.
Inactive ingredients: acetylsalicylic acid, alcohol, citric acid, FD&C blue #1, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate
Questions or comments?
 Call 1-877-753-3635 Monday-Friday 9AM-5PM EST

WALGREENS Nighttime Cold & Flu

NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9975
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9975-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2022	01/30/2027

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/30/2022	01/30/2027

Labeler - Walgreens (008965063)