

**ACETAMINOPHEN DIPHENHYDRAMINE HCL- acetaminophen diphenhydramine
hcl tablet
Walgreens**

**Pain Reliever PM
ACETAMINOPHEN USP, 500 mg / PAIN RELIEVER
DIPHENHYDRAMINE HCl USP, 25 mg / NIGHTTIME SLEEP AID
Extra Strength
Nighttime
Non-habit forming**

Active ingredients (in each caplet)

Acetaminophen USP, 500 mg

Diphenhydramine HCl USP, 25 mg

Purposes

Pain reliever

Nighttime sleep aid

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin

- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
- These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use

Keep out of reach of children

Keep out of reach of children

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ **do not take more than directed (see overdose warning)**

adults and children 12 years and over

■ take 2 caplets at bedtime

■ do not take more than 2 caplets of this product in 24 hours

children under 12 years

■ do not use

Other information

■ store between 20°-25°C (68°-77°F). See USP Controlled Room Temperature.

■ see end panel for lot number and expiration date

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, hypromellose, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone k-30, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

Principal Display panel

VALUE SIZE

Compare to the active ingredients
in Extra Strength Tylenol PM™



Pain Reliever PM

ACETAMINOPHEN USP, 500 mg / PAIN RELIEVER
DIPHENHYDRAMINE HCl USP, 25mg

- Nighttime
- Extra Strength
- Non-habit forming

250 CAPLETS* (*capsule shaped tablets)



ACTUAL SIZE

IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE
DANGER: EMERGENCY DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet)
Acetaminophen USP 500 mg
Diphenhydramine HCl USP 25mg

Purposes
Pain relief
Nighttime sleep aid

Uses
Temporary relief of occasional headache and minor aches and pains with accompanying discomforts

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you use more than 4,000 mg of acetaminophen in 24 hours.
Do not take more than 1 caplet every 4 hours and do not use this product for more than 10 days.
Do not take this product if you are taking any other medicine that contains acetaminophen or any other medicine that contains diphenhydramine.
If a pain reaction occurs, stop use and seek medical help right away.

Do not use
If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
If you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have
Liver disease ■ a breathing problem such as emphysema or chronic bronchitis
Trouble urinating due to an enlarged prostate gland ■ glaucoma

Ask a doctor or pharmacist before use if you are
Taking the blood thinning drug warfarin ■ taking medicines or tranquilizers
When using this product ■ drowsiness will occur ■ avoid alcoholic drinks
■ do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if
Drowsiness persists continuously for more than 2 weeks. Drowsiness may be a symptom of serious underlying medical illness. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: If value of overdose, get medical help or contact a Poison Control Center, night, phone, (1-800-222-1222). Dark medical attention is critical for adults, as well as for children even if you do not notice any signs or symptoms.

Directions Do not take more than directed (see overdose warning)
adults and children 12 years and over ■ 2 tablets at bedtime
■ do not take more than 2 caplets of this product in 24 hours
children under 12 years ■ do not use

Other information:
Contains no aspirin

Caution:
Keep out of reach of children.

Directions:
Do not take more than directed (see overdose warning)
adults and children 12 years and over ■ 2 tablets at bedtime
■ do not take more than 2 caplets of this product in 24 hours
children under 12 years ■ do not use

Other information:
Contains no aspirin

Caution:
Keep out of reach of children.

Directions:
Do not take more than directed (see overdose warning)
adults and children 12 years and over ■ 2 tablets at bedtime
■ do not take more than 2 caplets of this product in 24 hours
children under 12 years ■ do not use

Other information:
Contains no aspirin

Caution:
Keep out of reach of children.

Directions:
Do not take more than directed (see overdose warning)
adults and children 12 years and over ■ 2 tablets at bedtime
■ do not take more than 2 caplets of this product in 24 hours
children under 12 years ■ do not use

lot
exp

RECYCLE

W3ORQ224-F
W10761-0924-F

ITEM 625154

11917 04199 5

3

MADE IN INDIA

NDC 085-961-25

LOT

EXP

REV-0524

simulate a printed label,
fold along dotted line.

Inside (adhesive side)

Drug Facts (continued)

Other information:
Keep out of reach of children.

Directions:
Do not take more than directed (see overdose warning)
adults and children 12 years and over ■ 2 tablets at bedtime
■ do not take more than 2 caplets of this product in 24 hours
children under 12 years ■ do not use

Other information:
Contains no aspirin

Caution:
Keep out of reach of children.

Directions:
Do not take more than directed (see overdose warning)
adults and children 12 years and over ■ 2 tablets at bedtime
■ do not take more than 2 caplets of this product in 24 hours
children under 12 years ■ do not use

Other information:
Contains no aspirin

Caution:
Keep out of reach of children.

Directions:
Do not take more than directed (see overdose warning)
adults and children 12 years and over ■ 2 tablets at bedtime
■ do not take more than 2 caplets of this product in 24 hours
children under 12 years ■ do not use



ACETAMINOPHEN DIPHENHYDRAMINE HCL

acetaminophen diphenhydramine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9661
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE (UNII: 8GTS82S83M) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, CORN (UNII: O8232NY3S)	
POVIDONE K30 (UNII: U725QWY32X)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL (caplet-shaped)	Size	17mm
Flavor		Imprint Code	G651
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9661-15	150 in 1 BOTTLE; Type 0: Not a Combination Product	07/23/2024	
2	NDC:0363-9661-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	07/23/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/23/2024	

Labeler - Walgreens (008965063)

Revised: 7/2024

Walgreens