

COBURN FOAM-N-DIP- iodine liquid

Stearns Packaging Corporation

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

Iodine Teat Dip

IODOPHORS are considered relatively non-toxic, but should always be used in accordance with label directions. Consult your veterinarian in case of teat irritation.

TEAT DIPPING IN COLD WEATHER:

When the temperature falls below 10 degrees F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess
teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

PRODUCT INFORMATION: This teat dip is an IODOPHOR (a combination of iodine and complexing agents), and when properly used, is effective as an aid in reducing the spread of mastitis causing organisms.

DO NOT DILUTE. USE AT FULL STRENGTH. PROTECT FROM FREEZING.

Important Use Instructions

FOR PRE-MILKING

1. Wash each individual teat with a sanitizing udder wash.
- NOTE: For best results, keep hair around teat based clipped.
2. Hand strip teats to initial let down.
 3. Pre-dip entire teat with PRE-POST 1/2% on all four quarters. Repeat process on next 4 to 6 cows or until first cow prepped has at least 1 minute contact time.
 4. Wipe each individual teat with a clean paper towel. Utilize a "pulling down" motion which removes
iodine solution, germs, and sediment from teat skin. Very important!
 5. Place machine on cow and repeat wiping process on remaining cows that have been predipped.

FOR POST MILKING

DIRECTIONS FOR TEAT DIPPING. Dip entire teat with undiluted dip solution. Allow to air dry and remain on teats for prolonged contact time. Clean, sanitize and thoroughly dry each teat prior to next milking. Replace dip solution when visibly cloudy or dirty.

FIRST-AID:

EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes.

Consult a physician.

INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately. **24 HOUR EMERGENCY CONTACT: 1-800-255-3924**

**Coburn Foaming
Iodine Teat Dip**

FOAM-N-DIP

**NPE-Free Aid in Reducing the Spread of
Organisms Which May Cause Mastitis**

Active Ingredient.....0.5% Available Iodine
Emollient System Contains.....5% Glycerin
Minimum pH.....4.0

KEEP OUT OF THE REACH OF CHILDREN
See back panel for Precautionary Statements

The Coburn Company, Inc.
800 776 7024 | COBURN.COM

NET CONTENTS: 1 U.S. GALLON (3.78 L)



**NPE-Free Aid in Reducing the Spread of
Organisms Which May Cause Mastitis**

Active Ingredient0.5% Available Iodine
Emollient System Contains5% Glycerin
Minimum pH4.0

KEEP OUT OF THE REACH OF CHILDREN
See back panel for Precautionary Statements.

The Coburn Company, Inc.
800 776 7024 | COBURN.COM

NET CONTENTS: 1 U.S. GALLON (3.78 L)



DO NOT DILUTE. USE AT FULL STRENGTH. PROTECT FROM FREEZING.

Important Use Instructions

FOR PRE-MILKING

1. Wash each individual teat with a sanitizing udder wash.
NOTE: For best results, keep hair around teat base clipped.
2. Hand strip teats to initiate let down.
3. Pre-dip entire teat with FOAM-N-DIP on all four quarters. Repeat process on next 4 to 6 cows or until first cow prepped has at least 1 minute contact time.
4. Wipe each individual teat with a clean paper towel. Utilize a "pulling down" motion which removes iodine solution, germs, and sediment from teat skin. Very important!
5. Place machine on cow and repeat wiping process on remaining cows that have been prepped.

FOR POST MILKING

DIRECTIONS FOR TEAT DIPPING. Dip entire teat with undiluted dip solution. Allow to air dry and remain on teats for prolonged contact time. Clean, sanitize and thoroughly dry each teat prior to next milking. Replace dip solution when visibly cloudy or dirty.

PRODUCT INFORMATION. This teat dip is an IODOPHOR (a combination of iodine and complexing agents), and when properly used, is effective as an aid in reducing the spread of mastitis causing organisms. IODOPHORS are considered relatively non-toxic, but should always be used in accordance with label directions. Consult your veterinarian in case of teat irritation.

TEAT DIPPING IN COLD WEATHER:

When the temperature falls below 10°F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

FIRST-AID:

EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.
INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately. **24 HOUR EMERGENCY CONTACT: 1-800-255-3924**



COBURN FOAM-N-DIP

iodine liquid

Product Information

| | | | |
|--------------------------------|-----------------|---------------------------|----------------|
| Product Type | OTC ANIMAL DRUG | Item Code (Source) | NDC:60282-3106 |
| Route of Administration | TOPICAL | | |

| Active Ingredient/Active Moiety | | | | |
|--|--|-----------------------------|----------------------|--------------------|
| Ingredient Name | | | Basis of Strength | Strength |
| IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4) | | | IODINE | 0.5 kg in 100 L |
| | | | | |
| Inactive Ingredients | | | | |
| Ingredient Name | | | Strength | |
| GLYCERIN (UNII: PDC6A3C0OX) | | | 5 kg in 100 L | |
| | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:60282-3106-1 | 3.78 L in 1 BOTTLE, PLASTIC | | |
| | | | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | | Marketing Start Date | Marketing End Date |
| unapproved drug other | | | 02/07/2019 | |

Labeler - Stearns Packaging Corporation (006069256)

Registrant - Stearns Packaging Corporation (006069256)

| Establishment | | | |
|-------------------------------|---------|-----------|---------------------|
| Name | Address | ID/FEI | Business Operations |
| Stearns Packaging Corporation | | 006069256 | manufacture |

| Establishment | | | |
|---------------------------|---------|-----------|---------------------|
| Name | Address | ID/FEI | Business Operations |
| Lab Express International | | 800875106 | api manufacture |

Revised: 12/2025

Stearns Packaging Corporation