

**MOTION SICKNESS- dimenhydrinate tablet**  
**Geiss, Destin & Dunn Inc.**

-----  
**GoodSense 44-198**

***Active ingredient (in each tablet)***

Dimenhydrinate 50 mg

***Purpose***

Antiemetic

***Uses***

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

***Warnings***

**Do not use**

for children under 2 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

**Other information**

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

**Inactive ingredients**

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

**Questions or comments?**

**1-800-426-9391**

**Principal display panel**

**GoodSense®**

NDC 50804-198-02

*Original Formula*

**Motion Sickness**

*Dimenhydrinate 50 mg*

*Antiemetic*

***Prevents Nausea, Vomiting & Dizziness  
for Children & Adults***

**12** Tablets

*actual size*

**\*Compare to the** *active ingredient of*  
**Dramamine® Original Formula**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Medtech Products Inc.,  
owner of the registered trademark Dramamine® Original Formula.

50844 REV0518B19802

Distributed by: Perrigo Direct, Inc., Peachtree City, GA 30269

[www.PerrigoDirect.com](http://www.PerrigoDirect.com) (1-800-426-9391)

GoodSense® is a registered trademark of  
L. Perrigo Company.

GOODSENSE®

NDC 50804-198-02

# Motion Sickness

Dimenhydrinate 50 mg  
Antiemetic

12 Tablets

GOODSENSE®

# Motion Sickness

GOODSENSE®

NDC 50804-198-02

Original Formula

# Motion Sickness

Dimenhydrinate 50 mg  
Antiemetic

Prevents Nausea, Vomiting & Dizziness  
for Children & Adults



actual size

\*Compare to the active ingredient of  
Dramamine® Original Formula

12 Tablets

Z-Fold  
LEBG302104

\*This product is not manufactured or distributed by Medtech Products Inc.,  
owner of the registered trademark Dramamine® Original Formula.  
50844 REV0518B19802

Distributed by: Perrigo Direct, Inc., Peachtree City, GA 30269  
www.PerrigoDirect.com (1-800-426-9391)  
GoodSense® is a registered trademark of  
L. Perrigo Company.




8-0560-198-02-R  
REV0518B19802

no print / no varnish area  
lot no. & exp. date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING

<p><b>Warnings</b> Do not use for children under 2 years of age unless directed by a doctor. Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. When using this product ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	
<p><b>Other information</b> ■ each tablet contains: calcium 35 mg ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture ■ see end flap for expiration date and lot number <b>Inactive ingredients</b> croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid <b>Questions or comments?</b> 1-800-426-9391</p>	
<p>children 6 to under 12 years do not exceed 3 tablets in 24 hours, or as directed by a doctor</p>	<p>children 2 to under 6 years ½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor</p>

GOODSENSE®  
**Motion Sickness**  
Dimenhydrinate 50 mg  
Antiemetic



**Good Sense 44-198**

**MOTION SICKNESS**

dimenhydrinate tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50804-198
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DIMENHYDRINATE</b> (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	44;198
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50804-198-02	2 in 1 CARTON	04/07/2020	

<b>1</b>	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M009	04/07/2020	

**Labeler** - Geiss, Destin & Dunn Inc. (076059836)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		038154464	pack(50804-198)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	manufacture(50804-198) , pack(50804-198)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(50804-198)

Revised: 11/2025

Geiss, Destin & Dunn Inc.