

**NIGHTTIME COLD AND FLU RELIEF- acetaminophen dextromethorphan hbr
doxylamine succinate capsule, liquid filled
QUALITY CHOICE (Chain Drug Marketing Association)**

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/Fever Reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists more than 2 days, is

accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as

for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call **1-800-935-2362** Monday-Friday 9AM-5PM EST

Principal Display Panel

†Compare to the Active Ingredients in VICKS® NYQUIL™ COLD & FLU LIQUICAPS™

NightTime

Cold & Flu

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

Pain Reliever /Fever Reducer

Cough Suppressant | Antihistamine

Alcohol Free

Softgels

†This product is not manufactured or distributed by the Procter & Gamble Company. Vicks®, NyQuil™, and LiquiCaps™ are registered trademark of the Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
---	----------------------	---------

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE (oblong)	Size	21mm
Flavor		Imprint Code	PC10
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-148-24	24 in 1 CARTON	07/31/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/31/2024	

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)