

**NASAL- oxymetazoline hydrochloride spray**  
**Meijer, Inc.**

-----  
**Meijer Distribution, Inc. Nasal Spray Drug Facts**

**Active ingredient**

Oxymetazoline hydrochloride 0.05%

**Purpose**

Nasal decongestant

**Uses**

- temporarily relieves nasal congestion due to:
- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

**Warnings**

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product**

- **do not use more than directed**
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

**Stop use and ask a doctor if**

symptoms persist

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

**To Use:** Shake well before use. Hold white tabs, SQUEEZE grooved area of cap FIRMLY and turn counter clockwise. Before using the first time, prime metered pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use.

### **Other information**

- store at 20-25°C (68-77°F)
- retain carton for future reference on full labeling

### **Inactive ingredients**

benzalkonium chloride solution, benzyl alcohol, camphor, dibasic sodium phosphate, edetate disodium, eucalyptol, menthol, microcrystalline cellulose and carboxymethylcellulose sodium, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

meijer®

COMPARE TO AFRIN® NO DRIP ACTIVE INGREDIENT

MAXIMUM STRENGTH

Nasal Spray

Oxymetazoline Hydrochloride

12 HOUR NASAL DECONGESTANT

NO DRIP

With Cooling Menthol

Steroid Free

For Colds & Allergies

1 FL OZ (30 mL)

**meijer.** NDC 79481-0425-0

COMPARE TO AFRIN® NO DRIP ACTIVE INGREDIENT\*

**MAXIMUM STRENGTH**

**Nasal Spray**

Oxymetazoline Hydrochloride

**12 HOUR NASAL DECONGESTANT** **NO DRIP**

With Cooling Menthol

Steroid Free

For Colds & Allergies

**1 FL OZ (30 mL)**

**meijer.** NDC 79481-0425-0

**MAXIMUM STRENGTH**

**Nasal Spray**

Oxymetazoline Hydrochloride

**12 HOUR NASAL DECONGESTANT**

**NO DRIP**

With Cooling Menthol

1 FL OZ (30 mL)

Actual Size

**Drug Facts**

**Active ingredient** Oxymetazoline hydrochloride 0.05%...Nasal decongestant

**Purpose**

**Uses**

- temporarily relieves nasal congestion due to:
  - common cold
  - hay fever
  - upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

**Warnings**

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product**

- do not use more than directed
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

**To Use:** Shake well before use. Hold white tabs, **SQUEEZE** grooved area of cap **FIRMLY** and turn counter clockwise. Before using the first time, prime pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use.

**Other information**

- store at 20-25°C (68-77°F)
- retain carton for future reference on full labeling

**Drug Facts (continued)**

**Inactive ingredients** benzalkonium chloride solution, benzyl alcohol, camphor, dibasic sodium phosphate, edetate disodium, eucalyptol, menthol, microcrystalline cellulose and carboxymethylcellulose sodium, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water

**Questions or comments?** 1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Afrin®.

**How to use:** Hold white tabs, **SQUEEZE** grooved area of cap **FIRMLY** and turn counter clockwise. Secure cap after use.

**SQUEEZE** grooves **FIRMLY** while turning

Hold white tabs **FIRMLY**

7 19283 23093 5

7J810 6E C1

# NASAL

oxymetazoline hydrochloride spray

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-0425
<b>Route of Administration</b>	NASAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OXYMETAZOLINE HYDROCHLORIDE</b> (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	.05 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM</b> (UNII: GR686LBA74)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>EUCALYPTOL</b> (UNII: RV6J6604TK)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Product Characteristics

<b>Color</b>	WHITE (off white)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0425-0	1 in 1 CARTON	06/30/2025	
1		30 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	06/30/2025	

**Labeler** - Meijer, Inc. (006959555)

Revised: 7/2025

Meijer, Inc.