

OMEPRAZOLE MAGNESIUM- omeprazole magnesium capsule, delayed release

Chain Drug Marketing Association

Active ingredient (in each capsule)

*Omeprazole delayed-release capsules 20 mg
(equivalent to 20.6 mg omeprazole magnesium)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs ***2 or more*** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to omeprazole
- Omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if

you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush capsules.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients

black iron oxide, dibasic calcium phosphate, gelatin, glyceryl monostearate, hypromellose 3 cps, magnesium oxide, magnesium stearate, methacrylic acid copolymer dispersion, methacrylic acid copolymer Type B, microcrystalline cellulose, polysorbate 80, potassium hydroxide, propylene glycol, red iron oxide, shellac, silicon dioxide,

sodium lauryl sulphate, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions

call **1-888-375-3784**

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Carton Label



Drug Facts (continued)

- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months.
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush capsules. ▶



Drug Facts (continued)

- Repeated 14-Day Courses (if needed)**
- you may repeat a 14-day course every 4 months
 - do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 - children under 18 years of age; ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20 to 25°C (68 to 77°F) and protect from moisture

Inactive ingredients

black iron oxide, dibasic calcium phosphate, gelatin, glyceryl monostearate, hypromellose 3 cps, magnesium oxide, magnesium stearate, methacrylic acid copolymer dispersion, methacrylic acid copolymer Type B, microcrystalline cellulose, polysorbate 80, potassium hydroxide, propylene glycol, red iron oxide, shellac, silicic acid, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments?
call 1-888-375-3784

**This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®. Prilosec OTC® is a registered trademark of AstraZeneca AB.

Made in INDIA

REV 01/23

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

TAMPER EVIDENT: Do not use if roll seal under cap printed with "sealed for your protection" or pink band around center of capsule is broken or missing.
IMPORTANT: Read the directions and warnings before use. Keep the carton. It contains important information.

NDC 63868-177-15



"Compare to the Active Ingredient in Prilosec OTC®"

Treats Frequent Heartburn
See Current Drug Facts

Omeprazole
Delayed-Release Capsules, 20 mg*

Frequent Heartburn Relief

24 HR Acid Reducer



14 Capsules
(Safety Sealed)
One 14-Day Course of Treatment
May Take 1 to 4 Days for Full Effect



Distributed by CDMA, Inc.
Norwalk, CT 06855
www.qualbychoice.com
Questions: 800-925-2262



LOT
EXP

Drug Facts

Active ingredient (in each capsule) Purpose
*Omeprazole delayed-release capsule 20 mg (equivalent to 20.6 mg omeprazole magnesium).....Acid reducer

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert:
■ Do not use if you are allergic to omeprazole
■ Omeprazole may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition. ▶

Bottle Label

NDC 63868-177-15



Treats **Frequent Heartburn**

Omeprazole

Delayed-Release Capsules
20 mg*

Frequent Heartburn Relief

24 HR Acid Reducer

14 Capsules (Safety Sealed)
One 14-Day Course of Treatment
May Take 1 to 4 Days for Full Effect

IMPORTANT: This label does not contain full product information. See carton for complete information. Read all warnings and directions on carton before use. Retain carton for reference.

Active ingredient Purpose (in each capsule)
*Omeprazole delayed-release capsule 20 mg Acid omeprazole magnesium).....reducer

Use
■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Other Information
■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) and protect from moisture

Questions? call 1-888-375-3784

TAMPER EVIDENT: Do not use if foil seal under cap printed with "Sealed for your protection" or pink band around center of capsule is broken or missing.

Distributed by CDMA, Inc.
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

Made in India
REV01/23

150092444

LOT
EXP

OMEPRAZOLE MAGNESIUM

omeprazole magnesium capsule, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-177(NDC:55111-397)
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OMEPRAZOLE MAGNESIUM (UNII: 426QFE7XLK) (omeprazole - UNII:KG60484QX9)	omeprazole	20 mg	

Inactive Ingredients		
Ingredient Name		Strength
FERROSFERRIC OXIDE (UNII: XM0M87F357)		
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)		
Gelatin (UNII: 2G86QN327L)		
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)		
HYPROMELLOSE 2208 (3 MPA.S) (UNII: 9H4L9160BU)		
Magnesium Oxide (UNII: 3A3U0GI71G)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2) (UNII: 5KY68S2577)		
Methacrylic Acid - Ethyl Acrylate Copolymer (1:1) Type A (UNII: NX76LV5T8J)		
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)		
Polysorbate 80 (UNII: 6OZP39ZG8H)		
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
FERRIC OXIDE RED (UNII: 1K09F3G675)		
SHELLAC (UNII: 46N107B71O)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
RAW SUGAR (UNII: 8M707QY5GH)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)		

Sodium Lauryl Sulfate (UNII: 368GB5141J)

Product Characteristics

Color	WHITE, PINK	Score	no score
Shape	CAPSULE	Size	22mm
Flavor		Imprint Code	OMP20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-177-15	1 in 1 CARTON	01/01/2016	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:63868-177-42	3 in 1 CARTON	01/01/2016	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078878	01/01/2016	

Labeler - Chain Drug Marketing Association (011920774)

Revised: 7/2025

Chain Drug Marketing Association