# STOMACH RELIEF- bismuth subsalicylate tablet CDMA- QC

-----

#### **Stomach Relief**

#### Active ingredient(s)

Bismuth subsalicylate 262mg (each tablet contains 102mg salicylate)

## **Purpose**

Antidiarrheal/Antacid

#### Use(s)

- relieves
- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

# **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like

symptoms should not use this product. When using this product, if changes in behavior with nausea

and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's

syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### Do not use

- an ulcer
- a bleeding problem
- bloody or black stool

#### Ask a doctor before use if

fever

• mucus in the stool

#### Ask a doctor or pharmacist before use if

you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

#### When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

#### Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

#### Pregnancy/Breastfeeding

If pregnant or breast-feeding, ask a health professional before use.

#### Keep out of reach of children

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### **Directions**

Chew or dissolve in mouth

# Adults and children 12 years and over:

- 2 tablets (1 dose) every ½ hour or 4 tablets (2 doses) every hour as needed for diarrhea
- 2 tablets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days

# Children under 12 years of age: ask a doctor

drink plenty of clear fluids to help prevent dehydration caused by diarrhea

#### Other information

Phenylketonurics: contains phenylalanine 1.1mg per tablet Calcium content per tablet: 73mg

Do not use if cellophane unit is torn

#### Storage

Store at room temperature 15°- 30°C (59°- 86°F)

## **Inactive ingredients**

acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrates, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavor, silicon dioxide

#### **Principal Display Panel**



#### Stomach Relief

STOMACH RELIEF			
bismuth subsalicylate tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-622
Route of Administration	ORAL		

# Active Ingredient/Active Moiety Ingredient Name BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII: ZS9CD1I8YE) BISMUTH SUBSALICYLATE BISMUTH SUBSALICYLATE 262 mg

Inactive Ingredients			
Ingredient Name	Strength		
ASPARTAME (UNII: Z0H242BBR1)			
ACACIA (UNII: 5C5403N26O)			
CALCIUM CARBONATE (UNII: H0G9379FGK)			
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)			
DEXTRATES (UNII: G263MI44RU)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
PEPPERMINT (UNII: V95R5KMY2B)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			

Product Characteristics				
Color	pink	Score	no score	
Shape	ROUND	Size	17mm	
Flavor		Imprint Code	RH046	
Contains				

ı	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:63868-622- 30	30 in 1 CELLO PACK; Type 0: Not a Combination Product	06/16/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	12/23/2014	

# **Labeler - CDMA- QC (011920774)**

Revised: 6/2021 CDMA- QC