# HYDROCORTISONE ACETATE- hydrocortisone acetate suppository Golden State Medical Supply, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

## **Hydrocortisone Acetate**

Hydrocortisone Acetate Suppositories 25mg Rx Only

For Rectal Administration

#### **DESCRIPTION:**

Hydrocortisone acetate is a corticosteroid designed chemically as pregn-4-ene 3, 20-dione, 21-(acetyloxy)-11, 17-dihydroxy--(11  $\beta$ ) with the following structural formula:

Each rectal suppository contains hydrocortisone acetate, USP 25 mg in a specially blended hydrogenated vegetable oil base.

#### **CLINICAL PHARMACOLOGY:**

In normal subjects, about 26% of hydrocortisone acetate is absorbed when the suppository is applied to the rectum. Absorption of hydrocortisone acetate may vary across abraded or inflamed surfaces. Topical steroids are primarily effective because of their anti-inflammatory, anti-pruritic and vasoconstrictive action.

#### **INDICATIONS AND USAGE:**

Hydrocortisone Acetate suppositories are indicated for use in inflamed hemorrhoids, post-irradiation (factitial) proctitis; as an adjunct in the treatment of chronic ulcerative colitis; cryptitis; and other inflammatory conditions of anorectum and pruritus ani.

#### **CONTRAINDICATIONS:**

Hydrocortisone Acetate suppositories are contraindicated in those patients having a history of hypersensitivity to hydrocortisone acetate or any of the components.

#### **PRECAUTIONS:**

Do not use Hydrocortisone Acetate suppositories unless adequate proctologic examination is made. If irritation develops, the product should be discontinued and appropriate therapy instituted. In the presence of an infection, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, Hydrocortisone Acetate should be discontinued until the infection has been adequately controlled.

# Carcinogenesis:

No long term studies in animals have been performed to evaluate the carcinogenic potential of corticosteroid suppositories.

## **Pregnancy Category C:**

In laboratory animals, topical steroids have been associated with an increase in the incidence of fetal abnormalities when gestating females have been exposed to rather low dosage levels. There are no adequate and well controlled studies in pregnant women. Hydrocortisone Acetate suppositories should only be used during pregnancy if the potential benefit justifies the risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

It is not known whether this drug is excreted in human milk and because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from Hydrocortisone Acetate suppositories, a decision should be

made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

#### **ADVERSE REACTIONS:**

The following local adverse reactions have been reported with corticosteroid suppositories; burning, itching, irritation, dryness, folliculitis, hypopigmentation, allergic contact dermatitis, secondary infection.

#### **DRUG ABUSE AND DEPENDENCE:**

Drug abuse and dependence have not been reported in patients treated with Hydrocortisone Acetate suppositories.

#### **OVERDOSAGE:**

If signs and symptoms of systemic overdosage occur, discontinue use.

#### **DOSAGE AND ADMINISTRATION:**

For rectal administration. Detach one suppository from strip of suppositories. Remove the wrapper. Avoid excessive handling of the suppository which is designed to melt at body temperature. Insert suppository into the rectum with gentle pressure, pointed end first. Insert one suppository in the rectum twice daily, morning and night for two weeks, in nonspecific proctitis. In more severe cases, one suppository three times a day or two suppositories twice daily. In factitial proctitis, the recommended duration of therapy is six to eight weeks or less, according to the response of the individual case.

All prescription substitutions and/or recommendations using this product shall be made subject to state and federal statutes as applicable. PleaseNOTE: This is notanOrange Book product and hasnot been subjected to FDA therapeutic or other equivalencytesting. No representation is made asto generic status orbioequivalency. Each person recommending a prescription substitution using this product shall make such recommendation based on his/her professional knowledge and opinion, upon evaluating the active ingredients, inactive ingredients, excipients and chemical information provided herein.

#### **HOW SUPPLIED:**

Hydrocortisone Acetate suppositories, 25mg are off-white, smooth surfaced and bullet shaped with one pointed end.

Box of 12 and 24 suppositories, NDC 51407-665-12 and NDC 51407-665-24.

#### STORAGE:

Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Excursions permitted to 15°-30°C (59°-86°F). Store away from heat. Protect from freezing.

Manufactured for:

Laser Pharmaceuticals, LLC

1015 Nine North Drive, Ste 400


Alpharetta, GA 30004 770-754-9846

Rev. 08/24

Marketed by:

GSMS, Inc.

Camarillo, CA 93012 USA



# **HYDROCORTISONE ACETATE**

hydrocortisone acetate suppository

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:51407-665
Route of Administration	RECTAL		
Active Ingredient/Active	Majatur		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Stre	ength Strength

HYDROCORTISONE ACETATE

25 mg

Inactive	<b>Ingredients</b>

Ingredient Name	Strength
-----------------	----------

HYDROGENATED PALM KERNEL OIL (UNII: FM8D1RE2VP)

Product Characteristics			
Color	yellow (off-white)	Score	
Shape	BULLET	Size	
Flavor		Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51407-665- 12	2 in 1 BOX	07/21/2022	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:51407-665- 24	4 in 1 BOX	07/21/2022	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/12/2013	

# Labeler - Golden State Medical Supply, Inc. (603184490)

Establishment				
Name	Address	ID/FEI	<b>Business Operations</b>	
Speciality Pharma Manufacturing, LLC		013957125	manufacture(51407-665)	

Revised: 2/2025 Golden State Medical Supply, Inc.