

**BENZONATATE- benzonatate capsule**  
**Preferred Pharmaceuticals, Inc.**

-----  
**BENZONATATE CAPSULES**

**100 mg Capsules**


**150 mg Capsules**

**200 mg Capsules**

**(benzonatate, USP)**

**DESCRIPTION**

Benzonatate, a non-narcotic oral antitussive agent, is 2, 5, 8, 11, 14, 17, 20, 23, 26-nonaooctacosan-28-yl p-(butylamino) benzoate; with a molecular weight of 603.7.



Each benzonatate capsule USP contains:

Benzonatate, USP 100 mg

Each benzonatate capsule USP contains:

Benzonatate, USP 150 mg

Each benzonatate capsule USP contains:

Benzonatate, USP 200 mg

Benzonatate capsules USP also contain: bloom gelatin, glycerin, purified water, medium chain triglycerides, lecithin, isopropyl alcohol, nitrogen.

100 mg and 200 mg capsules also contain D&C Yellow No. 10.

150 mg capsules also contain FD&C Yellow No. 6 powder.

Each capsule also contains black iron oxide, propylene glycol, hypromellose as imprinting ink.

**CLINICAL PHARMACOLOGY**

Benzonatate capsules acts peripherally by anesthetizing the stretch receptors located in the respiratory passages, lungs, and pleura by dampening their activity and thereby

reducing the cough reflex at its source. It begins to act within 15 to 20 minutes and its effect lasts for 3 to 8 hours. Benzonatate capsules has no inhibitory effect on the respiratory center in recommended dosage.

## **INDICATIONS AND USAGE**

Benzonatate capsules is indicated for the symptomatic relief of cough.

## **CONTRAINDICATIONS**

Hypersensitivity to benzonatate or related compounds.

## **WARNINGS**

### **Hypersensitivity**

Severe hypersensitivity reactions (including bronchospasm, laryngospasm and cardiovascular collapse) have been reported which are possibly related to local anesthesia from sucking or chewing the capsule instead of swallowing it. Severe reactions have required intervention with vasopressor agents and supportive measures.

### **Psychiatric Effects**

Isolated instances of bizarre behavior, including mental confusion and visual hallucinations, have also been reported in patients taking benzonatate capsules in combination with other prescribed drugs.

### **Accidental Ingestion and Death in Children**

Keep benzonatate capsules out of reach of children. Accidental ingestion of benzonatate capsules resulting in death has been reported in children below age 10. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. If accidental ingestion occurs, seek medical attention immediately (see OVERDOSAGE).

## **PRECAUTIONS**

Benzonatate is chemically related to anesthetic agents of the para-amino-benzoic acid class (e.g. procaine; tetracaine) and has been associated with adverse CNS effects possibly related to a prior sensitivity to related agents or interaction with concomitant medication.

### **Information for patients**

Swallow benzonatate capsules whole. Do not break, chew, dissolve, cut, or crush benzonatate capsules. Release of benzonatate from the capsule in the mouth can produce a temporary local anesthesia of the oral mucosa and choking could occur. If numbness or tingling of the tongue, mouth, throat, or face occurs, refrain from oral ingestion of food or liquids until the numbness has resolved. If the symptoms worsen or persist, seek medical attention.

Keep benzonatate capsules out of reach of children. Accidental ingestion resulting in

death has been reported in children. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. Signs and symptoms may include restlessness, tremors, convulsions, coma and cardiac arrest. If accidental ingestion occurs, seek medical attention immediately.

Overdosage resulting in death may occur in adults.

Do not exceed a single dose of 200 mg and a total daily dosage of 600 mg. If you miss a dose of benzonatate capsules, skip that dose and take the next dose at the next scheduled time. Do not take 2 doses of benzonatate capsules at one time.

### **Usage in Pregnancy**

Pregnancy Category C

Animal reproduction studies have not been conducted with benzonatate capsules. It is also not known whether benzonatate capsules can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Benzonatate capsules should be given to a pregnant woman only if clearly needed.

### **Nursing mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk caution should be exercised when benzonatate capsules is administered to a nursing woman.

### **Carcinogenesis, mutagenesis, impairment of fertility**

Carcinogenicity, mutagenicity, and reproduction studies have not been conducted with benzonatate capsules.

### **Pediatric Use**

Safety and effectiveness in children below the age of 10 have not been established. Accidental ingestion resulting in death has been reported in children below age 10. Keep out of reach of children.

## **ADVERSE REACTIONS**

Potential Adverse Reactions to benzonatate capsules may include:

**Hypersensitivity reactions** including bronchospasm, laryngospasm, cardiovascular collapse possibly related to local anesthesia from chewing or sucking the capsule.

**CNS:** sedation; headache; dizziness; mental confusion; visual hallucinations.

**GI:** constipation; nausea; GI upset.

**Dermatologic:** pruritus; skin eruptions.

**Other:** nasal congestion; sensation of burning in the eyes; vague “chilly” sensation; numbness of the chest; hypersensitivity.

Deliberate or accidental overdose has resulted in death, particularly in children.

## **OVERDOSAGE**

Intentional and unintentional overdose may result in death, particularly in children.

The drug is chemically related to tetracaine and other topical anesthetics and shares various aspects of their pharmacology and toxicology. Drugs of this type are generally well absorbed after ingestion.

### **Signs and Symptoms**

The signs and symptoms of overdose of benzonatate have been reported within 15-20 minutes. If capsules are chewed or dissolved in the mouth, oropharyngeal anesthesia will develop rapidly, which may cause choking and airway compromise.

CNS stimulation may cause restlessness and tremors which may proceed to clonic convulsions followed by profound CNS depression. Convulsions, coma, cerebral edema and cardiac arrest leading to death have been reported within 1 hour of ingestion.

### **Treatment:**

In case of overdose, seek medical attention immediately. Evacuate gastric contents and administer copious amounts of activated charcoal slurry. Even in the conscious patient, cough and gag reflexes may be so depressed as to necessitate special attention to protection against aspiration of gastric contents and orally administered materials. Convulsions should be treated with a short-acting barbiturate given intravenously and carefully titrated for the smallest effective dosage. Intensive support of respiration and cardiovascular-renal function is an essential feature of the treatment of severe intoxication from overdosage.

Do not use CNS stimulants.

### **DOSAGE AND ADMINISTRATION**

Adults and Children over 10 years of age: Usual dose is one 100 mg, 150 mg or 200 mg capsule three times a day as needed for cough. If necessary to control cough, up to 600 mg daily in three divided doses may be given. **Benzonatate capsules should be swallowed whole.** Benzonatate capsules are not to be broken, chewed, dissolved, cut or crushed.

### **HOW SUPPLIED**

Benzonatate Capsules, USP are available in 100 mg, 150 mg and 200 mg dosage strengths.

The 200 mg capsules are yellow round capsules containing clear to light yellow liquid, printed with '3' sign in black ink.

NDC 68788-8336-1 bottles of 15 capsules

NDC 68788-8336-2 bottles of 20 capsules

NDC 68788-8336-3 bottles of 30 capsules

NDC 68788-8336-6 bottles of 60 capsules

**Store at 20°C to 25°C (68°F to 77°F) [see USP Controlled Room Temperature].**

Protect from light. Dispense in tight, light-resistant container as defined in the USP with a

child-resistant closure.

Manufactured by:

Ascent Pharmaceuticals, Inc.

Central Islip, NY 11722

Manufactured for:

Camber Pharmaceuticals, Inc.

Piscataway, NJ 08854

**Repackaged By: Referred Pharmaceuticals Inc.**

Rev: 07/21

**Benzonatate Capsules 200 mg**


**NDC:68788-8336**

**Benzonatate Capsules, USP 200mg**

Generic for Tessalon Perles  
Each soft gelatin capsule contains:  
Benzonatate, USP...200mg

**Pkg Size:** Exp Date:  
Lot#: Batch#: Ins:  
Mfg: Camber Pharmaceuticals, Inc.  
Prod#: Warning

Store at 20°- 25°C (68°- 77°). See USP controlled room temperature. Protect from light. Keep this and all medication out of the reach of children. RX Only. Capsule is round, yellow, and imprinted with 3




CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed

Benzonatate Capsules, USP 200mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):

Benzonatate Capsules, USP 200mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):


Benzonatate Capsules, USP 200mg  
Qty:  
Insurance NDC:  
Lot#: Bat#:  
Prod# (NDC):

Benzonatate Capsules, USP 200mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):



Directions English

Take \_\_\_ capsule(s)  
every \_\_\_ hour(s).



Instrucciones Espanol:

Toma \_\_\_ capsula(s)  
cada \_\_\_ horas.

Log  
Chart  
Billing  
Patient

| <b>BENZONATATE</b>  |                         |                           |                               |
|---|-------------------------|---------------------------|-------------------------------|
| benzonatate capsule   |                         |                           |                               |
| Product Information   |                         |                           |                               |
| <b>Product Type</b>   | HUMAN PRESCRIPTION DRUG | <b>Item Code (Source)</b> | NDC:68788-8336(NDC:31722-958) |
| <b>Route of Administration</b>  | ORAL                    |                           |                               |
| Active Ingredient/Active Moiety                                       |                         |                           |                               |
| <b>Ingredient Name</b>  |                         | <b>Basis of Strength</b>  | <b>Strength</b>               |
| <b>BENZONATATE</b> (UNII: 5P4DHS6ENR) (BENZONATATE - UNII:5P4DHS6ENR) |                         | BENZONATATE               | 200 mg                        |
| Inactive Ingredients  |                         |                           |                               |

| Ingredient Name                                      | Strength |
|--|----------|
| <b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)       |          |
| <b>WATER</b> (UNII: 059QF0KO0R)                      |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                   |          |
| <b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U) |          |
| <b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)          |          |
| <b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)          |          |
| <b>NITROGEN</b> (UNII: N762921K75)                   |          |
| <b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)      |          |
| <b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)         |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)           |          |
| <b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)  |          |

### Product Characteristics

|                 |        |                     |          |
|-----------------|--------|---------------------|----------|
| <b>Color</b>    | yellow | <b>Score</b>        | no score |
| <b>Shape</b>    | ROUND  | <b>Size</b>         | 8mm      |
| <b>Flavor</b>   |        | <b>Imprint Code</b> | 3        |
| <b>Contains</b> |        |                     |          |

### Packaging

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:68788-8336-1 | 15 in 1 BOTTLE; Type 0: Not a Combination Product | 04/06/2023           |                    |
| 2 | NDC:68788-8336-2 | 20 in 1 BOTTLE; Type 0: Not a Combination Product | 04/06/2023           |                    |
| 3 | NDC:68788-8336-3 | 30 in 1 BOTTLE; Type 0: Not a Combination Product | 04/06/2023           |                    |
| 4 | NDC:68788-8336-6 | 60 in 1 BOTTLE; Type 0: Not a Combination Product | 04/06/2023           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA211518                               | 04/06/2023           |                    |

**Labeler** - Preferred Pharmaceuticals, Inc. (791119022)

**Registrant** - Preferred Pharmaceuticals, Inc. (791119022)

### Establishment

| Name                            | Address | ID/FEI    | Business Operations |
|---------------------------------|---------|-----------|---------------------|
| Preferred Pharmaceuticals, Inc. |         | 791119022 | REPACK(68788-8336)  |