

**DEXCHLORPHENIRAMINE MALEATE, PHENYLEPHRINE HCL-**  
**dexchlorpheniramine maleate, phenylephrine hcl tablet**  
**Syntho Pharmaceuticals Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**RYMED TABLETS**

**Active Ingredients (in each tablet)**

1. Dexchlorpheniramine Maleate - 2 mg
2. Phenylephrine HCl - 10 mg

**Inactive ingredients**

Lactose Monohydrate, Microcrystalline Cellulose, Magnesium Stearate, Purified Water, PEG, HPMC

**Purpose**

1. Dexchlorpheniramine Maleate 2 mg - Antihistamine
2. Phenylephrine HCl 10 mg - Nasal Decongestant

**Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

- **Do not exceed recommended dosage.**

**Do not use this product**

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**

**When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by a fever
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions**

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor
Children under 6 years of age:	Consult a doctor.

**Questions or Comments?**

Call 1-800-664-1490

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

# Rymed Tablets - NDC 66576-080-01 - 100's Bottle Label

Exp. Date:  
Lot:

7  
100801  
66576  
3 N



**Syntho**  
NDC 66576-080-01  
**RYMED TABLETS**  
Antihistamine • Nasal Decongestant

Each tablet contains:  
Dexchlorpheniramine Maleate ..... 2 mg  
Phenylephrine HCl ..... 10 mg

Store at 59°-86°F (15°-30°C) (see USP Controlled Room Temperature).  
Temperatures above 30°C may result in degradation.  
Do not use if foil seal is broken or missing.

Manufactured for:  
**EDWARDS PHARMACEUTICALS**  
100 tablets

**Drug Facts**

**Active Ingredients Purpose (in each tablet)**

Dexchlorpheniramine Maleate 2 mg ..... Antihistamine

Phenylephrine HCl 10 mg ..... Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes
- nasal congestion ■ reduces swelling of nasal passages

**BACK OF FRONT**

**Drug Facts (continued)**

**Warnings**

- Do not exceed recommended dosage.

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma ■ heart disease
- high blood pressure ■ thyroid disease ■ diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**

**When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by a fever
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions**

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor
Children under 6 years of age:	Consult a doctor.

**Inactive ingredients**

Lactose Monohydrate, Microcrystalline Cellulose, Magnesium Stearate, Purified Water, PEG, HPMC

**Questions or Comments?**  
Call 1-800-664-1490

Rev. 9/22

## DEXCHLORPHENIRAMINE MALEATE, PHENYLEPHRINE HCL

dexchlorpheniramine maleate, phenylephrine hcl tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66576-080
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	10 mg
<b>DEXCHLORPHENIRAMINE MALEATE</b> (UNII: B10YD955QW) (DEXCHLORPHENIRAMINE - UNII:3Q9Q0B929N)	DEXCHLORPHENIRAMINE MALEATE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	

**MICROCRYSTALLINE CELLULOSE** (UNII: OP1R32D61U)

**MAGNESIUM STEARATE** (UNII: 70097M6I3O)

**WATER** (UNII: 059QF0KO0R)

**POLYETHYLENE GLYCOL** (UNII: 3WJQ0SDW1A)

**HYPROMELLOSE, UNSPECIFIED** (UNII: 3NXW29V3WO)

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	ED;DM
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66576-080-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/02/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/02/2022	

**Labeler** - Syntho Pharmaceuticals Inc. (088797407)

**Registrant** - Syntho Pharmaceuticals Inc. (088797407)

### Establishment

Name	Address	ID/FEI	Business Operations
Syntho Pharmaceuticals Inc.		088797407	manufacture(66576-080)

Revised: 11/2022

Syntho Pharmaceuticals Inc.