NRG TOPICAL ANESTHETIC- benzocaine gel IQ Dental Supply, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use. Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59° - 86°F (15°-30°C). Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions of comments?

973-488-7966 or www.igdentalsupply.com



Distributed by IQ Dental

Topical Anesthetic Gel

Net Content: 1 oz (30 g)



Visit our website at: www.iqdentalsupply.com

353 Rt 46 W, Building C Unit 120 Fairfield, NJ 07004 Telephone (973) 488-7966

99290451 Rev 06/2018



mouth symptoms do not improve directed by a physician. If sore in 7 days; irritation, pain, or

Benzocaine 200 mg (in each g)

Active ingredients

Drug Facts

f swelling, rash or fever develops, in children under 2 years of age redness persists or worsens; or see your physician promptly.

For oral mucosal use only, as

Oral Anesthetic

Purpose

temporary relief of pain due to minor dental procedures.

directed by dentist. For the

When using this product Avoid contact with eves. If

Do not exceed recommended contact occurs, flush with water. dosage. If more than used for

Methemoglobinemia

Warnings

may cause methemoglobinemia, a serious condition that must be occur even if you have used this reduces the amount of oxygen warning: Use of this product carried in the blood. This can treated promptly because it

f pregnant or breast feeding,

Keep out of reach of children

ask a physician before use.

product before. Stop use and

seek immediate medical

· Apply only amount needed to

Directions

the oral mucosa to prevent or

relieve pain.

children under 2 years of age

do not use

Store at 59° -86° F (15 - 30°C).

Protect from freezing.

Other information

Poison Control Center right away.

pain is accidentally swallowed,

get medical help or contact a

attention if you or a child in your care develops: (cyanosis)

 pale, gray or blue colored skin headache

 dizziness or lightheadedness shortness of breath rapid heart rate

fatigue or lack of energy

Allergy Alert: Do not use on allergies to local anesthetics such as procaine, butacaine, patients with a history of

Do not use anesthetics.

benzocaine or other "caine"

additive.

(tartrazine), yellow #6, as a color #3, red #28, red #40, yellow #5,

blue #1, green #3, green #5, red

sodium saccharin. May contain

flavoring, PEG 3350, PEG 400,

nactive ingredients

Questions or comments? www.iqdentalsupply.com 973-488-7966 or

for more than 7 days unless

NRG TOPICAL ANESTHETIC

benzocaine gel

Product	Inform	ation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:42756-1109

Route of Administration DENTAL

Active Ingredient/Active Moiety Ingredient Name BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)

Inactive Ingredients		
Ingredient Name	Strength	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)		
D&C RED NO. 28 (UNII: 767IP0Y5NH)		
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)		

Product Characteristics			
Color	orange	Score	
Shape		Size	
Flavor	MANGO	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:42756- 1109-3	30 g in 1 JAR; Type 0: Not a Combination Product	06/01/2018	

Marketing In	arketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not	no.+3E6	00/01/2012		

final par (230) 00/01/2013

Labeler - IQ Dental Supply, LLC (800349763)

Revised: 2/2022 IQ Dental Supply, LLC