MEMBERS MARK LANSOPRAZOLE- lansoprazole capsule, delayed release Sam's West Inc

Sam's West, Inc. Lansoprazole Drug Facts

Active ingredient (in each capsule)

Lansoprazole 15 mg

Purpose

Acid Reducer

Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to lansoprazole
- lansoprazole may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss

- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

 taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the package. It contains important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients

D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, gelatin, hypromellose, low substituted hydroxypropyl cellulose, mannitol, meglumine, methacrylic acid copolymer, pharmaceutical ink, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide

Questions or comments?

1-800-809-0469

Package/Label Principal Display Panel

Member's Mark™

COMPARE TO PREVACID® 24 HR ACTIVE INGREDIENT

VALUE 3 PACK

Lansoprazole

DELAYED-RELEASE CAPSULES 15 mg | ACID REDUCER

Treats Frequent Heartburn

24 HR | MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

SODIUM FREE

THREE 14-DAY COURSES OF TREATMENT

42 CAPSULES

ACTUAL SIZE



Drug Facts

Active ingredient (in each capsule) Lansoprazole 15 mg.....

Purpose .Acid Reducer

Þ

Ilee

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings Allergy alert

- do not use if you are allergic to lansop razole
- lansoprazole may cause severe skin reactions. Symptoms mayinclude:
 - skin reddening bilsters rash
- If an allergic reaction occurs, stop use and seek medical help right a way.

Do not use If you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- he artburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have liver disease

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with he artburn
- un explained weight loss nausea or vomiting stomach pain

Ask a doctor or pharmacist before use if you are

■ taking a prescription drug. Add reducers may interact with certain prescription drugs.

Stop use and a sk a doctor if your hear tourn continues or worsens

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea you develop a rash or joint pain

If pregnant or breast-feeding, aska health care professional before use. Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ adults 18 years of age and older

- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days

A7971 ZYS4

- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Drug Facts (continued)

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- m do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of a ge: ask a doctor before use. Heartburn in children may some times be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the package. It contains important information.
- protect product from moisture
 close cap tightly after use

Inactive ingredients D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, gelath, hypromellose, low substituted hydroxypropyl cellulose, mannitol, meglumine, methacrylic add copolymer, pharmaceutical hk, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide

Questions or comments? 1-800-809-0469



Tips for Managing Heartburn

- Avoid foods or drinks that a remore likely to cause heartburn, such as rich, spicy, fatty and fried foods, chood a te, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Donot eat late at night or just before bedtime.
- Do not lie flat or bend oversoon afte reating.
- Raise the head of your bed.
- ■Wearloose-fitting clothing around your stomach.
- If you are overweight, lose weight
- If you smoke, quit smoking.



KEEP THE PACKAGE, IT CONTAINS IMPORTANT INFORMATION.

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

*This product is not manufactured or distributed by Takeda Pharmaceutica is U.S.A., Inc., owner of the registered trademark Prevacid®.

DISTRIBUTED BY: SAM'S WEST, INC. BENTONVILLE, AR 727 16



MEMBERS MARK LANSOPRAZOLE

lansoprazole capsule, delayed release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68196-279

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength	

LANSOPRAZOLE (UNII: 0K5C5T2QPG) (LANSOPRAZOLE - UNII: 0K5C5T2QPG) LANSOPRAZOLE 15 mg

Inactive Ingredients

Ingredient Name		
DCC DED NO. 20 (UNIII, ZCZIDOVENIII)		

D&C RED NO. 28 (UNII: 767IP0Y5NH)

D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) **FD&C BLUE NO. 1** (UNII: H3R47K3TBD)

FD&C RED NO. 40 (UNII: WZB9127XOA)

GELATIN, UNSPECIFIED (UNII: 2G86QN327L)

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)

MANNITOL (UNII: 30WL53L36A)
MEGLUMINE (UNII: 6HG8UB2MUY)

METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

POLYSORBATE 80 (UNII: 60ZP39ZG8H)

SODIUM LAURYL SULFATE (UNII: 368GB5141J)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	PINK, GREEN	Score	no score	
Shape	CAPSULE	Size	15mm	
Flavor		Imprint Code	24HR	
Contains				

Packaging

ı		gg			
ı	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:68196-279- 03	3 in 1 PACKAGE	07/15/2024	
	1	NDC:68196-279- 01	14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing A
Category

Application Number or Monograph Citation Marketing Start Date

Marketing End Date ANDA ANDA202319 07/15/2024

Labeler - Sam's West Inc (051957769)

Revised: 7/2024 Sam's West Inc