

**HYDROGEN PEROXIDE- hydrogen peroxide solution**  
**McKesson Medical-Surgical Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Hydrogen Peroxide**

**Drug Facts**

**Active Ingredient**

Hydrogen Peroxide (stabilized) 3%

**Purpose**

First Aid Antiseptic/Oral Debriding Agent

**Uses**

- First aid to help prevent the risk of infection in minor cuts, scrapes and burns
- Aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

**Warnings**

- For external use only
- Do not use
  - in the eyes or apply over large areas of the body
  - longer than 1 week
- Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns
- Stop use and ask a doctor if
  - the condition persists or gets worse
  - sore mouth symptoms do not improve in 7 days
  - irritation, pain, or redness persists or worsens
  - swelling, rash, or fever develops
- **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

First aid antiseptic:

- Clean the affected area
- Apply a small amount of product on the area 1 to 3 times daily
- May be covered with a sterile bandage
- If bandaged, let dry first

Oral debriding agent (oral rinse):

adults and children 2 years of age and over:

- Mix with an equal amount of water
- Swish around in the mouth over the affected area for at least 1 minute and then spit out
- Use up to 4 times daily after meals

• If damaged, return first

and at bedtime or as directed by a  
dentist or doctor

- 
- Children under 12 years of age should be supervised in the use of this product
  - Children under 2 years of age: consult a dentist or doctor

### **Other Information**

Keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

### **Inactive Ingredient**

purified water

Distributed By McKesson Medical-Surgical Inc.  
Richmond, VA 23228

### **PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label**

NDC 68599-2301-6

**McKESSON**

Hydrogen

Peroxide 3%

TOPICAL SOLUTION USP  
FIRST AID ANTISEPTIC  
ORAL DEBRIDING AGENT

For treatment of minor cuts and abrasions.  
For use as a gargle or rinse. Tamper evident cap  
for your protection. If ring-band is detached from  
cap or missing, do not use.

**16 FL OZ (1 pt)**  
(473 mL)

MFR # 23-D0012

Distributed By McKesson Medical-Surgical Inc.

Richmond, VA 23233

LU-5750-1112 1-07-D0012R10

Made in USA

NDC 68599-2301-6

**McKESSON**

# Hydrogen Peroxide 3%

TOPICAL SOLUTION USP | FIRST AID

ANTISEPTIC | ORAL DEBRIDING AGENT

For treatment of minor cuts and abrasions.  
For use as a gargle or rinse. Tamper evident cap  
for your protection. If ring-band is detached from  
cap or missing, do not use.

**16 fl oz**

(473 mL)

Distributed by  
McKesson Medical-Surgical Inc.  
Richmond, VA 23233  
PVN B0719  
LU-5750-1112 1-07-D0012R10F  
Made in USA

MFR # 23-D0012

**PRINCIPAL DISPLAY PANEL - 473 mL Drug facts panel**

***Drug Facts***

## Drug Facts

Active Ingredient	Purpose
Hydrogen Peroxide (stabilized) 3%	First Aid Antiseptic/ Oral Debriding Agent

**Uses** ■ First aid to help prevent the risk of infection in minor cuts, scrapes and burns ■ Aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

**Warnings** For external use only.  
Do not use ■ in the eyes or apply over large areas of the body ■ longer than 1 week ■ Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns ■ Stop use and ask a doctor if ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** First aid antiseptic ■ Clean the affected area ■ Apply a small amount of product on the area 1 to 3 times daily ■ May be covered with a sterile bandage ■ If bandaged, let dry first ■ Oral debriding agent (oral rinse): adults and children 2 years of age and over: Mix with an equal amount of water ■ Swish around in the mouth over the affected area for at least 1 minute and then spit out ■ Use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

■ Children under 12 years of age should be supervised in the use of this product ■ Children under 2 years of age: consult a dentist or doctor

**Other Information**  
Keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

**Inactive Ingredient** purified water



## Drug Facts

Active Ingredient	Purpose
Hydrogen Peroxide (stabilized) 3%	First Aid Antiseptic/ Oral Debriding Agent

**Uses** ■ First aid to help prevent the risk of infection in minor cuts, scrapes and burns ■ Aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

**Warnings** For external use only.  
Do not use ■ in the eyes or apply over large areas of the body ■ longer than 1 week ■ Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns ■ Stop use and ask a doctor if ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** First aid antiseptic ■ Clean the affected area ■ Apply a small amount of product on the area 1 to 3 times daily ■ May be covered with a sterile bandage ■ If bandaged, let dry first ■ Oral debriding agent (oral rinse): adults and children 2 years of age and over: Mix with an equal amount of water ■ Swish around in the mouth over the affected area for at least 1 minute and then spit out ■ Use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

■ Children under 12 years of age should be supervised in the use of this product ■ Children under 2 years of age: consult a dentist or doctor

**Other Information**  
Keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

**Inactive Ingredient** purified water



# HYDROGEN PEROXIDE

hydrogen peroxide solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68599-2301
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>HYDROGEN PEROXIDE</b> (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	30 mg in 1 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:68599-2301-2	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/09/2013	
2	NDC:68599-2301-6	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/09/2013	
3	NDC:68599-2301-9	3785 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/09/2013	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part333A	10/09/2013	

**Labeler** - McKesson Medical-Surgical Inc. (023904428)

Revised: 12/2019

McKesson Medical-Surgical Inc.