

**VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU- acetaminophen,
dextromethorphan hbr, doxylamine succinate capsule, liquid filled
The Procter & Gamble Manufacturing Company**

VICKS® NyQuil™ HIGH BLOOD PRESSURE COLD & FLU, LiquiCaps™

Drug Facts

Active ingredients (in each liquicap)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Doxylamine succinate 6.25 mg

Active ingredients (in each liquicap) Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 Liquicaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs

adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

store at no greater than 25°C (77°F)

Inactive ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.

DIST. BY PROCTER & GAMBLE,

CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 24 LiquiCaps™

MAX STRENGTH

VICKS®

NyQuil™

HIGH BLOOD PRESSURE

COLD & FLU

HBP

Acetaminophen- Pain reliever/fever reducer

Doxylamine Succinate - Antihistamine

Dextromethorphan HBr - Cough suppressant

DECONGESTANT FREE

24 LIQUICAPS™

<p>Drug Facts</p> <p>Active ingredients (in each LiquiCap)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 10 mg Cough suppressant</p> <p>Doxylamine succinate 6.25 mg Antihistamine</p> <p>Purpose</p>	<p>Drug Facts (continued)</p> <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> • skin reddening • blisters • rash <p>If a skin reaction occurs, stop use and seek medical help right away</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>
<p>Uses temporarily relieves common cold/flu symptoms:</p> <ul style="list-style-type: none"> • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever • runny nose & sneezing 	<p>Do not use</p> <ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
<p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> • more than 8 Liquicaps in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product 	<p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • liver disease • glaucoma • cough that occurs with too much phlegm (mucus)

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • taking sedatives or tranquilizers • taking the blood thinning drug warfarin <p>When using this product</p> <ul style="list-style-type: none"> • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness 	<p>Drug Facts (continued)</p> <p>reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <table border="1"> <tr> <td>• take only as directed</td> <td>• do not exceed 8 LiquiCaps per 24 hrs</td> </tr> <tr> <td>adults & children 12 yrs & over</td> <td>2 LiquiCaps with water every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>	• take only as directed	• do not exceed 8 LiquiCaps per 24 hrs	adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
• take only as directed	• do not exceed 8 LiquiCaps per 24 hrs								
adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs								
children 4 to under 12 yrs	ask a doctor								
children under 4 yrs	do not use								
<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. <p>These could be signs of a serious condition.</p>	<p>Other information store at no greater than 25°C (77°F)</p> <p>Inactive Ingredients D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water</p>								
<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of</p>	<p>Questions? 1-800-362-1683</p>								

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-804
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
POLYVINYL ACETATE PHTHALATE (UNII: 58QVG85GW3)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MICA (UNII: V8A1AW0880)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	NQ
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-804-24	1 in 1 CARTON	07/11/2023	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/11/2023	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Establishment

Name	Address	ID/FEI	Business Operations
Patheon Softgels Inc.		002193829	manufacture(69423-804)

Revised: 11/2025

The Procter & Gamble Manufacturing Company