

**LINCOMIX 50- lincomycin hydrochloride granule**  
**Phibro Animal Health**

-----  
**LINCOMIX® 50(lincomycin)**

**LINCOMIX® 50**

***(lincomycin Type A medicated article)***

**50 grams/lb**

**Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**For further manufacturing only.**

**INDICATIONS FOR USE**

**INDICATIONS FOR USE**

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broiler chickens. Not for use in laying hens, breeder chickens, or turkeys.

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* in swine.

For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae* in swine.

**Each pound contains:**

**Active Drug Ingredient**

Lincomycin.....50 grams

(as lincomycin hydrochloride agricultural grade)

**Inactive Ingredients**

Soybean hulls, #20 grind; light liquid paraffin, Ph.Eur.

**STORAGE, HANDLING AND DISPOSAL**

Store opened bag in dry place to prevent caking. Store at room temperature.

Approved by FDA under NADA # 097-505

**LOT:**

**EXP:**

## **MFG:**

**NET WEIGHT 50 lb (22.7 kg)**

## **BROILERS**

### **BROILER CHICKENS**

#### **INDICATIONS FOR USE**

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broiler chickens. Not for use in laying hens, breeding chickens, or turkeys.

#### **MIXING DIRECTIONS (Type C medicated feeds)**

Thoroughly mix LINCOMIX 50 Type A medicated article with non-medicated broiler chicken feed to obtain 2 g/ton lincomycin in the complete Type C medicated feed. Prepare an intermediate pre-blend of the Type A medicated article prior to mixing in a complete Type C medicated feed. Thoroughly mix 0.04 lb (18 grams) of the Type A medicated article in a convenient quantity of feed ingredients, then thoroughly mix this pre-blend with the rest of the feed ingredients to ensure complete and uniform distribution of the Type A medicated article.

#### **FEEDING DIRECTIONS**

Feed a Type C medicated feed containing 2 g/ton lincomycin as the sole ration to broiler chickens.

## **SWINE**

### **SWINE**

#### **INDICATIONS FOR USE**

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* in swine. For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae* in swine.

#### **MIXING DIRECTIONS (Type B medicated feeds)**

Thoroughly mix LINCOMIX 50 Type A medicated article with non-medicated swine feed according to the table below to obtain the proper concentration in the Type B medicated feed (201 to 40,000 grams lincomycin per ton).

The following table gives examples of how some Type B medicated feed concentrations can be prepared:

---

<b>Pounds of LINCOMIX 50 Type A Medicated Article to Add per Ton to Make a Type B Medicated Feed</b>	<b>Resulting Lincomycin Concentration in Type B Medicated Feed (grams/ton)</b>
--	--

4.02	201
20	1,000
100	5,000
200	10,000
400	20,000
800	40,000

### **MIXING DIRECTIONS (Type C medicated feeds)**

Thoroughly mix LINCOMIX 50 Type A medicated article with non-medicated swine feed according to the table below to obtain the proper concentration of lincomycin in the complete Type C medicated feed. Prepare an intermediate pre-blend of the Type A medicated article prior to mixing in a complete Type C medicated feed. Thoroughly mix the required amount of Type A medicated article in a convenient quantity of feed ingredients, then thoroughly mix this pre-blend with the rest of the feed ingredients to ensure complete and uniform distribution of the Type A medicated article.

The following table gives examples of how some Type C medicated feed concentrations can be prepared:

<b>Pounds of LINCOMIX 50 Type A Medicated Article to Add per Ton to Make a Type C Medicated Feed</b>	<b>Resulting Lincomycin Concentration in Type C Medicated Feed (grams/ton)</b>
0.8	40
2	100
3	150
4	200

### **FEEDING DIRECTIONS**

For the treatment of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* in swine.

Feed a Type C medicated feed containing 100 g/ton lincomycin as the sole ration to swine for 3 weeks or until signs of disease (watery, mucoid, or bloody stools) disappear.

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* in swine.

Feed a Type C medicated feed containing 100 g/ton lincomycin as the sole ration to swine for 3 weeks or until signs of disease (water, mucoid, or bloody stools) disappear, followed by a Type C medicated feed containing 40 g/ton lincomycin as the sole ration.

For the control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* in swine.

For use in swine or on premises with a history of swine dysentery but where symptoms have not yet occurred. Feed a Type C medicated feed containing 40 g/ton lincomycin as the sole ration to swine.

For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae* in swine.

Feed a Type C medicated feed containing 100 to 200 g/ton lincomycin as the sole ration to swine for 21 days.

## **WITHDRAWAL PERIODS**

No withdrawal period is required when used according to labeling.

## **USER SAFETY WARNING**

Not for use in humans. Keep out of reach of children.

## **ANIMAL SAFETY WARNINGS**

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

## **ADDITIONAL RECOMMENDATIONS**

Good Manufacturing Practices should be observed in preparing feeds containing LINCOMIX 50. This includes appropriate clean-out procedures to avoid cross-contamination.

### **Restricted Drug (California)—Use Only as Directed**

Phibro, Phibro logo and Lincomix are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

**Questions/Comments?** For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 888-403-0074 or <http://www.pahc.com>. To report side effects, contact Phibro Animal Health Corporation at 1-888-475-7355.

For additional information about reporting side effects, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Distributed by:  
Phibro Animal Health Corporation 871-9015-04  
Teaneck, NJ 07666

871-9015-04

Made in China


**TAKE TIME**



**OBSERVE LABEL  
DIRECTIONS**

**Take time statement**

**PRINCIPAL DISPLAY PANEL**



Z Reference No.	FG Material No.	Description	Colors
NA	10000802	Lincomix 50 50lb bag	BLACK <span style="border: 1px solid red; padding: 2px;">Dielite</span>
Specification / DWG	Dimensions	Country	PMS 267 C
NA	985 x 308 x 232 mm	USA	
MPD No.	Date	Additional Information	
12889CR2	9/16/25		

**Lincomix 50 Label**

# LINCOMIX 50

lincomycin hydrochloride granule

## Product Information

### Product Type

VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG

### Item Code (Source)

NDC:66104-1494

### Route of Administration

ORAL

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
LINCOMYCIN HYDROCHLORIDE (UNII: M6T05Z2B68) (LINCOMYCIN - UNII:BOD072YW0F)	LINCOMYCIN	50 g in 0.453 kg

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:66104-1494-2	22.7 kg in 1 BAG		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NADA	NADA097505	06/30/1976	

**Labeler** - Phibro Animal Health (006989008)

Revised: 3/2026

Phibro Animal Health