

TREATONIC TATTOO NUMBING CREAM- tattoo numbing cream cream
Shenzhen Situya Trading Co., Ltd.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredient

LIDOCAINE 5%

Purpose

Local Anesthetic

Use

For temporary relief local discomfort, itching, pain, soreness, or burning in the perianal area associated with anorectal disorders and anorectal inflammation.

Do not use

Pregnant or breastfeeding, if there is any allergic reaction to this product.
In case of accidental overdose, get medical help, or contact the poison control center immediately.

When Using

Only take the recommended daily dosage unless otherwise instructed by a doctor.
Some individuals may experience allergic reactions to the ingredients contained in this product.
Never put it into the rectum by using the finger or any medical devices.

Stop Use

if discomfort persists for 7 days.

Ask Doctor

if discomfort persists for 7 days.

Keep Out Of Reach Of Children

and avoid contact with eyes.

In case of accidental ingestion, seek medical attention immediately

Directions

For adults: Cleanse, Rinse, and dry the affected area with warm water. Apply a thick numbing

cream onto the treated area. Apply up to 6 times a day.

For children under 12 years of age: Consult a doctor for guidance.

Other information

Store at room temperature and out of direct sunlight or heat.

Inactive ingredients

Carbomer 934, Emu Oil, Arnica, Lecithin(Soybean), Propylene Glycol, Tocopheryl Acetate, Benzyl Alcohol, Water.

Questions

service@treatonic.com

Warnings

For external use only.

PRINCIPAL DISPLAY PANEL

Recommended Steps for Preparing Before Tattooing:

- Cleanse, rinse, and dry the treated area with warm water.
- Apply a thick numbing cream onto the treated area, ensuring the cream covers the tattoo area and its immediate surroundings.
- When the treated area slightly warms up, re-apply and remain for 60 minutes.
- Remove the plastic wrap and wipe away any excess cream. Wait for an additional 10 minutes for peak numbness.
- Begin the procedure.

Directions:

- For adults: • Cleanse, rinse, and dry the affected area with warm water. • Apply a thick numbing cream onto the treated area. • Apply up to 6 times a day.
- For children under 12 years of age: • Consult a doctor for guidance.

Warnings:

- Do not use on children, pregnant women, or on the face.
- Do not use on broken skin, sunburned skin, or on areas with open wounds.
- Do not use on areas with severe allergic reactions, or on areas with severe skin irritation.
- Do not use on areas with severe allergic reactions, or on areas with severe skin irritation.
- Do not use on areas with severe allergic reactions, or on areas with severe skin irritation.

Other information:

- Contains 5% Lidocaine, 2% Emu Oil, and Arnica.
- Contains 5% Lidocaine, 2% Emu Oil, and Arnica.
- Contains 5% Lidocaine, 2% Emu Oil, and Arnica.

Customer Care:

- For any questions, please contact us for a quick and satisfactory solution.
- For a quick and satisfactory solution, please contact us for a quick and satisfactory solution.
- For a quick and satisfactory solution, please contact us for a quick and satisfactory solution.

Distributed By:

- 180 KARIMA INC.
- BROOKLYN, NY 11223

NET WT 1.7 OZ/50ML

178*26



60*60*47mm

TREATONIC TATTOO NUMBING CREAM			
tattoo numbing cream cream			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82739-013
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	5 g in 100 mL
Inactive Ingredients			
	Ingredient Name		Strength

CARBOMER 934 (UNII: Z135WT9208)	0.4 g in 100 mL
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	6 g in 100 mL
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	1 g in 100 mL
WATER (UNII: 059QF0KO0R)	86.8 g in 100 mL
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	0.3 g in 100 mL
BENZYL ALCOHOL (UNII: LKG8494WBH)	0.5 g in 100 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82739-013-01	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/04/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	M016	07/04/2023	

Labeler - Shenzhen Situya Trading Co., Ltd. (706154255)

Establishment

Name	Address	ID/FEI	Business Operations
Shenzhen Situya Trading Co., Ltd.		706154255	manufacture(82739-013) , label(82739-013)

Revised: 7/2023

Shenzhen Situya Trading Co., Ltd.