

**CORICIDIN HBP COUGH AND COLD- chlorpheniramine maleate,  
dextromethorphan hydrochloride tablet  
A-S Medication Solutions**

-----  
**Coricidin HBP Cough and Cold UI 1614831**

***Drug Facts***

***Active ingredients (in each tablet) Purpose***

Chlorpheniramine maleate 4 mg.....Antihistamine

Dextromethorphan hydrobromide 30 mg.....Cough suppressant

***Uses***

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves runny nose and sneezing due to the common cold

***Warnings***

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

**Do not use to sedate children.**

**Ask a doctor before use if you have**

- trouble urinating due to an enlarged prostate gland
- cough that occurs with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** cough persists for more than 7 days, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact

a Poison Control Center right away. **Abuse of this product can lead to serious injury.**

### **Directions**

- do not exceed recommended dose
- adults and children 12 years and over: take 1 tablet every 6 hours, not more

than 4 tablets in 24 hours

- children under 12 years of age: do not use

### **Other information**

- store between 20° to 25°C (68° to 77°F)
- see end flap for expiration date and lot number

**Inactive ingredients** croscarmellose sodium, D&C red #27 aluminum lake, FD&C yellow #6 aluminum lake, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, stearic acid, talc, titanium dioxide

**Questions or comments? 1-800-317-2165** (Mon-Fri 9AM - 5PM EST)

## **HOW SUPPLIED**

Product: 50090-5534

NDC: 50090-5534-0 8 TABLET in a BLISTER PACK / 2 in a CARTON

## **CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HYDROCHLORIDE**



## **CORICIDIN HBP COUGH AND COLD**

chlorpheniramine maleate, dextromethorphan hydrochloride tablet

### **Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50090-5534(NDC:11523-0008)
---------------------	----------------	---------------------------	--------------------------------

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg

Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>D&amp;C RED NO. 27 ALUMINUM LAKE</b> (UNII: ZK64F7XSTX)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	CC
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50090-5534-0	2 in 1 CARTON	05/04/2021	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/23/2020	

**Labeler -** A-S Medication Solutions (830016429)

**Establishment**

Name	Address	ID/FEI	Business Operations
A-S Medication Solutions		830016429	RELABEL(50090-5534)

Revised: 3/2024

A-S Medication Solutions