

51-022523-00/July, 2003




**NARCAN<sup>®</sup>**  
(Naloxone Hydrochloride Injection, USP)

*Opioid Antagonist*


**R<sub>x</sub> only**

**DESCRIPTION**

NARCAN (naloxone hydrochloride injection, USP), an opioid antagonist, is a synthetic congener of oxymorphone. In structure it differs from oxymorphone in that the methyl group on the nitrogen atom is replaced by an allyl group.



•HCl

NALOXONE HYDROCHLORIDE  
(-)-17-Allyl-4, 5 $\alpha$ -epoxy-3, 14 - dihydroxy   
morphinan-6-one hydrochloride

Naloxone hydrochloride occurs as a white to slightly off-white powder, and is soluble in water, in dilute acids, and in strong alkali; slightly soluble in alcohol; practically insoluble in ether and in chloroform.

NARCAN injection is available as a sterile solution for intravenous, intramuscular and subcutaneous administration in three concentrations: 0.02 mg, 0.4 mg and 1 mg of naloxone hydrochloride per mL.

pH is adjusted to  $3.5 \pm 0.5$  with hydrochloric acid.

The 0.02 mg/mL strength is an unpreserved, paraben-free formulation containing 9 mg/mL sodium chloride.

The 0.4 mg/mL vial contains 8.6 mg/mL of sodium chloride and 2 mg/mL of methylparaben and propylparaben as preservatives in a ratio of 9:1. The 0.4 mg/mL ampul is also available in an unpreserved, paraben-free formulation containing 9 mg/mL of sodium chloride.

The 1 mg/mL vial contains 8.35 mg/mL of sodium chloride and 2 mg/mL of methylparaben and propylparaben as preservatives in a ratio of 9:1. The 1 mg/mL ampul is also available in an unpreserved, paraben-free formulation containing 9 mg/mL of sodium chloride.

## CLINICAL PHARMACOLOGY

### Complete or Partial Reversal of Opioid Depression

NARCAN prevents or reverses the effects of opioids including respiratory depression, sedation and hypotension. Also, NARCAN can reverse the psychotomimetic and dysphoric effects of agonist-antagonists such as pentazocine.

NARCAN is an essentially pure opioid antagonist, i.e., it does not possess the “agonistic” or morphine-like properties characteristic of other opioid antagonists. When administered in usual doses and in the absence of opioids or agonistic effects of other opioid antagonists, it exhibits essentially no pharmacologic activity.

NARCAN has not been shown to produce tolerance or cause physical or psychological dependence. In the presence of physical dependence on opioids, NARCAN will produce withdrawal symptoms. However, in the presence of opioid dependence, opiate withdrawal symptoms may appear within minutes of NARCAN administration and subside in about 2 hours. The severity and duration of the withdrawal syndrome are related to the dose of NARCAN and to the degree and type of opioid dependence. While the mechanism of action of NARCAN is not fully understood, *in vitro* evidence suggests that NARCAN antagonizes opioid effects by competing for the  $\mu$ ,  $\kappa$  and  $\sigma$  opiate receptor sites in the CNS, with the greatest affinity for the  $\mu$  receptor.

When NARCAN is administered intravenously (I.V.), the onset of action is generally apparent within two minutes. The onset of action is slightly less rapid when it is administered subcutaneously (S.C.) or intramuscularly (I.M.). The duration of action is dependent upon the dose and route of administration of NARCAN. Intramuscular administration produces a more prolonged effect than intravenous administration. Since the duration of action of NARCAN may be shorter than that of some opiates, the effects of the opiate may return as the effects of NARCAN dissipates. The requirement for repeat doses of NARCAN will also be dependent upon the amount, type and route of administration of the opioid being antagonized.

### Adjunctive Use in Septic Shock

NARCAN has been shown in some cases of septic shock to produce a rise in blood pressure that may last up to several hours; however, this pressor response has not been demonstrated to improve patient survival. In some studies, treatment with NARCAN in the setting of septic shock has been associated with adverse effects, including agitation, nausea and vomiting, pulmonary edema, hypotension, cardiac arrhythmias, and seizures. The decision to use NARCAN in septic shock should be exercised with caution, particularly in patients who may have underlying pain or have previously received opioid therapy and may have developed opioid tolerance.

Because of the limited number of patients who have been treated, optimal dosage and treatment regimens have not been established.

## PHARMACOKINETICS

### Distribution

Following parenteral administration, NARCAN is rapidly distributed in the body and readily crosses the placenta. Plasma protein binding occurs but is relatively weak. Plasma albumin is the major binding constituent but significant binding of naloxone also occurs to plasma constituents other than albumin. It is not known whether naloxone is excreted into human milk.

### Metabolism and Elimination

NARCAN is metabolized in the liver, primarily by glucuronide conjugation with naloxone-3-glucuronide as the major metabolite. In one study the serum half-life in adults ranged from 30 to 81 minutes (mean  $64 \pm 12$  minutes). In a neonatal study the mean plasma half-life was observed to be  $3.1 \pm 0.5$  hours. After an oral or intravenous dose, about 25-40% of the drug is excreted as metabolites in urine within 6 hours, about 50% in 24 hours, and 60-70% in 72 hours.

## INDICATIONS AND USAGE

NARCAN is indicated for the complete or partial reversal of opioid depression, including respiratory depression, induced by natural and synthetic opioids, including propoxyphene, methadone and certain mixed agonist-antagonist analgesics: nalbuphine, pentazocine, butorphanol, and cyclazocine. NARCAN is also indicated for diagnosis of suspected or known acute opioid overdose.

NARCAN may be useful as an adjunctive agent to increase blood pressure in the management of septic shock (see **CLINICAL PHARMACOLOGY; Adjunctive Use in Septic Shock**).

## CONTRAINDICATIONS

NARCAN is contraindicated in patients known to be hypersensitive to naloxone hydrochloride or to any of the other ingredients in NARCAN.

## WARNINGS

### Drug Dependence

NARCAN should be administered cautiously to persons including newborns of mothers who are known or suspected to be physically dependent on opioids. In such cases an abrupt and complete reversal of opioid effects may precipitate an acute withdrawal syndrome.

The signs and symptoms of opioid withdrawal in a patient physically dependent on opioids may include, but are not limited to, the following: body aches, diarrhea, tachycardia, fever, runny nose, sneezing, piloerection, sweating, yawning, nausea or vomiting, nervousness, restlessness or irritability, shivering or trembling, abdominal cramps, weakness, and increased blood pressure. In the neonate, opioid withdrawal may also include: convulsions, excessive crying, and hyperactive reflexes.

## Repeat Administration

The patient who has satisfactorily responded to NARCAN should be kept under continued surveillance and repeated doses of NARCAN should be administered, as necessary, since the duration of action of some opioids may exceed that of NARCAN.

## Respiratory Depression due to Other Drugs

NARCAN is not effective against respiratory depression due to non-opioid drugs and in the management of acute toxicity caused by levopropoxyphene. Reversal of respiratory depression by partial agonists or mixed agonist/antagonists, such as buprenorphine and pentazocine, may be incomplete or require higher doses of naloxone. If an incomplete response occurs, respirations should be mechanically assisted as clinically indicated.

## PRECAUTIONS

### General

In addition to NARCAN, other resuscitative measures such as maintenance of a free airway, artificial ventilation, cardiac massage, and vasopressor agents should be available and employed when necessary to counteract acute opioid poisoning.

Abrupt postoperative reversal of opioid depression may result in nausea, vomiting, sweating, tremulousness, tachycardia, increased blood pressure, seizures, ventricular tachycardia and fibrillation, pulmonary edema, and cardiac arrest which may result in death. Excessive doses of NARCAN in postoperative patients may result in significant reversal of analgesia and may cause agitation (see **PRECAUTIONS** and **DOSAGE AND ADMINISTRATION; Usage in Adults-Postoperative Opioid Depression**). Several instances of hypotension, hypertension, ventricular tachycardia and fibrillation, pulmonary edema, and cardiac arrest have been reported in postoperative patients. Death, coma, and encephalopathy have been reported as sequelae of these events. These have occurred in patients most of whom had pre-existing cardiovascular disorders or received other drugs which may have similar adverse cardiovascular effects. Although a direct cause and effect relationship has not been established, NARCAN should be used with caution in patients with pre-existing cardiac disease or patients who have received medications with potential adverse cardiovascular effects, such as hypotension, ventricular tachycardia or fibrillation, and pulmonary edema. It has been suggested that the pathogenesis of pulmonary edema associated with the use of NARCAN is similar to neurogenic pulmonary edema, i.e., a centrally mediated massive catecholamine response leading to a dramatic shift of blood volume into the pulmonary vascular bed resulting in increased hydrostatic pressures.

### Drug Interactions

Large doses of naloxone are required to antagonize buprenorphine since the latter has a long duration of action due to its slow rate of binding and subsequent slow dissociation from the opioid receptor. Buprenorphine antagonism is characterized by a gradual onset of the reversal effects and a decreased duration of action of the normally prolonged respiratory depression. The barbiturate methohexital appears to block the acute onset of withdrawal symptoms induced by naloxone in opiate addicts.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Studies in animals to assess the carcinogenic potential of NARCAN have not been conducted. NARCAN was weakly positive in the Ames mutagenicity and in the *in vitro* human lymphocyte chromosome aberration test but was negative in the *in vitro* Chinese hamster V79 cell HGPRT mutagenicity assay and in the *in vivo* rat bone marrow chromosome aberration study. Reproduction studies conducted in mice and rats at doses 4-times and 8-times, respectively, the dose of a 50 kg human given 10 mg/day (when based on surface area or  $\text{mg}/\text{m}^2$ ), demonstrated no embryotoxic or teratogenic effects due to NARCAN.

## **Use in Pregnancy**

### **Teratogenic Effects: Pregnancy Category C:**

Teratology studies conducted in mice and rats at doses 4-times and 8-times, respectively, the dose of a 50 kg human given 10 mg/day (when based on surface area or  $\text{mg}/\text{m}^2$ ), demonstrated no embryotoxic or teratogenic effects due to NARCAN. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, NARCAN should be used during pregnancy only if clearly needed.

### **Non-teratogenic Effects:**

Risk-benefit must be considered before NARCAN is administered to a pregnant woman who is known or suspected to be opioid-dependent since maternal dependence may often be accompanied by fetal dependence. Naloxone crosses the placenta, and may precipitate withdrawal in the fetus as well as in the mother. Patients with mild to moderate hypertension who receive naloxone during labor should be carefully monitored as severe hypertension may occur.

## **Use in Labor and Delivery**

It is not known if NARCAN (naloxone hydrochloride injection, USP) affects the duration of labor and/or delivery. However, published reports indicated that administration of naloxone during labor did not adversely affect maternal or neonatal status.

## **Nursing Mothers**

It is not known whether NARCAN is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when NARCAN is administered to a nursing woman.

## **Pediatric Use**

NARCAN (naloxone hydrochloride injection, USP) may be administered intravenously, intramuscularly or subcutaneously in children and neonates to reverse the effects of opiates. The American Academy of Pediatrics, however, does not endorse subcutaneous or intramuscular administration in opiate intoxication since absorption may be erratic or delayed. Although the opiate-intoxicated child responds dramatically to NARCAN, he/she must be carefully monitored for at least 24 hours as a relapse may occur as naloxone is metabolized.

When NARCAN is given to the mother shortly before delivery, the duration of its effect lasts only for the first two hours of neonatal life. It is preferable to administer NARCAN

directly to the neonate if needed after delivery. NARCAN has no apparent benefit as an additional method of resuscitation in the newly born infant with intrauterine asphyxia which is not related to opioid use.

#### **Usage in Pediatric Patients and Neonates for Septic Shock:**

The safety and effectiveness of NARCAN in the treatment of hypotension in pediatric patients and neonates with septic shock have not been established. One study of two neonates in septic shock reported a positive pressor response; however, one patient subsequently died after intractable seizures.

#### **Geriatric Use**

Clinical studies of NARCAN did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

#### **Renal Insufficiency/Failure**

The safety and effectiveness of NARCAN in patients with renal insufficiency/failure have not been established in well-controlled clinical trials. Caution should be exercised when NARCAN is administered to this patient population.

#### **Liver Disease**

The safety and effectiveness of NARCAN in patients with liver disease have not been established in well-controlled clinical trials. Caution should be exercised when NARCAN is administered to patients with liver disease.

### **ADVERSE REACTIONS**

#### **Postoperative**

The following adverse events have been associated with the use of NARCAN in postoperative patients: hypotension, hypertension, ventricular tachycardia and fibrillation, dyspnea, pulmonary edema, and cardiac arrest. Death, coma, and encephalopathy have been reported as sequelae of these events. Excessive doses of NARCAN in postoperative patients may result in significant reversal of analgesia and may cause agitation (see **PRECAUTIONS** and **DOSAGE AND ADMINISTRATION; Usage in Adults-Postoperative Opioid Depression**).

#### **Opioid Depression**

Abrupt reversal of opioid depression may result in nausea, vomiting, sweating, tachycardia, increased blood pressure, tremulousness, seizures, ventricular tachycardia and fibrillation, pulmonary edema, and cardiac arrest which may result in death (see **PRECAUTIONS**).

#### **Opioid Dependence**

Abrupt reversal of opioid effects in persons who are physically dependent on opioids may precipitate an acute withdrawal syndrome which may include, but is not limited to, the

following signs and symptoms: body aches, fever, sweating, runny nose, sneezing, piloerection, yawning, weakness, shivering or trembling, nervousness, restlessness or irritability, diarrhea, nausea or vomiting, abdominal cramps, increased blood pressure, tachycardia. In the neonate, opioid withdrawal may also include: convulsions; excessive crying; hyperactive reflexes (see **WARNINGS**).

Adverse events associated with the postoperative use of NARCAN are listed by organ system and in decreasing order of frequency as follows:

**Cardiac Disorders:** pulmonary edema, cardiac arrest or failure, tachycardia, ventricular fibrillation, and ventricular tachycardia. Death, coma, and encephalopathy have been reported as sequelae of these events.

**Gastrointestinal Disorders:** vomiting, nausea

**Nervous System Disorders:** convulsions, paraesthesia, grand mal convulsion

**Psychiatric Disorders:** agitation, hallucination, tremulousness

**Respiratory, Thoracic and Mediastinal Disorders:** dyspnea, respiratory depression, hypoxia

**Skin and Subcutaneous Tissue Disorders:** nonspecific injection site reactions, sweating

**Vascular Disorders:** hypertension, hypotension, hot flushes or flushing.

See also **PRECAUTIONS** and **DOSAGE AND ADMINISTRATION; Usage in Adults; Postoperative Opioid Depression.**

## **DRUG ABUSE AND DEPENDENCE**

NARCAN is an opioid antagonist. Physical dependence associated with the use of NARCAN has not been reported. Tolerance to the opioid antagonist effect of NARCAN is not known to occur.

## **OVERDOSAGE**

There is limited clinical experience with NARCAN overdose in humans.

### **Adult Patients**

In one small study, volunteers who received 24 mg/70 kg did not demonstrate toxicity. In another study, 36 patients with acute stroke received a loading dose of 4 mg/kg (10 mg/m<sup>2</sup>/min) of NARCAN followed immediately by 2 mg/kg/hr for 24 hours. Twenty-three patients experienced adverse events associated with naloxone use, and naloxone was discontinued in seven patients because of adverse effects. The most serious adverse events were: seizures (2 patients), severe hypertension (1), and hypotension and/or bradycardia (3).

At doses of 2 mg/kg in normal subjects, cognitive impairment and behavioral symptoms, including irritability, anxiety, tension, suspiciousness, sadness, difficulty concentrating, and lack of appetite have been reported. In addition, somatic symptoms, including dizziness, heaviness, sweating, nausea, and stomachaches were also reported. Although complete information is not available, behavioral symptoms were reported to often persist for 2-3 days.

### **Pediatric Patients**

Up to 11 doses of 0.2 mg of naloxone (2.2 mg) have been administered to children following overdose of diphenoxylate hydrochloride with atropine sulfate. Pediatric reports include a 2-1/2 year-old child who inadvertently received a dose of 20 mg of naloxone for treatment of respiratory depression following overdose with diphenoxylate hydrochloride with atropine sulfate. The child responded well and recovered without adverse sequelae. There is also a report of a 4-1/2 year-old child who received 11 doses during a 12-hour period, with no adverse sequelae.

### **Patient Management**

Patients who experience a NARCAN overdose should be treated symptomatically in a closely supervised environment. Physicians should contact a poison control center for the most up-to-date patient management information.

### **DOSAGE AND ADMINISTRATION**

NARCAN may be administered intravenously, intramuscularly, or subcutaneously. The most rapid onset of action is achieved by intravenous administration, which is recommended in emergency situations.

Since the duration of action of some opioids may exceed that of NARCAN, the patient should be kept under continued surveillance. Repeated doses of NARCAN should be administered, as necessary.

### **Intravenous Infusion**

NARCAN may be diluted for intravenous infusion in normal saline or 5% dextrose solutions. The addition of 2 mg of NARCAN in 500 mL of either solution provides a concentration of 0.004 mg/mL. Mixtures should be used within 24 hours. After 24 hours, the remaining unused mixture must be discarded. The rate of administration should be titrated in accordance with the patient's response.

NARCAN should not be mixed with preparations containing bisulfite, metabisulfite, long-chain or high molecular weight anions, or any solution having an alkaline pH. No drug or chemical agent should be added to NARCAN unless its effect on the chemical and physical stability of the solution has first been established.

### **General**

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

## Usage in Adults

### Opioid Overdose—Known or Suspected:

An initial dose of 0.4 mg to 2 mg of NARCAN may be administered intravenously. If the desired degree of counteraction and improvement in respiratory functions are not obtained, it may be repeated at two- to three-minute intervals. If no response is observed after 10 mg of NARCAN have been administered, the diagnosis of opioid-induced or partial opioid-induced toxicity should be questioned. Intramuscular or subcutaneous administration may be necessary if the intravenous route is not available.

### Postoperative Opioid Depression:

For the partial reversal of opioid depression following the use of opioids during surgery, smaller doses of NARCAN are usually sufficient. The dose of NARCAN should be titrated according to the patient's response. For the initial reversal of respiratory depression, NARCAN should be injected in increments of 0.1 to 0.2 mg intravenously at two- to three-minute intervals to the desired degree of reversal, i.e., adequate ventilation and alertness without significant pain or discomfort. Larger than necessary dosage of NARCAN may result in significant reversal of analgesia and increase in blood pressure. Similarly, too rapid reversal may induce nausea, vomiting, sweating or circulatory stress.

Repeat doses of NARCAN may be required within one- to two-hour intervals depending upon the amount, type (i.e., short or long acting) and time interval since last administration of an opioid. Supplemental intramuscular doses have been shown to produce a longer lasting effect.

### Septic Shock:

The optimal dosage of NARCAN or duration of therapy for the treatment of hypotension in septic shock patients has not been established (see **CLINICALPHARMACOLOGY**).

## Usage in Children

### Opioid Overdose—Known or Suspected:

The usual initial dose in children is 0.01 mg/kg body weight given I.V. If this dose does not result in the desired degree of clinical improvement, a subsequent dose of 0.1 mg/kg body weight may be administered. If an I.V. route of administration is not available, NARCAN may be administered I.M. or S.C. in divided doses. If necessary, NARCAN can be diluted with sterile water for injection.

### Postoperative Opioid Depression:

Follow the recommendations and cautions under **Adult Postoperative Depression**. For the initial reversal of respiratory depression, NARCAN should be injected in increments of 0.005 mg to 0.01 mg intravenously at two- to three-minute intervals to the desired degree of reversal.

## Usage in Neonates

### Opioid-induced Depression:

The usual initial dose is 0.01 mg/kg body weight administered I.V., I.M. or S.C. This dose may be repeated in accordance with adult administration guidelines for postoperative opioid depression.

## HOW SUPPLIED

NARCAN (naloxone hydrochloride injection, USP) for intravenous, intramuscular, and subcutaneous administration is available as:

### Multiple Dose Vials

0.4 mg/mL            10 mL multiple dose vial-box of 1, NDC 63481-365-05  
1 mg/mL             10 mL multiple dose vial-box of 1, NDC 63481-368-05

### Preservative-Free Ampules

0.02 mg/mL        2 mL unit dose ampule-box of 10, NDC 63481-359-10  
0.4 mg/mL         1 mL unit dose ampule-box of 10, NDC 63481-358-10  
1 mg/mL            2 mL unit dose ampule-box of 10, NDC 63481-377-10

Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F). [See USP Controlled Room Temperature]. Protect from light.

Store in carton until contents have been used.

Manufactured for:

**Endo Pharmaceuticals Inc.**  
Chadds Ford, Pennsylvania 19317



NARCAN is a Registered Trademark of Endo Pharmaceuticals Inc.

Copyright © Endo Pharmaceuticals Inc. 2003

Printed in U.S.A.

51-022523-00/July, 2003