

FRONT LABEL



(b) (4)

2 3/4"
1 11/16"

Code 128
70236M

12 Hour Cough Relief
Children's Delsym
COUGH
Orange Flavored Liquid

NDC 63824-177-65

12 Hour Cough Relief
Children's Delsym
Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

COUGH

Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

COUGH

Also Available
In Grape Flavor



Age 4+
For Children & Adults



Please visit our web site:
www.delsym.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Orange Flavored Liquid 148 mL (5 fl oz)
Day or Night Alcohol-free

12 Hour Cough Relief
Children's Delsym
Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

COUGH

DELSYM® DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided.
Do not use dosing cup with other products.
mL = milliliter

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.




Dosing Cup Included

Contains
No Fever Reducer
or Pain Reliever

5 11/16"

3-63824-27665-6
(UPC @ 100% truncated)

Lot No.:
Exp. Date:

TAMPER EVIDENT: Do not use if the neckband printed with  is broken or missing.

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 ■ chronic cough that lasts as occurs with smoking, asthma or emphysema
 ■ cough that occurs with too much phlegm (mucus)
Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ **shake bottle well before use**
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: **sodium 7 mg**
 ■ store at 20-25°C (68-77°F) ■ dosing cup provided

Inactive ingredients citric acid, edetate disodium, ethylcellulose, FD&C Yellow No. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-888-963-3382
You may also report side effects to this phone number.


C I A 70236M
PT# C I A 60333
8137779
Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224 ©2014 RB 022114

(b) (4)

FRONT LABEL



(b) (4)



(b) (4)