

Seal is broken or missing or if the neckband is not printed. TAMPER EVIDENT.

Delsym

Drug Facts

Active ingredient (in each 5 mL teaspoonful)
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP. Cough suppressant.

Use Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or influenza.

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or if you have taken an MAOI in the last 14 days. Ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:
 ■ chronic cough that lasts as occurs with smoking, asthma or emphysema
 ■ cough that occurs with too much phlegm (mucus)
 ■ stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Directions
Shake bottle well before use
 ■ dose as follows or as directed by a doctor
 adults and children 12 years of age and over 2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
 children 6 to under 12 years of age 1 teaspoonful every 12 hours, not to exceed 2 teaspoonfuls in 24 hours
 children 2 to under 6 years of age 1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours
 children under 2 years of age consult a doctor

Other information
 ■ each 5 mL teaspoonful contains: sodium 6 mg
 ■ store at 20° to 25° C (68° to 77° F)

Inactive ingredients: citric acid, anhydrous; D&G Red #3, indigo disodium, ethylcellulose, FD&C Blue #1, leucine, high fructose corn syrup, methylparaben, polyethylene glycol 3350, polyethylene glycol, propylparaben, purified water, sorbitol, vegetable glycerin, xanthan gum.

Questions? 1-866-682-4639

ADAMS RESPIRATORY THERAPEUTICS
 © 2007. Distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76155
 010307 40006533

12 Hour Cough Relief
Delsym
 dextromethorphan polistirex extended-release suspension
COUGH SUPPRESSANT



Contains
 No Fever Reducer
 or Pain Reliever
 Great Grape Taste

Also Available In Great Orange Flavor

12 Hour Cough Relief
Delsym
 NDC 63824-171-65

12 Hour Cough Relief
Delsym
 dextromethorphan polistirex extended-release suspension
COUGH SUPPRESSANT

NEW GRAPE FLAVOR

Grape-Flavored Liquid
 Alcohol-free
 148 mL (5 fl. oz.)

3-63824-171-65-4

Lot No.
 Exp. Date

12 Hour Cough Relief
Delsym
 dextromethorphan polistirex extended-release suspension
COUGH SUPPRESSANT

DELSYM® DOSING
 SHAKE WELL BEFORE USE.

Age (yr)	Dose (1 TSP = 5 mL)
12 years to adult	2 TEASPOONFULS EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS

See back panel for full dosing directions.

Contains
 No Fever Reducer
 or Pain Reliever
 Great Grape Taste

Please visit our Web site:
www.delsym.com

FILE INFORMATION
 Job Description:
 Drawing #: #C02
 Carton Size (Built)
 Software: Adobe
 Date Work Performed
 Colors Used: 4/C
 Fonts Used: Futu
 Additional Inform
 Help



NDC 63824-171-65

Delsym[®]
 dextromethorphan polistirex
 extended-release suspension
COUGH SUPPRESSANT
12 Hour Cough Relief
 Contains No Fever Reducer
 or Pain Reliever
 Alcohol-free
 Grape-Flavored Liquid
 148 mL (5 fl oz)

© 2007, Registered by
 Adams Respiratory Technologies
 Fort Worth, TX 76155

3
 63824 17165
 4

Lot:
 Exp.:
 010307 4000521

INDICATIONS: Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants.

DIRECTIONS: SHAKE BOTTLE WELL BEFORE USING.

Dose as follows or as directed by a doctor.
Adults and Children 12 years of age and over: 2 teaspoons every 12 hours, not to exceed 4 teaspoons in 24 hours.
Children 6 to under 12 years of age: 1 teaspoonful every 12 hours, not to exceed 2 teaspoons in 24 hours.
Children 2 to under 6 years of age: 1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours.
Children under 2 years of age: Consult a doctor.

WARNINGS: Do not take this product for chronic cough that lasts as occurs with smoking, asthma, or emphysema, or if cough occurs with too much phlegm (mucus), unless directed by a doctor. If cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts, consult a doctor. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

DRUG INTERACTION PRECAUTION: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

ACTIVE INGREDIENT: Each teaspoonful (5 mL) contains dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.

TAMPER EVIDENT: Do not use if the neckband printed with  is broken or missing.

Store at 20°-25°C (68°-77°F).
 Questions? 1-866-582-4639

FILE INFORMATION

Job Description: Delsym Adult Grape 5-oz. Label

Drawing #: Not Available

Label Size: 6-5/16" w x 2-1/2" h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed, Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

Missing or broken Tamper Evident Band (TEB) printed on the neckband printed with the product name and lot number.

Drug Facts

Active ingredient (in each 5 mL teaspoonful) Purpose
 Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP Cough suppressant
 Use temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 chronic cough that lasts as occurs with smoking, asthma or emphysema
 cough that occurs with too much phlegm (mucus)
 Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Shake bottle well before use
 doses as follows or as directed by a doctor

adults and children 12 years of age and over	2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
children 6 to under 12 years of age	1 teaspoonful every 12 hours, not to exceed 2 teaspoonfuls in 24 hours
children 2 to under 6 years of age	1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours
children under 2 years of age	consult a doctor

Other information

each 5 mL teaspoonful contains: sodium 6 mg
 store at 20°-25°C (68°-77°F)

Inactive ingredients citric acid anhydrous, D&C Red #33, edetate disodium, ethylcellulose, FD&C Blue #1, flavor, high fructose corn syrup, methylparaben, polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tagaracanth, vegetable oil, xanthan gum
Questions? 1-866-682-4639



© 2007, Distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76155

GLUE AREA
 NO INK & COATING

12 Hour Cough Relief
Delsym

12 Hour Cough Relief
Delsym
 dextromethorphan polistirex extended-release suspension
 COUGH SUPPRESSANT

12 Hour Cough Relief
Delsym
 dextromethorphan polistirex extended-release suspension
 COUGH SUPPRESSANT



Grape-Flavored Liquid
 Alcohol-free
 89 mL (3 fl oz)



Contains No Fever Reducer or Pain Reliever
 Great Grape Taste

12 Hour Cough Relief
Delsym
 dextromethorphan polistirex extended-release suspension
 COUGH SUPPRESSANT

12 Hour Cough Relief
Delsym
 dextromethorphan polistirex extended-release suspension
 COUGH SUPPRESSANT

DELSYM® DOSING
 SHAKE WELL BEFORE USE.

Age (yr)	Dose (1 TSP = 5 mL)
12 years to adult	2 TEASPOONFULS EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS

See back panel for full dosing directions.

Contains No Fever Reducer or Pain Reliever
 Great Grape Taste

Please visit our Web site:
www.delsym.com

3-63824-171-63-0

Lot No.:
 Exp. Date:

GLUE AREA

NO INK & COATING

FILE INFORMATION
 Job Description:
 Drawing #: #C
 Carton Size (B
 Software: Ado
 Date Work Per
 Colors Used:
 Fonts Used: F
 Additional Infr



FILE INFORMATION

Job Description: Delsym Adult Grape 3-oz. Label

Drawing #: Not Available

Label Size: 5-5/8" w x 2-1/8" h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed, Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

NDC 63824-171-63

Delsym
dextromethorphan polistirex
extended-release suspension
COUGH SUPPRESSANT
12 Hour Cough Relief
Contains No Fever Reducer
or Pain Reliever
Alcohol-free
Grape-Flavored Liquid
89 mL (3 fl oz)

© 2007, Distributed by
Adams Respiratory Therapeutics
Fort Worth, TX 76135

Lot:
Exp:

3 63824 17163 0 010307 4000517

INDICATIONS: Temporarily relieves cough due to minor
croup and bronchitis. Irritation may occur with the common
cold or influenza viruses.

DIRECTIONS: SHAKE BOTTLE WELL BEFORE USING.

Dose as follows or as directed by a doctor.

Adults and Children 12 years of age and over: 2 teaspoonfuls
every 12 hours, not to exceed 4 teaspoonfuls in 24 hours.

Children 6 to under 12 years of age: 1 teaspoonful every
12 hours, not to exceed 2 teaspoonfuls in 24 hours.

Children 2 to under 6 years of age: ½ teaspoonful every
12 hours, not to exceed 1 teaspoonful in 24 hours.

Children under 2 years of age: Consult a doctor.


WARNINGS: Do not take this product for chronic cough that
lasts as occurs with smoking, asthma, or emphysema, or if
cough occurs with too much phlegm (mucus) unless directed
by a doctor. If cough lasts more than 7 days, cough comes back,
or occurs with fever, rash or headache that lasts, consult a
doctor. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before
use. **Keep out of reach of children.** In case of overdose, get
medical help or contact a Poison Control Center right away.

DRUG INTERACTION PRECAUTION: Do not use if you are now
taking a prescription medicine (cough suppressant) (Cetane
Parsons's disease), or for 2 weeks after stopping the MAOI
drug. If you do not know if your prescription drug contains an
MAOI, ask a doctor or pharmacist before taking this product.

ACTIVE INGREDIENT: Each teaspoonful (5 mL) contains
dextromethorphan polistirex equivalent
to 30 mg dextromethorphan
hydrobromide.

TAMPER EVIDENT:
Do not use if the
neck cap is broken or missing.

Store at 20°-25°C (68°-77°F).
Questions? 1-866-882-4639



FILE INFORMATION

Job Description: Delsym Pediatric Grape 5-oz. Label

Drawing #: Not Available

Label Size: 6-5/16" w x 2-1/2" h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed, Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

the
backprint printed with with
if you are not using the
backprint printed with with

Drug Facts

Active ingredient (in each 5 mL teaspoonful) Purpose
Dextromethorphan polistirex equivalent to
30 mg dextromethorphan hydrobromide..... Cough suppressant
Use temporarily relieves cough due to minor throat and bronchial
irritation as may occur with the common cold or inhaled irritants

Warnings

Do not use if you are now taking a prescription monoamine oxidase
inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional
conditions, or Parkinson's disease, or for 2 weeks after stopping the
MAOI drug. If you do not know if your prescription drug contains an
MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ chronic cough that lasts as occurs with smoking, asthma or emphysema
■ cough that occurs with too much phlegm (mucus)
■ Stop use and ask a doctor if cough lasts more than 7 days, cough comes
back, or occurs with fever, rash or headache that lasts. These could be
signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or
contact a Poison Control Center right away.

Directions

■ Shake bottle well before use
■ dose as follows or as directed by a doctor
adults and children 2 teaspoonfuls every 12 hours,
12 years of age and over not to exceed 4 teaspoonfuls in 24 hours
children 6 to under 12 1 teaspoonful every 12 hours,
not to exceed 2 teaspoonfuls in 24 hours
children 2 to under 6 1/2 teaspoonful every 12 hours,
not to exceed 1 teaspoonful in 24 hours
children under 2 years of age consult a doctor

Other information

■ each 5 mL teaspoonful contains sodium 6 mg
■ store at 20°-25° C (68°-77° F)

Inactive ingredients: citric acid anhydrous, D&C Red #3, edetate
disodium, ethylcellulose, FD&C Blue #2, ferrous hydroxide, iron
oxide, methylparaben, polyethylene glycol 550, polyvinylpyrrolidone
K30, polyvinylpyrrolidone K90, polyvinylpyrrolidone K120, xanthan gum
Questions? 1-866-682-4639



© 2007. Distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76155

021207
4000530

GLUE AREA
NO INK & COATING

12 Hour Cough Relief
Delsym

NDC 63824-172-63

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
COUGH SUPPRESSANT

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**



**NEW GRAPE
FLAVOR**



Contains
No Fever Reducer
or Pain Reliever
Great Grape Taste

Also Available
In Great
Orange Flavor

Grape-Flavored Liquid
for Children & Adults
Alcohol-free
89 mL (3 fl oz)



DELSYM® DOSING SHAKE WELL BEFORE USE.	Dose
Age (Yr)	(1 TSP = 5 mL)
Under 2	CONSULT DOCTOR
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS

Please visit our Web site:
www.delsym.com

Contains
No Fever Reducer
or Pain Reliever
Great Grape Taste

3-63824-172-63-7

Lot No.:
Exp. Date:

GLUE AREA

NO INK & COATING

Job Description:
Drawing #: #C03
Carton Size (Built)
Software: Adobe
Date Work Perfor
Colors Used: 4/C
Fonts Used: Frut
Helv
Additional Infor



FILE INFORMATION

Job Description: Delsym Pediatric Grape 3-oz. Label

Drawing #: Not Available

Label Size: 5-5/8" w x 2-1/8" h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed, Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

Additional information:

Contains
 No Fever Reducer
 or Pain Reliever
 Great Grape Taste

Delsym
 dextromethorphan polistirex
 extended-release suspension
**COUGH
 SUPPRESSANT**

Delsym
 dextromethorphan polistirex
 extended-release suspension
**COUGH
 SUPPRESSANT**

12 Hour Cough Relief



Delsym
 dextromethorphan polistirex
 extended-release suspension
**COUGH
 SUPPRESSANT**

NDC 63824-172-75

12 Hour Cough Relief

TAMPER EVIDENT:
 Do not use if the neckband
 printed with is broken or missing.

Delsym 12 Hour Cough Relief

See side panel for full dosing directions.

Age (yr)	Dose (1 TSP = 5 mL)
Under 2	CONSULT DOCTOR
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each 5 mL teaspoonful) Purpose</p> <p>Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... Cough suppressant</p> <p>Use temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants</p> <p>Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the</p>	<p>MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> chronic cough that lasts as occurs with smoking, asthma or emphysema cough that occurs with too much phlegm (mucus) <p>Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>

LOT & EXP DATE AREA
 NO INK

Grape-Flavored
 Liquid for
 Children & Adults

Delsym
 dextromethorphan polistirex
 extended-release suspension
**COUGH
 SUPPRESSANT**

Alcohol-free
 12 bottles
 containing
 15 mL each
 PHYSICIAN SAMPLES —
 NOT TO BE SOLD

Directions	Drug Facts (continued)
<ul style="list-style-type: none"> shake well before use use as follows or as directed by doctor adults and children 12 years of age and over 2 teaspoonful every 12 hours, not to exceed 4 teaspoonful in 24 hours children 6 to under 12 years of age 1 teaspoonful every 12 hours, not to exceed 2 teaspoonful in 24 hours children 2 to under 6 years of age 1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours children under 2 years of age consult a doctor 	<p>Drug Facts (continued)</p> <p>Inactive ingredients citric acid anhydrous, D&C Red #33, edate disodium, ethylcellulose, FD&C Blue #1, flavor, high fructose corn syrup, methylparaben, polyethylene glycol 3350, polyorbital 80, propylene glycol, polypropylene, printed paper, sucrose, tragacanth, vegetable oil, xanthan gum.</p> <p>Questions? 1-866-682-4639</p>



© 2017 Distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76165
 021907
 400026

2 1/2"

3 3/4"

PHYSICIAN SAMPLE - NOT TO BE SOLD

Delsym
 dextromethorphan polistirex
 extended-release suspension

COUGH SUPPRESSANT
 15 mL (1/2 fl oz)

ACTIVE INGREDIENT
 (in each 5 mL teaspoonful)
 Dextromethorphan polistirex
 equivalent to 30 mg
 dextromethorphan hydrobromide..... suppressant

Purpose
 To temporarily relieve cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants.

Warnings: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (serotonin reuptake inhibitor, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Use: temporarily relieve cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants.

Warnings: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (serotonin reuptake inhibitor, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Use: temporarily relieve cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants.

Warnings: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (serotonin reuptake inhibitor, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Cover

Ask a doctor before use if you have chronic cough that lasts as long as 8 weeks, occurs with asthma or emphysema, cough that occurs with too much phlegm (mucus). Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
 Shake bottle well before use. Dose as follows or as directed by a doctor.

adults and children 12 years of age and over	2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
children 6 to under 12 years of age	1 teaspoonful every 12 hours, not to exceed 2 teaspoonfuls in 24 hours
children 2 to under 6 years of age	1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours
children under 2 years of age	consult a doctor

Back of Cover

PHYSICIAN SAMPLE - NOT TO BE SOLD

Delsym
 dextromethorphan polistirex
 extended-release suspension

COUGH SUPPRESSANT
 15 mL (1/2 fl oz)

Lot No.:
 Exp. Date:
 022207 4000514

Other information: each 5 mL teaspoonful contains sodium 5 mg. Store at 20°-25°C (68°-77°F).


Inactive ingredients: citric acid and anhydrous D&C Red #33, eucalyptol, ethylcellulose, FD&C Blue #1, flavor, high fructose corn syrup, methylparaben, polyethylene glycol, 3350, polyethylene glycol, propylparaben, purified water, sucrose, xanthan gum, vegetable oil, xanthan gum.

Questions? 1-866-682-4639

© 2007 Distributed by
 Adams Respiratory Therapeutics
 Fort Worth, TX 76155

Base

Hinge Area
 Reseal Area



FILE INFORMATION

Job Description: Delsym Grape 15mL Expanded Content Label

Drawing #: Not Available

Label Size: 4-1/16" w x 15/16" h

Software: Adobe Illustrator 12