

**RB Artwork and
 Print Specification**

RB514682
TR2479418
RBL2203003
G
 Delsym
 Adult
 Cough
 12 Hour Grape
 15 ml
 USA
 21/04/2023

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ shake bottle well before use
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information
 ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77° F)
 ■ dosing cup provided

Drug Facts (continued)
Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

DOSING CUP INCLUDED

Delsym
 10 mL
 5 mL
 2.5 mL
NOT ACTUAL SIZE

Contains No Fever Reducer or Pain Reliever
 Please visit our website: www.delsym.com

Delsym
 Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

COUGH
12 HOUR RELIEF

grape Flavored Liquid
 DAY or NIGHT
 ALCOHOL FREE
15 mL (1/2 fl oz)

Drug Facts
Active ingredient Purpose (in each 5 mL)
 Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 Ask a doctor before use if you have
 ■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema
 ■ cough that occurs with too much phlegm (mucus)
 Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

LOT NO.:
 EXP. DATE:

reckitt
 Dist by:
 RB Health (US)
 Parsippany, NJ
 07054-0224
 ©2022 RB Health
 C I A 000000
 PT# C I A 60346
 RB000000 042123

6 7/8

7 11/32

7 1/2

TAMPER EVIDENT: Do not use if the neckband printed with is broken or missing.

00000000

Delsym®
 NDC 63824-171-63
Delsym®
 Dextromethorphan Polistirex
 Extended-Release Suspension
 (Cough Suppressant)

Drug Facts
Active ingredient (in each 5 mL) Purpose
 Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 ■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema
 ■ cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information
 ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77° F) ■ dosing cup provided

Drug Facts (continued)
Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639

ALSO AVAILABLE IN ORANGE FLAVOR

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

Dist. by: RB Health (US) Parsippany, NJ 07054-0224 ©2022 RB Health
 C I A 000000 PT# C I A 60329 RB000000 042123

ALCOHOL FREE

grape Flavored Liquid
 DAY or NIGHT

COUGH 12 HOUR RELIEF

89 mL (3 fl oz)

DELSYM® DOSING
 SHAKE WELL BEFORE USE.
 Measure only with dosing cup provided. Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions
DOSING CUP INCLUDED

Delsym®
 10 mL
 5 mL
 2.5 mL

ACTUAL SIZE
 Contains no Fever Reducer or Pain Reliever
 Please visit our website: www.delsym.com


000000000000

Lot No.:
 Exp. Date:

RB Artwork and Print Specification


RB514508
 TR2478306
 RBL2202999
F
 Delsym
 Adult
 Cough
 12 Hour Grape
 3 fl oz
 USA
 21/04/2023

9 15/32




RB Artwork and Print Specification

RB515328
TR2462502
A
Delsym
Adult
Cough
12 Hour Grape
3 fl oz
USA
26/04/2022



RB Artwork and Print Specification

RBS14094
TR2476938
RBL2202984
H
Delsym
Adult
Cough
12 Hour Grape
3 fl oz
USA
16/03/2023



Drug Facts
Active ingredient (in each 5 mL) Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... Cough suppressant
Purpose
Uses temporarily relieves
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the impulse to cough to help you get to sleep
Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema
■ cough that occurs with too much phlegm (mucus)
Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ shake bottle well before use
■ measure only with dosing cup provided
■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor
■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information
■ each 5 mL contains: sodium 7 mg
■ store at 20-25°C (68-77° F) ■ dosing cup provided

Drug Facts (continued)
Inactive ingredients citric acid anhydrous, edetate disodium, ethylcellulose, FD&C yellow no. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum
Questions? 1-866-682-4639

DELSYM® DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided. Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions
DOSING CUP INCLUDED

Delsym®
10 mL
5 mL
2.5 mL
ACTUAL SIZE
Contains no Fever Reducer or Pain Reliever
Please visit our website: www.delsym.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Dist. by: RB Health (US) Parsippany, NJ 07054-0224 ©2023 RB Health
C I A 000000 PT# C I A 60329 RB000000 042123

00000000

0 000000 000000 0

Lot No.:
Exp. Date:

RB Artwork and Print Specification




RB514507
TR2478311
RBL2203000
G
Delsym
Adult
Cough
12 Hour Orange
3 fl oz
USA
21/04/2023

4⁵/₁₆

2³/₄ 1²¹/₃₂


RB Artwork and Print Specification

RB515320
TR2462496
A
 Delsym
 Adult
 Cough
 12 Hour Orange
 3 fl oz
 USA
 26/04/2022



RB Artwork and Print Specification

RB514098
TR2494983
RBL2202985
H
Delsym
Adult
Cough
12 Hour Orange
3 fl oz
USA
16/03/2023





RB Artwork and Print Specification

RB514681
TR2478315
RBL2203001
G
Delsym
Adult
Cough
12 Hour Grape
5 fl oz
USA
21/04/2023

Drug Facts

Active ingredient (in each 5 mL) Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... **Purpose** Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 ■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema
 ■ cough that occurs with too much phlegm (mucus)
Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77°F) ■ dosing cup provided


Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639

DELSYM DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided.
Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL Every 12 hours
6 to under 12	5 mL Every 12 hours
4 to under 6	2.5 mL Every 12 hours
Under 4	DO NOT USE

See back panel for full dosing directions
DOSING CUP INCLUDED



PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



©2023 RB Health
Dist. by: RB Health (US) Parsippany, NJ 07054-0224
C I A 000000
P T & C I A 60333
R B 000000 042123

grape
Flavored Liquid
DAY or NIGHT

ALCOHOL FREE
148 mL (5 fl oz)

ACTUAL SIZE
Contains No Fever Reducer or Pain Reliever
Please visit our website: www.delsym.com



Lot No.:
Exp. Date:

**RB Artwork and
Print Specification**

RB515332
TR2462522
A
Delsym
Adult
Cough
12 Hour Grape
5 fl oz
USA
20/04/2022

PEEL OPEN HERE TO READ COMPLETE WARNINGS AND INFORMATION ▶

TAMPER EVIDENT: Do not use if the neckband printed with is broken or missing.



Keep carton for full information.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2022 RB Health

Please visit our website: www.delsym.com

PT# C I A 60377
RB000000 042022

Lot: _____
Exp.: _____

C I A 0 0 0 0 0 0 0

2-1/4"

Drug Facts

Active ingredient (in each 5 mL) Purpose
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide...Cough suppressant

Uses: temporarily relieves: cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants; the impulse to cough to help you get to sleep.

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **Ask a doctor before use if you have** chronic cough that lasts such as occurs with smoking, asthma, or emphysema; cough that occurs with too much phlegm (mucus). **Stop use and ask a doctor** if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Shake bottle well before use. Measure only with dosing cup provided. Do not use dosing cup with other products. Dose as follows or as directed by a doctor.

2-7/8"

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information:
Each 5 mL contains: **sodium 7 mg.**
Store at 20-25°C (68-77°F). Dosing cup provided.

Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639
You may also report side effects to this phone number.

RB Artwork and Print Specification

RB514117
TR2476952
RBL2202986
G
Delsym
Adult
Cough
12 Hour Grape
5 fl oz
USA
14/03/2023



8 7/8

5 11/16

Delsym[®]

Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

NDC 63824-175-65

Drug Facts

Active ingredient (in each 5 mL) Purpose
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... Cough suppressant

Uses temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have

- chronic cough that lasts such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use

- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77° F) ■ dosing cup provided

Inactive ingredients citric acid anhydrous, edetate disodium, ethylcellulose, FD&C yellow no. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639

Delsym[®]

Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

COUGH

12 HOUR RELIEF

orange
Flavored Liquid
DAY or NIGHT

ALCOHOL FREE

148 mL (5 fl oz)

Delsym[®]

Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

DELSYM DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided.
Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL Every 12 hours
6 to under 12	5 mL Every 12 hours
4 to under 6	2.5 mL Every 12 hours
Under 4	DO NOT USE

See back panel for full dosing directions

ALSO AVAILABLE IN GRAPE FLAVOR

Dosing Cup Included



ACTUAL SIZE

Contains No Fever Reducer or Pain Reliever
Please visit our website: www.delsym.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

© 2023 RB Health
Dist. by: RB Health (US) C I A 000000
Parolippany, NJ PTF# C I A 60333
07054-0224 3233999 061323

reckitt

Lot No.:
Exp. Date:

RB Artwork and Print Specification




RB514668
TR2893441
RBL2304188
B
Delsym
Adult
Cough
12 Hour Orange
5 fl oz
USA
13/06/2023

9 15/32



**RB Artwork and
Print Specification**

RB515329
TR2462517
B
Delsym
Adult
Cough
12 Hour Orange
5 fl oz
USA
13/05/2022



**RB Artwork and
Print Specification**

RB514116
TR2496484
RBL2202987
G
Delsym
Adult
Cough
12 Hour Orange
5 fl oz
USA
14/03/2023



9⁶³/₆₄



Twin Pack
2 x 5 fl oz Bottles
GRAPE FLAVORED LIQUID

TAMPER EVIDENT: Do not use if the neckband printed with is broken or missing.

Drug Facts

Active ingredient (In each 5 mL)
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide... Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 Ask a doctor before use if you have
 ■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema
 ■ cough that occurs with too much phlegm (mucus)
 Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ shake bottle well before use
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information
 ■ each 5 mL contains sodium 7 mg
 ■ store at 20-25°C (68-77° F)
 ■ dosing cup provided

Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639
Please visit our website: www.delsym.com

NDC 63824-171-22



GRAPE FLAVORED LIQUID

DAY OR NIGHT

Twin Pack
For Adults & Children
AGES 4+
GRAPE FLAVORED LIQUID

Twin Pack
For Adults & Children
AGES 4+
GRAPE FLAVORED LIQUID

COUGH



12 HOUR RELIEF

ALCOHOL FREE

TWO - 148 mL (5 fl oz) Bottles TOTAL - 296 mL (10 fl oz)



DELSYM® DOSING
SHAKE WELL BEFORE USE. Measure only with dosing cup provided. Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing instructions
DOSING CUP INCLUDED



ACTUAL SIZE
Contains no Fever Reducer or Pain Reliever

ALSO AVAILABLE IN ORANGE FLAVOR

reckitt


Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2022 RB Health 00000000 042/23

RB Artwork and Print Specification

RB537209
TR2616547
RBL2209707
F
Delsym
Adult
Cough
12 Hour Grape
2 x 5 fl oz
USA
24/04/2023

15¹⁵/₃₂

9⁶³/₆₄



Twin Pack
2 x 5 fl oz Bottles
ORANGE FLAVORED LIQUID

TAMPER EVIDENT: Do not use if the neckband printed with is broken or missing.

Drug Facts

Active ingredient (in each 5 mL)
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide... Cough suppressant

Purpose
Cough suppressant

Uses
temporarily relieves
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema
■ cough that occurs with too much phlegm (mucus)
Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ shake bottle well before use
■ measure only with dosing cup provided
■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor
■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information
■ each 5 mL contains sodium 7 mg
■ store at 20-25°C (68-77°F)
■ dosing cup provided

Inactive ingredients citric acid anhydrous, edetate disodium, ethylcellulose, FD&C yellow no. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639
Please visit our website: www.delsym.com

NDC 63824-175-22



DAY OR NIGHT


Twin Pack
For Adults & Children
AGES 4+
ORANGE FLAVORED LIQUID

Twin Pack
For Adults & Children
AGES 4+
ORANGE FLAVORED LIQUID

DELSYM® DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided.
Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing instructions
DOSING CUP INCLUDED



ACTUAL SIZE
Contains no Fever Reducer or Pain Reliever

ALSO AVAILABLE IN **GRAPE FLAVOR**

reckitt
Dist. by: RB Health (US)
Parippany, NJ 07654-0224
©2023 RB Health 00000000 042423

Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

COUGH

12 HOUR RELIEF

ALCOHOL FREE

TWO - 148 mL (5 fl oz) Bottles TOTAL - 296 mL (10 fl oz)


Lot No.:
Exp. Date:

0 00000 00000 0

RB Artwork and Print Specification

RB537039
TR2610668
RBL2209706
F
Delsym
Adult
Cough
12 Hour Orange
2 x 5 fl oz
USA
24/04/2023

15¹⁵/₃₂



RB Artwork and Print Specification

RB514511
 TR2490427
 RBL2203004
G
 Delsym
 Children's
 Cough
 12 Hour Grape
 3 fl oz
 USA
 21/04/2023

Drug Facts

Active ingredient (in each 5 mL)
 Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... Cough suppressant

Purpose
 Cough suppressant

Uses temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- chronic cough that lasts such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use

- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 7 mg
- dosing cup provided

Drug Facts (continued)

Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639



PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

Dist. by:
 RB Health (US)
 Parsippany,
 NJ 07054-0224
 ©2023 RB Health

C I A 000000
 PT# C I A 60329
 RB000000 042123

children's
Delsym[®]
 Dextromethorphan Polistirex
 Extended-Release Suspension
 (Cough Suppressant)

AGES
4+

COUGH

12 HOUR RELIEF

grape
 Flavored Liquid
 For Children & Adults

DAY or NIGHT
 ALCOHOL FREE


89 mL (3 fl oz)

children's
Delsym[®]
 Dextromethorphan Polistirex
 Extended-Release Suspension
 (Cough Suppressant)

DELSYM[®] DOSING
 SHAKE WELL BEFORE USE.
 Measure only with dosing cup provided.
 Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions
DOSING CUP INCLUDED



4 5/16




1 11/16

2 3/4

Lot No.:
 Exp. Date:


RB Artwork and Print Specification

RB517704
TR2476874
A
 Delsym
 Children's
 Cough
 12 Hour Grape
 3 fl oz
 USA
 26/04/2022



**RB Artwork and
Print Specification**

RB514120
TR2489256
RBL2202993
H
Delsym
Children's
Cough
12 Hour Grape
3 fl oz
USA
16/03/2023





RB Artwork and Print Specification

RB514509
 TR2490410
 RBL2203005
G
 Delsym
 Children's
 Cough
 12 Hour Orange
 3 fl oz
 USA
 21/04/2023

Drug Facts

Active ingredient (in each 5 mL) Purpose
 Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 ■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema
 ■ cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77°F) ■ dosing cup provided

Drug Facts (continued)

Inactive ingredients citric acid anhydrous, edetate disodium, ethylcellulose, FD&C yellow no. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639



PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

Dist. by:
 RB Health (US)
 Parsippany,
 NJ 07054 0224
 ©2023 RB Health

C I A 000000
 PT# C I A 60328
 RB000000 042123

NDC 63824-177-63

children's
Delsym[®]

Dextromethorphan Polistirex
 Extended-Release Suspension
 (Cough Suppressant)

AGES
4+

COUGH

12 HOUR RELIEF

orange

Flavored Liquid
 For Children & Adults

DAY or NIGHT
 ALCOHOL FREE

89 mL (3 fl oz)

children's
Delsym[®]

Dextromethorphan Polistirex
 Extended-Release Suspension
 (Cough Suppressant)

DELSYM[®] DOSING

SHAKE WELL BEFORE USE.
 Measure only with dosing cup provided.
 Do not use dosing cup with other products.

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions
DOSING CUP INCLUDED



ACTUAL SIZE
 Contains no Fever Reducer or Pain Reliever
 Please visit our website: www.delsym.com




4 5/16

2 3/4

1 21/32


**RB Artwork and
 Print Specification**

RB517697
TR2476856
A
 Delsym
 Children's
 Cough
 12 Hour Orange
 3 fl oz
 USA
 26/04/2022



RB Artwork and Print Specification

RB514119
TR2489236
RBL2202994
H
Delsym
Children's
Cough
12 Hour Orange
3 fl oz
USA
16/03/2023



8 7/8



00000000

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... Cough suppressant

Uses temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have

- chronic cough that lasts such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use

- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77°F) ■ dosing cup provided

Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639



ALSO AVAILABLE IN
ORANGE
FLAVOR

children's
Delsym[®]

NDC 63824-173-65

children's
Delsym[®]

Dextromethorphan Polistirex
Extended-Release Suspension
(Cough Suppressant)

AGES
4+

COUGH



12 HOUR RELIEF

DAY or NIGHT
ALCOHOL FREE

grape
Flavored Liquid
For Children & Adults

148 mL (5 fl oz)



Delsym[®]

10 mL
5 mL
2.5 mL

ACTUAL SIZE

Contains No Fever Reducer or Pain Reliever
Please visit our website: www.delsym.com



0 00000000000 0

Lot No.:
Exp. Date:

RB Artwork and Print Specification

RB514674
TR2490449
RBL2203006
G
 Delsym
 Children's
 Cough
 12 Hour Grape
 5 fl oz
 USA
 21/04/2023

9 15/32


**RB Artwork and
 Print Specification**

RB517708
TR2476894
A
 Delsym
 Children's
 Cough
 12 Hour Grape
 5 fl oz
 USA
 20/04/2022

2-1/4"

PEEL OPEN HERE TO READ COMPLETE WARNINGS AND INFORMATION ▶

**TAMPER EVIDENT: Do not use
 if the neckband printed with
 is broken or missing.**



0 000000 000000 0

Keep carton for full information.

Dist. by: RB Health (US)
 Parsippany, NJ 07054-0224
 ©2022 RB Health

Please visit our website: www.delsym.com

PT# C | A 60377
 RB000000 042022

Lot:
 Exp:

C I A 0 0 0 0 0 0 0

2-1/4"

Drug Facts

Active ingredient (in each 5 mL) Purpose
 Dextromethorphan polistirex equivalent to
 30 mg dextromethorphan hydrobromide.....Cough suppressant

Uses: temporarily relieves: cough due to minor throat and
 bronchial irritation as may occur with the common cold or
 inhaled irritants; the impulse to cough to help you get to
 sleep.

Warnings
Do not use if you are now taking a prescription monoamine
 oxidase inhibitor (MAOI) (certain drugs for depression,
 psychiatric, or emotional conditions, or Parkinson's
 disease), or for 2 weeks after stopping the MAOI drug. If
 you do not know if your prescription drug contains an MAOI,
 ask a doctor or pharmacist before taking this product. **Ask**
a doctor before use if you have chronic cough that lasts
 such as occurs with smoking, asthma, or emphysema;
 cough that occurs with too much phlegm (mucus). **Stop**
use and ask a doctor if cough lasts more than 7 days,
 comes back, or occurs with fever, rash, or headache that
 lasts. These could be signs of a serious condition. **If**
pregnant or breast-feeding, ask a health professional
 before use. **Keep out of reach of children.** In case of
 overdose, get medical help or contact a Poison Control
 Center right away.

Directions: Shake bottle well before use. Measure
 only with dosing cup provided. Do not use dosing cup with
 other products. Dose as follows or as directed by a doctor.

2-7/8"

2-1/4"

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information:
 Each 5 mL contains: sodium 7 mg.
 Store at 20-25°C (68-77°F). Dosing cup provided.

Inactive ingredients citric acid anhydrous, D&C red
 no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1,
 flavor, high fructose corn syrup, methylparaben, partially
 hydrogenated vegetable oil (soybean, cottonseed), polyethylene
 glycol 3350, polysorbate 80, propylene glycol, propylparaben,
 purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639
 You may also report side effects to this phone number.

2-7/8"

RB Artwork and Print Specification

RB514123
TR2489261
RBL2202995
G
Delsym
Children's
Cough
12 Hour Grape
5 fl oz
USA
14/03/2023



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
03/15/2024 09:57:45 AM