

BACK



AREA FOR LOT AND EXP
 TO BE PRINTED IN DARK COLOR
 26 mm x 5.5 mm

SEAL AREA FOR LOT & EXP
 25 mm x 4.5 mm

000067194

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.75 pt
• SUBHEADINGS/BODY TEXT	4.75 pt
• LEADING	4.75 pt
• BULLETS	3.75 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	N/A

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID)*Fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> headache toothache backache menstrual cramps the common cold muscular aches minor pain of arthritis temporarily reduces fever <p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> hives facial swelling asthma (wheezing) shock skin reddening rash blisters 	<p>Purpose Pain reliever/ Fever reducer</p> <p>Drug Facts (continued)</p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>
--	--

PANEL 1

<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> under a doctor's care for any serious condition taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug <p>When using this product</p> <ul style="list-style-type: none"> take with food or milk if stomach upset occurs 	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> feel faint vomit blood have bloody or black stools have stomach pain that does not get better you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> chest pain trouble breathing weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear
---	--

PANEL 2


<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed the smallest effective dose should be used adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 tablet, 2 tablets may be used do not exceed 6 tablets in 24 hours, unless directed by a doctor children under 12 years: ask a doctor 	<p>Drug Facts (continued)</p> <p>Other information</p> <ul style="list-style-type: none"> read all warnings and directions before use. Keep carton. store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F) <p>Inactive ingredients</p> <p>acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax</p> <p>Questions or comments? call toll free 1-800-88-ADVIL</p>
--	--

000067196


BASE PANEL

000067196

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



UNVARNISHED AREA
FOR LOT & EXP. DATE




000067197

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces




FRONT




BACK

000067198

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces




READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION



FPO
3 0573 0151 10 8

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Ibuprofen 200 mg (NSAID)*Pain reliever/Fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever <p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters</p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug 	<p>Drug Facts (continued)</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ take with food or milk if stomach upset occurs <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ read all warnings and directions before use. Keep card. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F) <p>Inactive ingredients</p> <p>acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, methylparaben, microcrystalline cellulose, starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax</p> <p>Questions or comments? call toll free 1-800-88-ADVIL</p>
--	---

For most recent product information, visit www.Advil.com
 Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor
 Trademarks owned or licensed by GSK.



000070156


↑
 LOT & EXP AREA

000070156

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION



FPO
3 0573 0151 20 7

Drug Facts

Active ingredient (in each tablet) Purpose
 Ibuprofen 200 mg (NSAID) Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

For most recent product information, visit www.Advil.com
 Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor
 Trademarks owned or licensed by GSK

Drug Facts (continued)

When using this product

- take with food or milk. 1 stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use. Keep card.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
 acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, polyethylene glycol, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments? call toll free 1-800-88-ADVIL

000070157

UNPRINTED/UNVARNISHED
 AREA FOR LOT & EXP

000070157

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Reference ID: 4903919

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts
Active ingredient (in each tablet): Ibuprofen 200 mg (NSAID) *Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Uses

- Temporarily relieves minor aches and pains due to:
 - backache
 - headache
 - muscular cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- Temporarily reduces fever

Warnings
Allergy: A severe allergic reaction may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- swollen lymph nodes
- skin redness
- rash
- itching

If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or nonsteroidal NSAID, aspirin, ibuprofen, naproxen, or steroid drug
- take other drugs containing aspirin or ibuprofen
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risks are higher if you use more than directed or for longer than directed.

Do not use: If you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as indigestion
- you have high blood pressure, heart disease, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin or heart attack or stroke medicine because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs

000070104

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

50% MORE FREE

12 Free Tablets

Advil
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

24 36
Coated Tablets

Advil
Tablets


000070104

	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Advil

**Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)**

50 Coated Tablets



Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts
Active ingredient (in each tablet) **Purpose**
Ibuprofen (200 mg (NSAID)) Pain reliever/fever reducer
Hydrocodone bitartrate Pain reliever/fever reducer

Uses
Temporarily relieves minor aches and pains due to:

- headache
- backache
- muscle cramps
- the common cold
- menstrual aches
- minor pain of arthritis
- temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using the product
- take more or for a longer time than directed

Drug Facts (continued)
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risks are higher if you use more than directed or for longer than directed.
Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as indigestion
- you have high blood pressure, heart disease, liver problems, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart disease or stroke, because ibuprofen may increase this benefit of aspirin
- taking any other drug

When using the product

- take with food or milk, if stomach upset occurs
- stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
 - red stool
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better

Drug Facts (continued)

- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest amount to be used should be used
- take with food or milk
- take 1 tablet every 4 to 6 hours as needed for pain or fever
- do not exceed 1 tablet (200 mg) ibuprofen in 24 hours unless directed by a doctor
- children under 12 years, ask a doctor


Other information

- read all warnings and directions before use. Keep carton.
- store at 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000070158



3 0573 0150 30 9

Drug Facts (continued)
Do Not Use if seal under bottle cap is impr or broken or missing.
Questions or comments?
 For most recent product information, visit www.Advil.com
 Dist. by GSK CH, Warren, NJ 07059
 ©2021 GSK or license or
 Trademarks owned or licensed by GSK

	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

100 Coated Tablets

Advil

**Ibuprofen Tablets, 200 mg
Pain Reliever/ Fever Reducer (NSAID)**

100 Coated Tablets

Advil
Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts
(in each tablet)

Active ingredient
Ibuprofen 200 mg (NSAID)*.....Fever reducer
nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/
fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - the common cold
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: The product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - faint pain
 - vomit blood
 - have bloody or black stools
- your stomach pain that does not get better
- you have symptoms of heart problems or stroke
- chest pain
- trouble breathing
- weakness in one part of your body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 3 days
- ever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Do Not Use If seal and/or dotto cap impaled with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information, visit www.Advil.com.
©2021 GSK or its affiliates.
Distributed by GSKCH, Warren, NJ 07069.
*FDA approved and licensed by GSK

UNPRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000070106

3 0573 0150 40 8

FPO

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

000070106

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Advil

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

200 Coated Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts
(in each tablet)

Active ingredient (in each tablet) **Purpose**
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - muscle aches
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Always alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- rash
- skin redness/itching
- dizziness

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take aspirin or other drugs
- use alcohol
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or other NSAIDs)
- take 3 or more alcoholic drinks every day while using this product.

Use more or for longer time than directed

Heart attack and stroke warning, NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin or heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product:

- be with food or milk if stomach upset occurs

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - bad acid
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - big swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Advil

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

200 Coated Tablets

Do Not Use If seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information, visit www.advil.com
Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or Hanser
Trademarks owned or licensed by GSK

000070105

3 0573 0154 75 8


FPO

PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

UNPRINTED AREA FOR
DATE & LOT LASER CODING
(UNVARNISHED)

000070105

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



LASER ETCHING AREA FOR LOT & EXP.
PRINT A SOLID DARK COLOR.

DRUG FACTS TEXT DEFINED	TYPE SIZE
* DRUG FACTS TITLE	9 pt
* DRUG FACTS CONTINUED	8 pt
* HEADINGS	8 pt
* SUBHEADINGS/BODY TEXT	6 pt
* LEADING	6.5 pt
* # OF CHARACTERS PER INCH	<39
* BULLETS	5 pt
* SPACE BEFORE BULLET	2 ems
* BARLINES, HAIRLINES	1.5 pt, .5 pt
* SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP
PANEL 1

Easy Open ARTHRITIS CAP

Arthritis Foundation
Ease of Use
CAP

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Advil
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

200 Coated Tablets

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Drug Facts

Active ingredient (in each tablet)
Ibuprofen 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/
Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

000067200

LIFT HERE for More Drug Facts

FPO

3 0573 0154 21 5

FOR LOT & EXPIRATION AREA.
UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE.

BASE
PANEL 3

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use.
- store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Drug Facts (continued)

Questions or comments? call toll free 1-800-88-ADVIL

Advil

For most recent product information, visit www.Advil.com
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2021 GSK group of companies or its licensor
Trademarks are owned by or licensed to the GSK group of companies

RELEASED BACK
PANEL 2

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product! ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body
 - slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area ■ any new symptoms appear

000067200

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LAYER FACE

Advil
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Advil
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

000069978

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

3/16" X 3/16" DATA MATRIX CODE (ECC 200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND WITH A MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.

UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

Drug Facts

Active ingredient (in each tablet)	Purpose
Ibuprofen 200 mg (NSAID)*	Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

3 0573 0154 98 7

TOP LAYER BACK

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

NO COPY HINGED AREA

BASE LAYER

Advil

For most recent product information, visit www.Advil.com

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2021 GSK group of companies or its licensor
Trademarks are owned by or licensed to the GSK group of companies

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments?
call toll free 1-800-88-ADVIL

NO COPY HINGED AREA

NO COPY AREA

000069978

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LAYER FACE

Advil
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Advil
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

000069980

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

3/16" X 3/16" DATA MATRIX CODE (ECC 200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND WITH A MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES

UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

Drug Facts

Active ingredient (in each tablet)	Purpose
Ibuprofen 200 mg (NSAID)*	Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

FPO
3 0573 0154 60 4

TOP LAYER BACK

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Drug Facts (continued)

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

NO COPY HINGED AREA

BASE LAYER

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments?
call toll free 1-800-88-ADVIL

Advil

For most recent product information, visit www.Advil.com

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2021 GSK group of companies or its licensor
Trademarks are owned by or licensed to the GSK group of companies

NO COPY HINGED AREA

NO COPY AREA

000069980

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Advil

**Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)**

24 Coated Gel Caplets*
*Capsule-Shaped Gelatin-Coated Tablets



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts
(in each gel caplet) Purpose: Ibuprofen 200 mg (NSAID)*. Pain reliever/fever reducer/anti-inflammatory drug.

Uses
Temporarily relieves minor aches and pains due to:

- headache
- toothache
- muscle aches
- nerve pain
- menstrual cramps
- the common cold
- muscular aches
- minor pain or arthritis
- temporal headaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- itching
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)
Some health warnings: This product contains an NSAID, which may cause severe stomach bleeding, the more you take. It may also cause stomach bleeding, the more you take. It may also cause stomach bleeding, the more you take. It may also cause stomach bleeding, the more you take.

Do not use if you have ever had an allergic reaction to any of the ingredients listed above, or if you have a history of stomach pain, ulcers, or bleeding in the stomach or intestines. Do not use if you have a history of stomach pain, ulcers, or bleeding in the stomach or intestines.

Ask a doctor before use if you have problems or stomach side effects from taking pain relievers or fever reducers, or you have a history of stomach problems, such as heartburn, indigestion, or ulcers. Do not use if you have a history of stomach pain, ulcers, or bleeding in the stomach or intestines.

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin, taking any other drug, or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Drug Facts (continued)
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- stomach pain that does not get better
- black or bloody stools
- dark or bloody urine
- stomach pain that does not get better
- stomach pain that does not get better
- stomach pain that does not get better

Directions Do not take more than directed. The smallest amount should be used. Adults and children 12 years and over: take 1 gel caplet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used. Do not exceed 6 gel caplets in 24 hours, unless directed by a doctor. Children under 12 years: ask a doctor.

Other information read all warnings and directions before use. Keep carton.

000070115

3 0573 0165 20 4

FPO

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000070115	
DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
12/14/2021 09:29:25 AM