

**Drug Facts (continued)**

**If pregnant or breast feeding, ask a health professional before use.** It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed

adults and children 12 years and older	<ul style="list-style-type: none"> <li>take 1 caplet every 4 to 6 hours while symptoms persist</li> <li>if pain or fever does not respond to 1 caplet, 2 caplets may be used, but do not exceed 6 caplets in 24 hours, unless directed by a doctor</li> <li>the smallest effective dose should be used</li> </ul>
children under 12 years	ask a doctor

**Other information**  
 ■ do not use if container is open or if tear strip is broken or missing  
 ■ store between 20 - 25°C (68 - 77°F).  
 ■ see container label for lot and expiration date

**Inactive ingredients**  
 camuba wax, corn starch, FD&C Yellow #6, hydroxypropyl methylcellulose, iron oxide, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide

**Questions or comments?**  
 Call toll free 1-800-962-5357

**McNeil** XXXXXXXX  
 McNeil Consumer Healthcare,  
 DIVISION OF MCNEIL-PPC, INC.,  
 FORT WASHINGTON, PA 19034 USA  
 © MCN-PPC, Inc. '01 [www.motrin.com](http://www.motrin.com)

EXP. 0 00000 00000 0

**Motrin**  
 IB Pain Reliever/  
 Fever Reducer  
 8 Caplets

**Drug Facts**

Active ingredient (in each caplet)	Purposes
ibuprofen 200 mg.....	Pain reliever/ fever reducer

**Uses** temporarily relieves minor aches and pains due to:

<ul style="list-style-type: none"> <li>headache</li> <li>minor pain of arthritis</li> <li>backache</li> <li>menstrual cramps</li> <li>reduces fever</li> </ul>	<ul style="list-style-type: none"> <li>muscular aches</li> <li>toothache</li> <li>the common cold</li> </ul>
--	--

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:  
 ■ hives ■ facial swelling  
 ■ asthma (wheezing) ■ shock

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

**Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer**

**Ask a doctor before use if you have**  
 ■ stomach pain  
 ■ problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition  
 ■ taking any other drug  
 ■ taking any other product that contains ibuprofen, or any other pain reliever/fever reducer

**When using this product give with food or milk if stomach upset occurs**

**Stop use and ask a doctor if**  
 ■ an allergic reaction occurs. Seek medical help right away.  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ stomach pain or upset gets worse or lasts  
 ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear

**Labeling Format Information:**

<b>Fonts:</b> Helvetica regular, bold and bold italic.	
<b>Drug Facts:</b> 8.5 pt	<b>Leading:</b> 6.5 pt
<b>Header:</b> 8 pt	<b>Bullets:</b> 5 pt
<b>Subheader:</b> 6 pt	<b>Barlines:</b> 2.5 pt
<b>Body Text:</b> 6 pt	<b>Hairlines:</b> 0.5 pt
<b>Drug Facts (continued):</b> 8 pt	

Horizontal Scale: 94-100%      Average Kerning: 0

**Drug Facts (continued)**

**If pregnant or breast feeding, ask a health professional before use.** It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

■ do not take more than directed

adults and children 12 years and older

■ take 1 caplet every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used, but do not exceed 6 caplets in 24 hours, unless directed by a doctor  
■ the smallest effective dose should be used

children under 12 years

■ ask a doctor

**Other information**

do not use if container is open or if tear strip is broken or missing  
store between 20 - 25 C (68 - 77 F).  
see container label for lot and expiration date

**Inactive ingredients**

cambea wax, corn starch, FD&C Yellow #6, hydroxypropyl methylcellulose, iron oxide, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide

**Questions or comments?**

Call toll free 1-800-962-5357



McNeil Consumer Healthcare,  
DIVISION OF MCNEIL-PPC, INC.,  
FORT WASHINGTON, PA 19034 USA  
© MCN-PPC, Inc. '01

XXXXXXXX

[www.motrin.com](http://www.motrin.com)

EXP.



0 00000-000-00 0

**Motrin**  
IB Pain Reliever/  
Fever Reducer

8

**Drug Facts**

Active Ingredient (in each caplet)	Purposes
Ibuprofen 200 mg.....	Pain reliever/ fever reducer

**Uses** temporarily relieves minor aches and pains due to:

- headache
- minor pain of arthritis
- backache
- menstrual cramps
- reduces fever
- muscular aches
- toothache
- the common cold

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- facial swelling
- shock

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.

**Ask a doctor before use if you have**

- stomach pain
- problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains ibuprofen, or any other pain reliever/fever reducer

**When using this product give with food or milk if stomach upset occurs**

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

↓

---

**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**

---

/s/

-----  
Charles Ganley  
9/7/01 06:05:31 PM