

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) --- NO PRINT
- 5) [Hatched Box] LOT & EXP. DATE AREA (NO ADDITIONAL COPY, NO COLOR)
- 6) [Cross-hatched Box] UPC AREA (NO COLOR)
- 7) [Diagonal Hatched Box] GLUE AREA (NO COPY, NO COLOR, NO VARNISH)

(b) (4)	7	6/16/97	ADDED IV INFO	ING. NO.	 CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA. 19034 TITLE: CARTON, SEAL END 1.50 L X 1.44 W X 2.63 D	
	6	3-15-94	ADDED 'EXP.' TEXT	FC-C-111		
	5	10-18-93	REVISED NO VARNISH AREA	DRAWN		(b) (4)
	4	9-1-93	ADDED UPC AREA	PLT		NONE
	3	7-1-92	REDRAWN ON NEW FORMAT	SCALE		NONE
	2	6-6-88	ING. FLIPPED AS LH CARTON	DATE		8-31-87
	1	3-21-88	NOTE ADDED PRINT SIDE UP	REVISIONS		
DET.	NO.	DATE	REVISIONS			

00000000

Drug Facts (continued)

Do not use if you have ever had allergic reaction to any other pain reliever/fever reducer. Do not use if you have asthma, hives, itching, or other allergic reactions.

Warnings:

- Do not take if you are taking any other NSAID or aspirin.
- Do not take if you are taking any other blood thinners.
- Do not take if you are taking any other drugs that may increase the risk of bleeding.
- Do not take if you are taking any other drugs that may increase the risk of kidney problems.
- Do not take if you are taking any other drugs that may increase the risk of liver problems.
- Do not take if you are taking any other drugs that may increase the risk of stomach or intestinal problems.
- Do not take if you are taking any other drugs that may increase the risk of heart problems.
- Do not take if you are taking any other drugs that may increase the risk of high blood pressure.
- Do not take if you are taking any other drugs that may increase the risk of diabetes.
- Do not take if you are taking any other drugs that may increase the risk of low blood sodium.
- Do not take if you are taking any other drugs that may increase the risk of dizziness or lightheadedness.
- Do not take if you are taking any other drugs that may increase the risk of ringing in the ears.
- Do not take if you are taking any other drugs that may increase the risk of blurred vision.
- Do not take if you are taking any other drugs that may increase the risk of drowsiness or dizziness.
- Do not take if you are taking any other drugs that may increase the risk of constipation.
- Do not take if you are taking any other drugs that may increase the risk of diarrhea.
- Do not take if you are taking any other drugs that may increase the risk of nausea or vomiting.
- Do not take if you are taking any other drugs that may increase the risk of indigestion or heartburn.
- Do not take if you are taking any other drugs that may increase the risk of stomach pain or discomfort.
- Do not take if you are taking any other drugs that may increase the risk of bloating or gas.
- Do not take if you are taking any other drugs that may increase the risk of constipation.
- Do not take if you are taking any other drugs that may increase the risk of diarrhea.
- Do not take if you are taking any other drugs that may increase the risk of nausea or vomiting.
- Do not take if you are taking any other drugs that may increase the risk of indigestion or heartburn.
- Do not take if you are taking any other drugs that may increase the risk of stomach pain or discomfort.
- Do not take if you are taking any other drugs that may increase the risk of bloating or gas.

See New Warnings Information

NDC 50580-109-02

Motrin^{IB}

Ibuprofen Tablets usp 200 mg

Pain Reliever / Fever Reducer (NSAID)

24 Coated Tablets

Drug Facts (continued)

Warnings:

- Do not take if you are taking any other NSAID or aspirin.
- Do not take if you are taking any other blood thinners.
- Do not take if you are taking any other drugs that may increase the risk of bleeding.
- Do not take if you are taking any other drugs that may increase the risk of kidney problems.
- Do not take if you are taking any other drugs that may increase the risk of liver problems.
- Do not take if you are taking any other drugs that may increase the risk of stomach or intestinal problems.
- Do not take if you are taking any other drugs that may increase the risk of heart problems.
- Do not take if you are taking any other drugs that may increase the risk of high blood pressure.
- Do not take if you are taking any other drugs that may increase the risk of diabetes.
- Do not take if you are taking any other drugs that may increase the risk of low blood sodium.
- Do not take if you are taking any other drugs that may increase the risk of dizziness or lightheadedness.
- Do not take if you are taking any other drugs that may increase the risk of ringing in the ears.
- Do not take if you are taking any other drugs that may increase the risk of blurred vision.
- Do not take if you are taking any other drugs that may increase the risk of drowsiness or dizziness.
- Do not take if you are taking any other drugs that may increase the risk of constipation.
- Do not take if you are taking any other drugs that may increase the risk of diarrhea.
- Do not take if you are taking any other drugs that may increase the risk of nausea or vomiting.
- Do not take if you are taking any other drugs that may increase the risk of indigestion or heartburn.
- Do not take if you are taking any other drugs that may increase the risk of stomach pain or discomfort.
- Do not take if you are taking any other drugs that may increase the risk of bloating or gas.



00000000

<p>Description: Ctn Motrin IB Tab 24 (LP Bulk)</p> <p>File #: FDA3009897 Version: A-1</p> <p>Date: 08/27/2010 Contact: (b) (4)</p>	<p>Labeling Format Information:</p> <p>Fonts: Mcn DF Condensed, Bold, Bold Italic</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td>Drug Facts:</td> <td>8.25 pt</td> <td>Leading:</td> <td>6.00 pt</td> </tr> <tr> <td>Header:</td> <td>7.00 pt</td> <td>Bullets:</td> <td>5.00 pt</td> </tr> <tr> <td>Subheader:</td> <td>6.00 pt</td> <td>Barlines:</td> <td>2.50 pt</td> </tr> <tr> <td>Body Text:</td> <td>6.00 pt</td> <td>Hairlines:</td> <td>0.50 pt</td> </tr> <tr> <td>Drug Facts (continued):</td> <td>7.00 pt</td> <td></td> <td></td> </tr> </table> <p>Horizontal Scale: 70% Average Kerning: 0</p>	Drug Facts:	8.25 pt	Leading:	6.00 pt	Header:	7.00 pt	Bullets:	5.00 pt	Subheader:	6.00 pt	Barlines:	2.50 pt	Body Text:	6.00 pt	Hairlines:	0.50 pt	Drug Facts (continued):	7.00 pt		
Drug Facts:	8.25 pt	Leading:	6.00 pt																		
Header:	7.00 pt	Bullets:	5.00 pt																		
Subheader:	6.00 pt	Barlines:	2.50 pt																		
Body Text:	6.00 pt	Hairlines:	0.50 pt																		
Drug Facts (continued):	7.00 pt																				

Submission Artwork


Consumer Healthcare

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5)  LOT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6)  VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

INIT.	NO.	DATE	REVISIONS	DWG. NO.	DATE
(b) (4)	7	1/7/99	ADDED LOT/EXP AREA	L.B.L.43	
	6	6/16/97	ADDED DV INFO.		
	5	10-24-94	CHANGED NO PRINT AREA TO 1/16		
	4	3-15-94	ADDED "EXP. DATE" TEXT		
	3	5-11-92	REDRAWN ON NEW FORMAT		
	2	11-21-89	LENGTH & WIDTH CHANGED		
	1	11-15-89	REVISION OF NO-PRINT AREA		
				DATE	11-15-89

McNEIL
CONSUMER PRODUCTS CO.
CAMP HILL ROAD, FORT WASHINGTON, PA 19084 USA
TITLE:
HA LABEL
4.25 L X 1.25 W



Do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing.

Motrin[®] IB
Ibuprofen Tablets usp. 200 mg
Pain Reliever/Fever Reducer (NSAID)
24 Coated Tablets

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)*Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever
Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear


Directions ■ do not take more than directed ■ the smallest effective dose should be used
Adults and children 12 years and older: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
Children under 12 years: ■ ask a doctor. **Other information** ■ store between 20 - 25°C (68 - 77°F) **7844475**

Dist. by: McNeil-PPC, Inc., Fort Washington, PA 19084 USA
© McNeil-PPC, Inc. 2009 www.motrin.com
EXP. DATE: LOT:

200% OF SIZE



07844475



Do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing.

Motrin[®] IB
Ibuprofen Tablets usp. 200 mg
Pain Reliever/Fever Reducer (NSAID)
24 Coated Tablets

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)*Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever
Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Directions ■ do not take more than directed ■ the smallest effective dose should be used
Adults and children 12 years and older: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
Children under 12 years: ■ ask a doctor. **Other information** ■ store between 20 - 25°C (68 - 77°F) **7844475**

Dist. by: McNeil-PPC, Inc., Fort Washington, PA 19084 USA
© McNeil-PPC, Inc. 2009 www.motrin.com
EXP. DATE: LOT:

ACTUAL SIZE

Description: Lbl Motrin IB Tablet 24 - (b) (4)
File #: 3009897 Version: F-2
Date: 2/4/2010 Contact: (b) (4)

Labeling Format Information:





Fonts: McNeil DF Medium, Bold, Bold Italic


Drug Facts:	NA pt	Leading:	4.75 pt
Header:	4.75 pt	Bullets:	3.75 pt
Subheader:	4.75 pt	Barlines:	NA pt
Body Text:	4.75 pt	Hairlines:	NA pt
Drug Facts (continued):	NA pt		

Horizontal Scale: 75% Average Kerning: -10

McNeil
Consumer Healthcare

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) --- NO PRINT
- 5)  LOT AREA (NO ADDITIONAL COPY, NO COLOR)
- 6)  EXP. DATE AREA (NO ADDITIONAL COPY, NO COLOR)
- 7)  UPC AREA (NO COLOR)
- 8)  GLUE AREA (NO COPY, NO COLOR, NO VARNISH)

(b) (4)	7	8/6/98	SPLIT LOT & EXP AREA	DWG. NO.	FC-C-110	 CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA 19034 TITLE: CARTON, SEAL END 1.56 L X 1.56 W X 2.94 D	
	6	1/7/98	IMPORTED INTO MPSS	DRAWN			(b) (4)
	5	4-21-94	ADDED NO VARNISH AREA	PLT			1:1
	4	5-11-92	REDRAWN ON NEW FORMAT	SCALE			
	3	8-28-89	FLAP ANGLE AND RADIO CORRECTED	DATE			8-26-87
	2	10-11-88	DEPTH DIM ADDED TO JWS				
	1	3-21-88	NOTE ADDED: PRINT SIDE UP				
INIT.	NO	DATE	REVISIONS				

00000000

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Right before or after heart surgery.**

Ask a doctor before use if:

- you have problems or serious side effects from taking pain relievers or fever reducers
- you have stomach bleeding or problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

- taking aspirin or heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
- red, dark, or vomit blood in your stool or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- rashes or swelling is present in the painful area
- any new symptoms appear

Drug Facts

Active ingredient (in each tablet): ibuprofen 200 mg (NSAID) ... **Purpose:** Pain reliever/fever reducer

Uses: temporarily relieves minor aches and pains due to:

- headache
- minor pain of arthritis
- toothache
- the common cold
- menstrual cramps

Warnings:

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness/itching
- rash
- dizziness

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 65 or older
- have had stomach ulcers or bleeding problems
- take alcohol (drinking) or aspirin, blood thinners, or other drugs that increase the risk of stomach bleeding
- are on prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) or have a history of stomach ulcers, bleeding, or other problems

Every day while using this product, take more or every day while using this product, take more or

Drug Facts (continued)

Inactive ingredients: camellia wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide

Questions or comments?
Call 1-877-895-3665, weekdays 8:00 AM to 10:00 PM EST

Manufactured by:
McNeil Consumer Healthcare, LLC, LAS P EDRAS, PUERTO RICO 00771

McNeil Consumer Healthcare, DIVISION OF MCNEIL-PPC, INC.
FRT WASH NGTON, PA 19034 USA
MCN-PPC, NC 2010 www.motrin.com

00000000

Description: Ctn Motrin IB Tab 50 (LP Bulk)
File #: FDA3009897 Version: K-1
Date: 08/27/2010 Contact: (b) (4)

Labeling Format Information:

Fonts: Mcn DF Condensed, Bold, Bold Italic

Drug Facts:	8.25 pt	Leading:	6.50 pt
Header:	8.00 pt	Bullets:	5.00 pt
Subheader:	6.00 pt	Barlines:	2.50 pt
Body Text:	6.00 pt	Hairlines:	0.50 pt
Drug Facts (continued):	8.00 pt		

Horizontal Scale: 80% Average Kerning: 0


McNeil Submission Artwork
Consumer Healthcare

NOTES:


- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5) ▨ LDT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6) ▩ VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

INIT.	NO.	DATE	REVISIONS	DWG. NO.	DATE
(b) (4)	7	1/7/99	ADDED LDT/EXP AREA	L.B.L. 43	
	6	6/16/97	ADDED DV INFO.		
	3	10-24-94	CHANGED NO PRINT AREA TO 1/16		
	4	3-15-94	ADDED "EXP. DATE" TEXT		
	3	5-11-92	REBRAIN ON NEW FORMAT		
	2	11-21-89	LENGTH & WIDTH CHANGED		
	1	11-15-89	REVISION OF NO-PRINT AREA		

McNEIL	TITLE:
CONSUMER PRODUCTS CO.	HA LABEL
CAMP HILL ROAD, FORT WASHINGTON, PA 19034	4.25 L X 1.25 W



200% OF SIZE



ACTUAL SIZE

Description: <u>Lbl Motrin IB Tablet 50 - (b) (4)</u>		Labeling Format Information:	
File #: <u>3009897</u>	Version: <u>J-2</u>	Fonts: McNeil DF Medium, Bold, Bold Italic	
Date: <u>2/4/2010</u>	Contact: <u>(b) (4)</u>	Drug Facts: NA pt	Leading: 4.75 pt
		Header: 4.75 pt	Bullets: 3.75 pt
		Subheader: 4.75 pt	Barlines: NA pt
		Body Text: 4.75 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 75%	Average Kerning: -10

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/32 ANGLES ±1°
- 4) --- NO PRINT
- 5) [Hatched Box] LOT AREA (NO ADDITIONAL COPY, NO COLOR)
- 6) [Hatched Box] EXP. DATE AREA (NO ADDITIONAL COPY, NO COLOR)
- 7) [Hatched Box] UPC AREA (NO COLOR)
- 8) [Hatched Box] GLUE AREA (NO COPY, NO COLOR, NO VARNISH)

INIT.	NO.	DATE	REVISIONS	DWG. NO.	DATE
(b) (4)	7	8/6/98	SPLIT LOT & EXP AREA	FC-C-110	
	6	1/7/98	IMPORTED INTO MPSS		
	5	4-21-94	ADDED NO VARNISH AREA		
	4	5-11-92	REDRAWN ON NEW FORMAT		
	3	8-28-89	FLAP ANGLE AND RADIO CORRECTED		
	2	10-11-88	DEPTH DIM ADDED TO JWS		
	1	3-21-88	NOTE ADDED: PRINT SIDE UP		
				PLAT SCALE: 1:1	
				DATE: 8-26-87	

McNEIL
CONSUMER PRODUCTS CO.
CAMP HILL ROAD, FORT WASHINGTON, PA 19034
TITLE: CARTON, SEAL END
1.56 L X 1.56 W
X 2.94 D

00000000

00000000

Description: Ctn Motrin IB Tab 50+25 (LP Bulk)
File #: FDA3009897 Version: C-1
Date: 08/27/2010 Contact: (b) (4)

Labeling Format Information:

Fonts: Mcn DF Condensed, Bold, Bold Italic



Drug Facts:	8.25 pt	Leading:	6.50 pt
Header:	8.00 pt	Bullets:	5.00 pt
Subheader:	6.00 pt	Barlines:	2.50 pt
Body Text:	6.00 pt	Hairlines:	0.50 pt
Drug Facts (continued):	8.00 pt		

Horizontal Scale: 80%

Average Kerning: 0


McNeil Submission Artwork
Consumer Healthcare

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5)  LOT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6)  VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

INIT.	NO.	DATE	REVISIONS	DWG. NO.	DATE
(b) (4)	7	1/7/99	ADDED LOT/EXP AREA	L.B.L.43	
	6	6/16/97	ADDED DV INFO.		
	5	10-24-94	CHANGED NO PRINT AREA TO 1/16		
	4	3-15-94	ADDED "EXP. DATE" TEXT		
	3	5-11-92	REDRAWN ON NEW FORMAT		
	2	3-21-89	LENGTH & WIDTH CHANGED		
	1	3-15-89	REVISION OF NO-PRINT AREA		


McNEIL	TITLE:
CONSUMER PRODUCTS CO.	HA LABEL
CAMP HILL ROAD, FORT WASHINGTON, PA 19034	4.25 L X 1.25 W



200% OF SIZE




07844479








ACTUAL SIZE

Description: <u>Lbl Motrin IB Tablet 75</u>		Labeling Format Information:	
File #: <u>3009897</u>	Version: <u>D-1</u>	Fonts: McNeil DF Medium, Bold, Bold Italic	
Date: <u>12/2/09</u>	Contact: <u>(b) (4)</u>	Drug Facts: NA pt	Leading: 4.75 pt
		Header: 4.75 pt	Bullets: 4 pt
		Subheader: 4.75 pt	Barlines: NA pt
		Body Text: 4.75 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 75%	Average Kerning: -10



NOTES:


- 1)  NO COPY AREA. COLORS MAY BLEED IN THIS ZONE.
- 2)  DATA MATRIX AREA.
- 3)  DATA MATRIX QUIET ZONE. NO COPY, NO COLOR.
- 4)  LOT & EXP AREA. (TO BE NON-VARNISHED)
- 5) REFERENCE NATIONAL LABEL DRAWING #6-08-13C.

				DWG. NO. LB-PS-1193	 7050 Camp Hill Road, Fort Washington, PA 19034 TITLE: LABEL, SINGLE PLY 75cc BOTTLE
(b) (4)	4	11/3/08	UPDATED NOTES	DRAWN: (b) (4)	
	3	10/15/08	ADDED 1/2 DIM, NOTE AND LABEL P/N	SCALE: NONE	
	2	9/18/08	UPDATED LOT/EXP AREA	DATE: 6/30/08	
	1	6/30/08	INITIAL DRAFT	APPROVED:	
INIT.	NO.	DATE	REVISIONS	APPROVED:	








150% OF SIZE



Description: Lbl Motrin IB Tablet 100 (NCR)	(b) (4)	Labeling Format Information:	
File #: 3009897	Version: N-1	Fonts: McNeil DF Medium, Bold, Bold Italic	
Date: 12/18/09	Contact: (b) (4)	Drug Facts: NA pt	Leading: 5 pt
	(b) (4)	Header: 5 pt	Bullets: 4 pt
		Subheader: 5 pt	Barlines: NA pt
		Body Text: 5 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 84-85%	Average Kerning: -10
			

NOTES:

- 1)  NO COPY AREA. COLORS MAY BLEED IN THIS ZONE.
- 2)  DATA MATRIX AREA.
- 3)  DATA MATRIX QUIET ZONE. NO COPY, NO COLOR.
- 4)  LOT & EXP AREA. (TO BE NON-VARNISHED)
- 5) REFERENCE NATIONAL LABEL DRAWING #6-08-13C.


		DWG. NO.		LB-PS-1193			
		DRAWN: (b) (4)				Consumer Healthcare 7050 Camp Hill Road, Fort Washington, PA 19034	
(b) (4)	4	11/3/08	UPDATED NOTES	SCALE: NONE	TITLE: LABEL, SINGLE PLY		
	3	10/15/08	ADDED 1/2 DIM. NOTE AND LABEL P/N	DATE: 6/30/08	75cc BOTTLE		
	2	9/18/08	UPDATED LOT/EXP AREA	APPROVED:			
	1	6/30/08	INITIAL DRAFT				
INIT.	NO.	DATE	REVISIONS				







150% OF SIZE




ACTUAL SIZE

Description: Lbl Motrin IB Tablet 125 - LP BULK		Labeling Format Information:	
File #: 3009897	Version: T-1	Fonts: McNeil DF Medium, Bold, Bold Italic	
Date: 12/18/09	Contact: (b) (4)	Drug Facts: NA pt	Leading: 5 pt
		Header: 5 pt	Bullets: 4 pt
		Subheader: 5 pt	Barlines: NA pt
		Body Text: 5 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 84-85%	Average Kerning: -10
			

NOTES:


- 1)  NO COPY AREA. COLORS MAY BLEED IN THIS ZONE.
- 2)  DATA MATRIX AREA.
- 3)  DATA MATRIX QUIET ZONE. NO COPY, NO COLOR.
- 4)  LOT & EXP AREA. (TO BE NON-VARNISHED)
- 5) REFERENCE NATIONAL LABEL DRAWING #6-08-13C.

				DWG. NO.	 7050 Camp Hill Road, Fort Washington, PA 19034 TITLE: LABEL, SINGLE PLY 75cc BOTTLE
				LB-PS-1193	
(b) (4)	4	11/3/08	UPDATED NOTES	DRAWN: (b) (4)	
	3	10/15/08	ADDED 1/2 DIM, NOTE AND LABEL P/P/N	SCALE: NONE	
	2	9/18/08	UPDATED LOT/EXP AREA	DATE: 6/30/08	
	1	6/30/08	INITIAL DRAFT	APPROVED:	
INIT.	NO.	DATE	REVISIONS		







150% OF SIZE



Description: Lbi Motrin IB Tablet 150		Labeling Format Information:											
File #: 3009897	Version: B-1	Fonts: McNeil DF Medium, Bold, Bold Italic											
Date: 11/30/09	Contact: (b) (4)	<table border="1"> <tr> <td>Drug Facts: NA pt</td> <td>Leading: 5 pt</td> </tr> <tr> <td>Header: 5 pt</td> <td>Bullets: 4 pt</td> </tr> <tr> <td>Subheader: 5 pt</td> <td>Barlines: NA pt</td> </tr> <tr> <td>Body Text: 5 pt</td> <td>Hairlines: NA pt</td> </tr> <tr> <td>Drug Facts (continued): NA pt</td> <td></td> </tr> </table>		Drug Facts: NA pt	Leading: 5 pt	Header: 5 pt	Bullets: 4 pt	Subheader: 5 pt	Barlines: NA pt	Body Text: 5 pt	Hairlines: NA pt	Drug Facts (continued): NA pt	
Drug Facts: NA pt	Leading: 5 pt												
Header: 5 pt	Bullets: 4 pt												
Subheader: 5 pt	Barlines: NA pt												
Body Text: 5 pt	Hairlines: NA pt												
Drug Facts (continued): NA pt													
		Horizontal Scale: 84-85%	Average Kerning: -10										
													

NOTES:

1. ALL DIMENSIONS ARE IN INCHES
2. BASED ON OSG IVERS-LEE DRAWING #: 87743.
3.  SEAL AREA: NO PRINT
4.  SLIT
5.  LOT/EXP CODING AREA: NO VARNISH, NO PRINT.
6.  EYEMARK
7. "FRONT" USE STANDARD UNWIND #3.
8. "BACK" USE STANDARD UNWIND #4.
9. * .0156 OF SCRAP ON BOTH SIDES.


				DWG. NO. UNWIND-192	
				DRAWN : (b) (4)	
				PLOT : NONE	
				SCALE :	
				DATE : 6/20/06	
				APPROVED :	
(b) (4)	3	11/30/06	UPDATED PER AS BUILT		
	2	9/27/06	ADJUST WEB DIMENSIONS		
	1	8/12/06	UPDATE TEAR NOTCH DIMENSIONS		
INIT.	NO.	DATE	REVISIONS		

McNeil
Consumer Healthcare
Camp Hill Road, Fort Washington, PA 19034

TITLE
**POUCH MATERIAL
WEB LAYOUT**
2.031 L x 2.640 W



150% OF SIZE





Description: Pouch Motrin IB Caplet CR-LP Bulk		Labeling Format Information:	
File #: 3010048	Version: B-1	Fonts: McNeil DF Cond. Bold and Italic, Helv. Black and Italic	
Date: 03/21/10	Contact: (b) (4)	Drug Facts: NA pt	Leading: 4.5 pt
		Header: 4.5 pt	Bullets: 3.5 pt
		Subheader: 4.5 pt	Barlines: NA pt
		Body Text: 4.5 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 73%	Average Kerning: 0

(b) (4)



NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5)  LOT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6)  VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

(b) (4)	7	1/7/99	ADDED LOT/EXP AREA	DWG. NO.	McNEIL CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA 19034 TITLE: HA LABEL 4.25 L X 1.25 W	
	6	6/16/97	ADDED DV INFO.	L.B.L. 43		
	5	10-24-94	CHANGED NO PRINT AREA TO 1/16	DRAWN		(b) (4)
	4	3-15-94	ADDED "EXP. DATE" TEXT	PLOT SCALE		NONE
	3	5-11-92	REDRAWN ON NEW FORMAT	DATE		11-15-89
	2	3-21-89	LENGTH & WIDTH CHANGED			
	1	3-15-89	REVISION OF NO-PRINT AREA			
INIT.	NO.	DATE	REVISIONS			

Do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing.

NDC 50580-110-03

Motrin[®] IB

Ibuprofen Tablets usp. 200 mg
 Pain Reliever/Fever Reducer (NSAID)
 24 Coated Caplets * Capsule-Shaped Tablets

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet) Purpose
 Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer
 nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**
Directions ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F)

Manufactured by: McNeil Healthcare LLC, US PEDS/AS, PUERTO RICO 00711
 For: McNeil-PPC, Inc., FORT WASHINGTON, PA 19034 USA
 ©McNeil-PPC, Inc. 2010 www.motrin.com 7844436
 EXP. DATE: _____ LOT: _____

200% OF SIZE



Do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing.

NDC 50580-110-03

Motrin[®] IB

Ibuprofen Tablets usp. 200 mg
 Pain Reliever/Fever Reducer (NSAID)
 24 Coated Caplets * Capsule-Shaped Tablets

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet) Purpose
 Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer
 nonsteroidal anti-inflammatory drug




Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis


■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**
Directions ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F)

Manufactured by: McNeil Healthcare LLC, US PEDS/AS, PUERTO RICO 00711
 For: McNeil-PPC, Inc., FORT WASHINGTON, PA 19034 USA
 ©McNeil-PPC, Inc. 2010 www.motrin.com 7844436
 EXP. DATE: _____ LOT: _____

Description: Lbl Motrin IB Caplet 24-LP Bulk File #: 3009923 Version: P-1 Date: 1/12/2010 Contact: (b) (4)	Labeling Format Information: Fonts: Helvetica Condensed, Bold, Bold Italic <table border="1" style="width: 100%; border-collapse: collapse; font-size: x-small;"> <tr> <td>Drug Facts:</td> <td>NA pt</td> <td>Leading:</td> <td>4.75 pt</td> </tr> <tr> <td>Header:</td> <td>4.75 pt</td> <td>Bullets:</td> <td>3.75 pt</td> </tr> <tr> <td>Subheader:</td> <td>4.75 pt</td> <td>Barlines:</td> <td>NA pt</td> </tr> <tr> <td>Body Text:</td> <td>4.75 pt</td> <td>Hairlines:</td> <td>NA pt</td> </tr> <tr> <td>Drug Facts (continued):</td> <td>NA pt</td> <td></td> <td></td> </tr> </table> Horizontal Scale: 74-75% Average Kerning:-10	Drug Facts:	NA pt	Leading:	4.75 pt	Header:	4.75 pt	Bullets:	3.75 pt	Subheader:	4.75 pt	Barlines:	NA pt	Body Text:	4.75 pt	Hairlines:	NA pt	Drug Facts (continued):	NA pt		
Drug Facts:	NA pt	Leading:	4.75 pt																		
Header:	4.75 pt	Bullets:	3.75 pt																		
Subheader:	4.75 pt	Barlines:	NA pt																		
Body Text:	4.75 pt	Hairlines:	NA pt																		
Drug Facts (continued):	NA pt																				

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) --- NO PRINT
- 5)  LOT & EXP. DATE AREA (NO ADDITIONAL COPY, NO COLOR)
- 6)  UPC AREA (NO COLOR)
- 7)  GLUE AREA (NO COPY, NO COLOR, NO VARNISH)

(b) (4)	7	6/16/97	ADDED IV INFO	IVG. NO.	FC-C-111	 CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA. 19034 TITLE: CARTON, SEAL END 1.50 L X 1.44 W X 2.63 D	
	6	3-15-94	ADDED 'EXP.' TEXT	DRAWN			(b) (4)
	5	10-18-93	REVISED NO VARNISH AREA	PLT. SCALE			NONE
	4	9-1-93	ADDED UPC AREA	DATE			8-31-87
	3	7-1-92	REDRAWN ON NEW FORMAT				
	2	6-6-88	IVG. FLIPPED AS LH CARTON				
	1	3-21-88	NOTE ADDED PRINT SIDE UP				
DET.	NO.	DATE	REVISIONS				

0000000

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to another pain reliever or fever reducer. An allergic reaction may include hives, rash, or difficulty breathing. Stop use and ask a doctor if you experience any of the following symptoms:

- skin rash or hives
- difficulty breathing
- wheezing
- swelling of the face, lips, tongue, or throat
- dizziness
- fainting
- severe dizziness
- severe lightheadedness
- severe weakness or fatigue
- severe stomach pain
- severe constipation
- severe diarrhea
- severe nausea or vomiting
- severe headache
- severe drowsiness
- severe confusion
- severe irritability
- severe changes in behavior
- severe changes in vision
- severe changes in hearing
- severe changes in taste or smell
- severe changes in touch or feeling
- severe changes in temperature
- severe changes in sweating
- severe changes in heart rate
- severe changes in blood pressure
- severe changes in breathing
- severe changes in circulation
- severe changes in digestion
- severe changes in excretion
- severe changes in reproduction
- severe changes in development
- severe changes in aging
- severe changes in death

Drug Facts

Active ingredient (in each tablet): Ibuprofen 200 mg (NSAID). **Purpose:** Pain reliever/fever reducer.

Use: Use temporarily to relieve minor aches and pains, such as headache, muscle aches, minor pain of arthritis, rheumatoid arthritis, osteoarthritis, menstrual cramps, and temporary relief of fever.

Warnings: Ibuprofen may cause serious allergic reactions, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- shock
- skin redness, rash
- dizziness
- fainting
- severe dizziness
- severe lightheadedness
- severe weakness or fatigue
- severe stomach pain
- severe constipation
- severe diarrhea
- severe nausea or vomiting
- severe headache
- severe drowsiness
- severe confusion
- severe irritability
- severe changes in behavior
- severe changes in vision
- severe changes in hearing
- severe changes in taste or smell
- severe changes in touch or feeling
- severe changes in temperature
- severe changes in sweating
- severe changes in heart rate
- severe changes in blood pressure
- severe changes in breathing
- severe changes in circulation
- severe changes in digestion
- severe changes in excretion
- severe changes in reproduction
- severe changes in development
- severe changes in aging
- severe changes in death

Drug Facts (continued)

Inactive ingredients: carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide.

Questions or comments?
Call 1-877-895-3665; weekdays 8:00 AM to 8:00 PM EST.

Distributed by:
McNeil Consumer Healthcare
DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
©MCN-PPC, INC. 2010 www.motrin.com

0000000

Description: Ctn Motrin IB Caplet 24+6
File #: FDA3010135 Version: 1-1
Date: 9/08/2010 Contact: (b) (4)

Labeling Format Information:



Fonts: McN DF Condensed, Bold, Bold Italic


Drug Facts:	8.25 pt	Leading:	6.00 pt
Header:	7.00 pt	Bullets:	5.00 pt
Subheader:	6.00 pt	Barlines:	2.50 pt
Body Text:	6.00 pt	Hairlines:	0.50 pt
Drug Facts (continued):	7.00 pt		


Horizontal Scale: 70% Average Kerning: 0



NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5)  LDT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6)  VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

(b) (4)	7	1/7/99	ADDED LOT/EXP AREA	DWG. NO.	 CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA 19034 TITLE: HA LABEL 4.25 L X 1.25 W	
	6	6/16/97	ADDED DV INFO	L.B.L. 43		
	5	10-24-94	CHANGED NO PRINT AREA TO 1/16	DRAWN		(b) (4)
	4	3-15-94	ADDED "EXP. DATE" TEXT	PLOT SCALE		NONE
	3	5-11-92	REDRAWN ON NEW FORMAT	DATE		11-15-89
	2	3-21-89	LENGTH & WIDTH CHANGED			
INIT.	NO.	DATE	REVISIONS			



Do not use if neck wrap or foil inner seal imprinted "Safety Seal" is broken or missing.

Motrin[®] IB
 ibuprofen Tablets USP, 200 mg
 Pain Reliever/Fever Reducer (NSAID)
 24+6 Coated Caplets* *Capsule-Shaped Tablets

Important: Read all product information before using. Keep the box for important information.


Active ingredient (in each caplet)
 Ibuprofen 200 mg (NSAID)*Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Purpose
 *nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children. Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F)

200% of size


Description: Lbl Motrin IB Caplet 24+6 - LP BULK		Labeling Format Information:	
File #: 3009923	Version: V-1	Fonts: Helvetica Condensed, Bold, Bold Italic	
Date: 1/6/2010	Contact: (b) (4)	Drug Facts: NA pt	Leading: 4.75 pt
		Header: 4.75 pt	Bullets: 3.75 pt
		Subheader: 4.75 pt	Barlines: NA pt
		Body Text: 4.75 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 74-75%	Average Kerning:-10

McNeil
Consumer Healthcare


NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5) [X] LDT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6) [X] VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

(b) (4)	7	1/7/99	ADDED LDT/EXP AREA	DWG. NO.	McNEIL CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA 19034 TITLE: HA LABEL 4.25 L X 1.25 W	
	6	6/16/97	ADDED DV INFO.	L.B.L. 43		
	5	10-24-94	CHANGED NO PRINT AREA TO 1/16	DRAWN		(b) (4)
	4	3-15-94	ADDED "EXP. DATE" TEXT	PLOT SCALE		NONE
	3	5-11-92	REDRAWN ON NEW FORMAT	DATE		11-15-89
	2	3-21-89	LENGTH & WIDTH CHANGED			
	1	3-15-89	REVISION OF NO-PRINT AREA			
INIT.	NO.	DATE	REVISIONS			



200% OF SIZE



Description: Lbl Motrin IB Caplet 50 -LP Bulk		Labeling Format Information:	
File #:	3009923	Version:	L-1
Date:	1/12/2010	Contact:	(b) (4)
		Fonts: Helvetica Condensed, Bold, Bold Italic Drug Facts: NA pt Leading: 4.75 pt Header: 4.75 pt Bullets: 3.75 pt Subheader: 4.75 pt Barlines: NA pt Body Text: 4.75 pt Hairlines: NA pt Drug Facts (continued): NA pt Horizontal Scale: 74-75% Average Kerning:-10	

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) --- NO PRINT
- 5) [Hatched Box] LOT AREA (NO ADDITIONAL COPY, NO COLOR)
- 6) [Hatched Box] EXP. DATE AREA (NO ADDITIONAL COPY, NO COLOR)
- 7) [Hatched Box] UPC AREA (NO COLOR)
- 8) [Hatched Box] GLUE AREA (NO COPY, NO COLOR, NO VARNISH)

INT.	NO.	DATE	REVISIONS	DWG. NO.	DATE
(b) (4)	7	8/6/98	SPLIT LOT & EXP AREA	FC-C-110	
	6	1/7/98	IMPORTED INTO MPSS		
	5	4-21-94	ADDED NO VARNISH AREA		
	4	5-11-92	REDRAWN ON NEW FORMAT		
	3	8-29-89	FLAP ANGLE AND RADII CORRECTED		
	2	10-11-88	DEPTH DIM ADDED TO DWG.		
	1	3-21-88	NOTE ADDED: PRINT SIDE UP		
				DRAWN (b) (4)	
				PLOT SCALE 1:1	
					DATE 8-26-87

McNEIL
CONSUMER PRODUCTS CO.
CAMP HILL ROAD, FORT WASHINGTON, PA. 19034
TITLE:
CARTON, SEAL END
1.56 L X 1.56 W
X 2.94 D

0000000

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Aspirin** before or after heart surgery.

Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers. The stomach bleeding warning applies to you if you have a history of stomach problems, such as heartburn, you take high blood pressure, heart disease, liver disease, or kidney disease, you have asthma, or you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin.

Under a doctor's care for any serious condition. **Do not take any other drug.**

When using this product:

- Take with food or milk. (Stomach upset occurs.)
- Do not take more than directed.
- Do not take more than directed or for longer than directed.

Stop use and ask a doctor if:

- You experience any of the following signs of stomach bleeding:
- Heartburn, vomit blood, or have bloody or black stools.
- Have stomach pain that does not get better.
- Pain gets worse or lasts more than 10 days.
- Other gets worse or lasts more than 3 days.
- Redness or swelling is present in the painful area.
- Any new symptoms appear.

Drug Facts

Active ingredient (in each caplet): Ibuprofen 200 mg (NSAID). **Pain reliever or fever reducer.** "Nonsteroidal anti-inflammatory drug."

Uses: temporarily relieves minor aches and pains due to:

- Headache
- Muscle aches
- Minor pain of arthritis
- Backache
- Menstrual cramps
- Temporarily reduces fever.

Warnings:

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- Hives
- Swelling of the face, lips, tongue, or throat (may be life threatening)
- Difficulty breathing
- Stomach pain or cramping
- Itchy skin
- Blurred vision or dizziness

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance of bleeding is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, are taking blood thinning (anticoagulant) or aspirin, or taking other drugs containing aspirin or NSAIDs. Stomach bleeding can occur without warning and may be serious. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- Heartburn, vomit blood, or have bloody or black stools.
- Have stomach pain that does not get better.
- Pain gets worse or lasts more than 10 days.
- Other gets worse or lasts more than 3 days.
- Redness or swelling is present in the painful area.
- Any new symptoms appear.

Drug Facts (continued)

Inactive ingredients: ceresin wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide.

Questions or comments?
Call 1-877-895-3665, weekdays 8:00 AM to 8:00 PM EST.

Distributed by:
McNeil Consumer Healthcare
DIVISION OF MCNEIL PPC, INC.
FORT WASHINGTON, PA 19034 USA
©MNC PPC, INC. 2010 www.motrin.com

0000000

0000000

Description: CTN Motrin IB Cpt 60 H/G
File #: FDA3010135 Version: K-1
Date: 9/08/2010 Contact: (b) (4)

Labeling Format Information:

Fonts: McN DF Condensed, Bold, Bold Italic



Drug Facts:	8.25 pt	Leading:	6.50 pt
Header:	7.00 pt	Bullets:	5.00 pt
Subheader:	6.00 pt	Barlines:	2.50 pt
Body Text:	6.00 pt	Hairlines:	0.50 pt
Drug Facts (continued):	7.00 pt		

Horizontal Scale: 75%

Average Kerning: 0

McNeil Submission Artwork
Consumer Healthcare

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5)  LOT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6)  VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

(b) (4)	7	1/7/99	ADDED LOT/EXP AREA	DWG. NO.	
	6	6/16/97	ADDED DV INFO	L.B.L. 43	
	3	10-24-94	CHANGED NO PRINT AREA TO 1/16		
	4	3-15-94	ADDED 'EXP. DATE' TEXT	DRAWN	(b) (4)
	2	5-11-92	REDRAWN ON NEW FORMAT	PLOT SCALE	NONE
	2	3-21-89	LENGTH & WIDTH CHANGED	DATE	11-15-89
	1	3-15-89	REVISION OF NO-PRINT AREA		
INIT.	NO.	DATE	REVISIONS		

McNEIL
CONSUMER PRODUCTS CO.
CAMP HILL ROAD, FORT WASHINGTON, PA 19034
TITLE:
HA LABEL
4.25 L X 1.25 W

For Hospital and Government Use Only
NDC 50580-110-60

Motrin[®] IB
Ibuprofen Tablets
USP, 200 mg
Pain Reliever
Fever Reducer (NSAID)

60 Coated Caplets* *Capsule Shaped Tablets

Keep the box for important information.

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** **Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25° C (68-77° F)

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
 For McNeil, P.C., INC., FORT WASHINGTON, PA 19034, USA
 800-444-495
 www.motrin.com
 7844495

200% OF SIZE



For Hospital and Government Use Only
NDC 50580-110-60

Motrin[®] IB
Ibuprofen Tablets
USP, 200 mg
Pain Reliever
Fever Reducer (NSAID)

60 Coated Caplets* *Capsule Shaped Tablets

Keep the box for important information.

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** **Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25° C (68-77° F)

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
 For McNeil, P.C., INC., FORT WASHINGTON, PA 19034, USA
 800-444-495
 www.motrin.com
 7844495

Description: Lbl Motrin IB Caplet 60 H/G - (b) (4)

File #: 3009923 Version: B-1

Date: 12/23/2009 Contact: (b) (4)

Labeling Format Information:

Fonts: Helvetica Condensed, Bold, Bold Italic

Drug Facts:	NA pt	Leading:	4.75 pt
Header:	4.75 pt	Bullets:	3.75 pt
Subheader:	4.75 pt	Barlines:	NA pt
Body Text:	4.75 pt	Hairlines:	NA pt
Drug Facts (continued):	NA pt		

Horizontal Scale: 74-75% Average Kerning:-10

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) --- NO PRINT
- 5) [Hatched Box] LOT AREA (NO ADDITIONAL COPY, NO COLOR)
- 6) [Hatched Box] EXP. DATE AREA (NO ADDITIONAL COPY, NO COLOR)
- 7) [Hatched Box] UPC AREA (NO COLOR)
- 8) [Hatched Box] GLUE AREA (NO COPY, NO COLOR, NO VARNISH)

(b) (4)	7	8/6/98	SPLIT LOT & EXP AREA	DWG. NO.	FC-C-110	McNEIL CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA 19024
	6	1/7/98	IMPORTED INTO MPSS	DRAWN	(b) (4)	
	5	4-21-94	ADDED NO VARNISH AREA	PLOT SCALE	1:1	TITLE: CARTON, SEAL END
	4	5-11-92	REBORN ON NEW FORMAT	DATE	8-26-87	1.56 L X 1.56 W X 2.94 D
	3	8-28-89	FLAP ANGLE AND RADIUS CORRECTED			
	2	12-11-88	DEPTH DIM ADDED TO DWG			
	1	3-21-88	NOTE ADDED: PRINT SIDE UP			
INIT.	MD	DATE	REVISIONS			

08737381

See New Warnings Information NDC 50390-110-76

Motrin[®] IB
Ibuprofen Tablets USP 200 mg
Pain Reliever/Fever Reducer (NSAID)
50+25[®] Coated Caplets[®] *Capsule-Shaped Tablets

BONUS! 50% MORE

Drug Facts (continued)
Inactive ingredients: carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hydroxypropyl methylcellulose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shear, stearic acid, titanium dioxide

Questions or comments?
Call 1-877-895-3665; weekdays 8:00AM to 8:00PM EST

Manufactured by: McNeil Healthcare LLC
LAS PIEDRAS, PUERTO RICO 00771
For: McNeil Consumer Healthcare
DIVISION OF McNEIL PPC, INC.
FORT WASHINGTON, PA 19054 USA
©MCMC PPC, INC. 2010 www.motrin.com

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless it is clearly needed to do so by a doctor because it may cause problems in the unborn child or complications during the birth of children.
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222).

Directions
Do not take more than directed.
The smallest effective dose should be used.
Adults and children 12 years and older: Take 1 caplet every 4 to 6 hours while in pain. Do not take more than 4 caplets in 24 hours.
Children under 12 years: Do not use without a doctor's advice.
Do not use if you are allergic to any of the ingredients listed above or if you have ever had an allergic reaction to ibuprofen.
Ask a doctor if you are taking any other medicines, especially aspirin, because they may increase the risk of bleeding.
Ask a doctor if you are taking any other medicines, especially aspirin, because they may increase the risk of bleeding.

Other Information
Store between 20° to 25°C (68° to 77°F).
Do not use if neck wrap or foil liner seal is imprinted with "Safety Seal" or is broken or missing.
Seventeen panels for 17 numbers are expansion date.

08737381

Description:	Ctn Motrin IB Caplet 50+25	(b) (4)
File #:	3009923	Version: G-1
Date:	01/04/2010	Contact: (b) (4)

(b) (4)

Labeling Format Information:



Fonts: McNeil DF Condensed, Bold, Bold Italic

Drug Facts:	8.25 pt	Leading:	6.25 pt
Header:	7 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	2.5 pt
Body Text:	6 pt	Hairlines:	0.5 pt
Drug Facts (continued):	7 pt		

Horizontal Scale: 70-80% Average Kerning: 0


McNeil
Consumer Healthcare

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES.
- 3) TOLERANCES UNLESS SPECIFIED FRACTIONS ±1/64
- 4) ----- NO PRINT AREA
- 5)  LDT & EXP. DATE AREA (NO COLOR, NO COPY)
- 6)  VERIFICATION BARCODE AREA (NO COLOR, NO COPY)

(b) (4)	7	1/7/99	ADDED LOT/EXP AREA	DUG NO.	L.B.L. 4.3	McNEIL
	8	10-24-04	CHANGED NO PRINT AREA TO 1/16	CONSUMER PRODUCTS CO.		
	9	10-24-04	ADDED "EXP. DATE" TEXT	CHP, HLL, KDA, FORT WASHINGTON, PA 19044		
	10	12-12-04	REWORKED ON NEW FORMAT			
	11	12-12-04	LENGTH & WIDTH CHANGED	DATE	NONE	TITLE
	12	12-12-04	REVISION OF NO-PRINT AREA	DATE	11-15-89	HA LABEL
						4.25 L X 1.25 W

Do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing.



Motrin^{IB}
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
75 Coated Caplets* *Capsule-Shaped Tablets

Important: Read all product information before using. Keep the box for important information.

Active Ingredient (in each caplet) **Purpose**
 ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to: headache
 muscular aches minor pain of arthritis toothache backache the

EXP. DATE

LOT

7844442

common cold menstrual cramps temporarily reduces fever. **Warnings: Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery. **Ask a doctor before use if** you have problems or serious side effects from taking pain relievers or fever reducers the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you have asthma you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** taking aspirin to heart attack or stroke, because ibuprofen may decrease this benefit of aspirin under a doctor's care for any serious condition taking any other drug. **When using this product** take with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. **Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions** do not take more than directed at the smallest effective dose should be used. **Adults and children 12 years and older:** take 1 caplet every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 caplet, 2 caplets may be used do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ask a doctor. **Other information** store between 20-25°C (68-77°F)

200% OF SIZE



DO NOT USE IF NECK WRAP OR FOLIO INNER SEAL imprinted "Safety Seal®" is broken or missing.



Motrin^{IB}
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
75 Coated Caplets* *Capsule-Shaped Tablets

Important: Read all product information before using. Keep the box for important information.

Active Ingredient (in each caplet) **Purpose**
 ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug


Uses temporarily relieves minor aches and pains due to: headache
 muscular aches minor pain of arthritis toothache backache the

EXP. DATE

LOT

7844442

common cold menstrual cramps temporarily reduces fever. **Warnings: Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery. **Ask a doctor before use if** you have problems or serious side effects from taking pain relievers or fever reducers the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you have asthma you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** taking aspirin to heart attack or stroke, because ibuprofen may decrease this benefit of aspirin under a doctor's care for any serious condition taking any other drug. **When using this product** take with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. **Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions** do not take more than directed at the smallest effective dose should be used. **Adults and children 12 years and older:** take 1 caplet every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 caplet, 2 caplets may be used do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ask a doctor. **Other information** store between 20-25°C (68-77°F)

<p>Description: Lbl Motrin IB CplT 75 - (b) (4)</p> <p>File #: 3009923 Version: H-1</p> <p>Date: 1/6/2010 Contact: (b) (4)</p>	<p>Labeling Format Information:</p> <p>Fonts: McNeil Cond Medium, Bold, Bold Italic</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td>Drug Facts:</td> <td>NA pt</td> <td>Leading:</td> <td>4.25 pt</td> </tr> <tr> <td>Header:</td> <td>4.5 pt</td> <td>Bullets:</td> <td>3.5 pt</td> </tr> <tr> <td>Subheader:</td> <td>4.5 pt</td> <td>Barlines:</td> <td>NA pt</td> </tr> <tr> <td>Body Text:</td> <td>4.5 pt</td> <td>Hairlines:</td> <td>NA pt</td> </tr> <tr> <td>Drug Facts (continued):</td> <td>NA pt</td> <td></td> <td></td> </tr> </table> <p>Horizontal Scale: 65% Average Kerning: -35</p> <div style="text-align: center;">  </div>	Drug Facts:	NA pt	Leading:	4.25 pt	Header:	4.5 pt	Bullets:	3.5 pt	Subheader:	4.5 pt	Barlines:	NA pt	Body Text:	4.5 pt	Hairlines:	NA pt	Drug Facts (continued):	NA pt		
Drug Facts:	NA pt	Leading:	4.25 pt																		
Header:	4.5 pt	Bullets:	3.5 pt																		
Subheader:	4.5 pt	Barlines:	NA pt																		
Body Text:	4.5 pt	Hairlines:	NA pt																		
Drug Facts (continued):	NA pt																				

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) [Hatched Box] LDT & EXP. DATE AREA
- 5) [Cross-hatched Box] UPC AREA
- 6) [Diagonal Lines] NO VARNISH AREA
- 7) - - - - NO PRINT

INIT.	NL.	DATE	REVISIONS	DWG. NO.	DATE
(b) (4)	5	4/30/08	EXPORTED INTO NPES	FC-C-183	
	4	4/3/95	ADDED NO VARNISH AREA		
	3	2-2-95	UPDATED TO CURRENT STANDARDS		
	2	12-31-93	REBROWN ON NEW FORMAT		
	1	3-21-90	NOTE ADDED PRINT SIDE UP		

McNEIL	TITLE
CONSUMER PRODUCTS CO.	CARTON, SEAL END
CAMP HILL ROAD, FORT WASHINGTON, PA 19034	1.88 L X 1.88 W
	X 3.50 D

08737372



Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn, indigestion, ulcers, or bleeding disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
- you are serious pain reliever does not get better
- you use serious pain reliever more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient (in each caplet): Purpose
Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

USES

- temporarily relieves minor aches and pains, due to:
 - headache
 - muscle aches
 - minor pain of arthritis
 - toothache
 - the common cold
 - menstrual cramps
 - temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- skin redness/itching
- asthma (wheezing)
- shock
- skin swelling
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take aspirin, blood thinners or bleeding problems
- take alcohol during (or soon after) use of this drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments?
Call 1-877-895-3665; weekdays 8:00 AM to 8:00 PM EST

McNeil

Manufactured by:
McNeil Hea thcare LLC
LAS PEDRAS, PUERTO RICO 00771
For McNeil Consumer Hea thcare
DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
©MCN-PPC, INC. 2010
www.motrin.com



Description: Ctn Motrin IB Caplet 100- LP Bulk File #: 3009923 Version: M-1 Date: 1/8/2010 Contact: (b) (4)	Labeling Format Information: Fonts: McNeil Condensed Med, Bold, Bold Italic <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 2px;">Drug Facts: 8.25 pt</td> <td style="padding: 2px;">Leading: 6.25 pt</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">Header: 7 pt</td> <td style="padding: 2px;">Bullets: 5 pt</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">Subheader: 6 pt</td> <td style="padding: 2px;">Barlines: 2.5 pt</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">Body Text: 6 pt</td> <td style="padding: 2px;">Hairlines: .5 pt</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">Drug Facts (continued): 7 pt</td> <td style="padding: 2px;"></td> </tr> </table> <p style="font-size: small; margin-top: 5px;">Horizontal Scale: 78-80% Average Kerning: 0</p>	Drug Facts: 8.25 pt	Leading: 6.25 pt	Header: 7 pt	Bullets: 5 pt	Subheader: 6 pt	Barlines: 2.5 pt	Body Text: 6 pt	Hairlines: .5 pt	Drug Facts (continued): 7 pt	
Drug Facts: 8.25 pt	Leading: 6.25 pt										
Header: 7 pt	Bullets: 5 pt										
Subheader: 6 pt	Barlines: 2.5 pt										
Body Text: 6 pt	Hairlines: .5 pt										
Drug Facts (continued): 7 pt											

NOTES:

- 1) [Pattern] NO COPY AREA. COLORS MAY BLEED IN THIS ZONE.
- 2) [Pattern] DATA MATRIX AREA.
- 3) [Pattern] DATA MATRIX QUIET ZONE. NO COPY, NO COLOR.
- 4) [Pattern] LOT & EXP AREA. (TO BE NON-VARNISHED)
- 5) REFERENCE NATIONAL LABEL DRAWING #6-08-13C.

		DVG. NO.		LB-PS-1193		McNeil Consumer Healthcare	
		DRAWN		(b) (4)		1909 Camp Hill Road, Fort Washington, PA 19004	
		SCALE:		NONE		TITLE	
		DATE:		6/30/08		LABEL, SINGLE PLY	
		APPROVED				75cc BOTTLE	
DIT.	NO.	DATE	REVISIONS				
(b) (4)	4	11/2/08	UPDATED NOTES				
	3	10/15/08	ADDED 1/2 INK NOTE AND LABEL P/N				
	2	9/18/08	UPDATED LOT/EXP AREA				
	1	6/30/08	INITIAL DRAFT				

784434

LOT: EXP:

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever

NDC 50580-110-01

Motrin IB
Ibuprofen Tablets usp. 200 mg
Pain Reliever /Fever Reducer (NSAID)
100 Coated Caplets* *Capsule-Shaped Tablets

reducer right before or after heart surgery. Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers or the stomach bleeding warning applies to you. you have a history of stomach problems, such as heartburn or you have high blood pressure, heart disease, liver cirrhosis, or kidney disease. you have asthma. you are taking a diuretic. Ask a doctor or pharmacist before use if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin. under a doctor's care for any serious condition. taking any other drug. When using this product, take with food or milk if stomach upset occurs. the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Directions: do not take more than directed. the smallest effective dose should be used. Adults and children 12 years and older: take 1 caplet every 4 to 6 hours while symptoms persist. if pain or fever does not respond to 1 caplet, 2 caplets may be used. do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ask a doctor. Other information: store between 20-25°C (68-77°F). do not use if neck wrap or foil inner seal imprinted "Safety Seal" is broken or missing.

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
For: MCNEIL PPC, INC., FORT WASHINGTON, PA 19034 USA
©MCN-PPC, INC. 2010 www.motrin.com

150% OF SIZE

784434

LOT: EXP:

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever


NDC 50580-110-01

Motrin IB
Ibuprofen Tablets usp. 200 mg
Pain Reliever /Fever Reducer (NSAID)
100 Coated Caplets* *Capsule-Shaped Tablets




reducer right before or after heart surgery. Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers or the stomach bleeding warning applies to you. you have a history of stomach problems, such as heartburn or you have high blood pressure, heart disease, liver cirrhosis, or kidney disease. you have asthma. you are taking a diuretic. Ask a doctor or pharmacist before use if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin. under a doctor's care for any serious condition. taking any other drug. When using this product, take with food or milk if stomach upset occurs. the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Directions: do not take more than directed. the smallest effective dose should be used. Adults and children 12 years and older: take 1 caplet every 4 to 6 hours while symptoms persist. if pain or fever does not respond to 1 caplet, 2 caplets may be used. do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ask a doctor. Other information: store between 20-25°C (68-77°F). do not use if neck wrap or foil inner seal imprinted "Safety Seal" is broken or missing.

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
For: MCNEIL PPC, INC., FORT WASHINGTON, PA 19034 USA
©MCN-PPC, INC. 2010 www.motrin.com

Description: LB1 Motrin IB Caplet 100 - LP Bulk		Labeling Format Information:	
File #: 3009923	Version: N-1	Fonts: McNeil Condensed Med, Bold, Bold Italic	
Date: 1/8/2010	Contact: (b) (4)	Drug Facts: NA pt	Leading: 5 pt
		Header: 5 pt	Bullets: 4 pt
		Subheader: 5 pt	Barlines: NA pt
		Body Text: 5 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 82-83%	Average Kerning: 0
			

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4)  LOT & EXP. DATE AREA
- 5)  UPC AREA
- 6)  NO VARNISH AREA
- 7) - - - - NO PRINT

INIT.	NO.	DATE	REVISIONS	DWG. NO.	DATE
(b) (4)	5	4/30/08	EXPORTED INTO NPES	FC-C-183	
	4	4/3/96	ADDED NO VARNISH AREA		
	3	2-2-95	UPDATED TO CURRENT STANDARDS		
	2	12-31-93	REBORN ON NEW FORMAT		
	1	3-21-90	NOTE ADDED PRINT SIDE UP		

McNEIL	
CONSUMER PRODUCTS CO.	
CAMP HILL ROAD, FORT WASHINGTON, PA 19034	
TITLE: CARTON, SEAL END	
1.88 L X 1.88 W X 3.50 D	

08737382



Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems such as heartburn, indigestion, ulcers, or bleeding
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
- the stomach pain does not get better
- you see black or tarry stools in your stool
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient (in each caplet): Purpose
Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains, due to:
 - headache
 - muscle aches
 - minor pain of arthritis
 - toothache
 - the common cold
 - menstrual cramps
 - temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- skin swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take other drugs containing aspirin or bleeding problems
- take alcohol during (or soon after) use of this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments?
Call 1-877-895-3665; weekdays 8:00 AM to 8:00 PM EST

McNeil
Manufactured by:
McNeil Hea thcare LLC
LAS PIEDRAS, PUERTO RICO 00771
For McNeil Consumer Hea thcare
DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
©MCN-PPC, INC. 2010
www.motrin.com



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Warnings

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 caplet every 6 to 8 hours while pain or fever lasts
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

ask a doctor if you are under 12 years

Other information

- store between 20°-25°C (68°-77°F)
- do not use if neck wrap or foil inner seal is printed
- "Safety Seal" is broken or missing
- see end panel for lot number and expiration date


Inactive ingredients: cannauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose,

EXP: 3 0045-0481-955





08737382

Description: Ctn Motrin IB Caplet 100+25- (b) (4) File #: 3009923 Version: C-2 Date: 2/8/2010 Contact: (b) (4)	Labeling Format Information: Fonts: McNeil Condensed Med, Bold, Bold Italic <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td>Drug Facts:</td> <td style="text-align: right;">8.25 pt</td> <td>Leading:</td> <td style="text-align: right;">6.25 pt</td> </tr> <tr> <td>Header:</td> <td style="text-align: right;">7 pt</td> <td>Bullets:</td> <td style="text-align: right;">5 pt</td> </tr> <tr> <td>Subheader:</td> <td style="text-align: right;">6 pt</td> <td>Barlines:</td> <td style="text-align: right;">2.5 pt</td> </tr> <tr> <td>Body Text:</td> <td style="text-align: right;">6 pt</td> <td>Hairlines:</td> <td style="text-align: right;">.5 pt</td> </tr> <tr> <td>Drug Facts (continued):</td> <td style="text-align: right;">7 pt</td> <td></td> <td></td> </tr> </table>	Drug Facts:	8.25 pt	Leading:	6.25 pt	Header:	7 pt	Bullets:	5 pt	Subheader:	6 pt	Barlines:	2.5 pt	Body Text:	6 pt	Hairlines:	.5 pt	Drug Facts (continued):	7 pt		
Drug Facts:	8.25 pt	Leading:	6.25 pt																		
Header:	7 pt	Bullets:	5 pt																		
Subheader:	6 pt	Barlines:	2.5 pt																		
Body Text:	6 pt	Hairlines:	.5 pt																		
Drug Facts (continued):	7 pt																				
Horizontal Scale: 78-80% Average Kerning: 0																					
																					

NOTES:

- 1) [Pattern] NO COPY AREA. COLORS MAY BLEED IN THIS ZONE.
- 2) [Pattern] DATA MATRIX AREA.
- 3) [Pattern] DATA MATRIX QUIET ZONE. NO COPY, NO COLOR.
- 4) [Pattern] LOT & EXP AREA. (TO BE NON-VARNISHED)
- 5) REFERENCE NATIONAL LABEL DRAWING #6-08-13C.

		DVG# NO.		LB-PS-1193		McNeil	
		DRAWN		(b) (4)		Consumer Healthcare	
		SCALE		NONE		1900 Camp Hill Road, Fort Washington, PA 19034	
		DATE		6/30/08		TITLE	
		APPROVED				LABEL, SINGLE PLY	
						75cc BOTTLE	
DIT.	NO.	DATE	REVISIONS				

784443

LOT: [Pattern]

EXP: [Pattern]

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet)
 ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer.

NDC 50580-110-95

Motrin IB

ibuprofen TABLETS USP 200 mg

Pain Reliever /Fever Reducer (NSAID)

125 Coated Caplets *Capsule-Shaped Tablets

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** do not take more than directed; the smallest effective dose should be used. **Adults and children 12 years and older:** take 1 caplet every 4 to 6 hours while symptoms persist; if pain or fever does not respond to 1 caplet, 2 caplets may be used; do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ask a doctor. **Other information:** store between 20°-25°C (68°-77°F); do not use if neck wrap or foil inner seal imprinted "Safety Seal" is broken or missing.

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
 For: MCNEIL PPC, INC., FORT WASHINGTON, PA 19034 USA
 ©MCNEIL-PPC, INC. 2010 www.motrin.com

150% OF SIZE

784443

LOT: [Pattern]

EXP: [Pattern]

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet)
 ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer.

NDC 50580-110-95

Motrin IB


ibuprofen TABLETS USP 200 mg

Pain Reliever /Fever Reducer (NSAID)

125 Coated Caplets *Capsule-Shaped Tablets

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** do not take more than directed; the smallest effective dose should be used. **Adults and children 12 years and older:** take 1 caplet every 4 to 6 hours while symptoms persist; if pain or fever does not respond to 1 caplet, 2 caplets may be used; do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ask a doctor. **Other information:** store between 20°-25°C (68°-77°F); do not use if neck wrap or foil inner seal imprinted "Safety Seal" is broken or missing.

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
 For: MCNEIL PPC, INC., FORT WASHINGTON, PA 19034 USA
 ©MCNEIL-PPC, INC. 2010 www.motrin.com

Description: Lbl Motrin IB Caplet 125 - (b) (4)		Labeling Format Information:	
File #: 3009923	Version: D-1	Fonts: McNeil Condensed Med, Bold, Bold Italic	
Date: 1/5/2010	Contact: (b) (4)	Drug Facts: NA pt	Leading: 5 pt
		Header: 5 pt	Bullets: 4 pt
		Subheader: 5 pt	Barlines: NA pt
		Body Text: 5 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 82-83%	Average Kerning: 0
			


NOTES:

- 1) ALL DIMENSIONS IN INCHES
- 2) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/64
- 3) ---- NO PRINT
- 4) ☒ NO COLOR AREA
- 5) ☒ UPC AREA

				DWG. NO.	McNEIL	
				LBL 48	CONSUMER PRODUCTS CO.	
				DRAWN	CAMP HILL ROAD, FORT WASHINGTON, PA. 19034	
(b) (4)	3	10/6/97	IMPORTED INTO MPSS	(b) (4)	TITLE	
	2	6-1-92	REDRAWN ON NEW FORMAT	1:1	HA LABEL MV	
	1	2-9-90	DIMENSION CHANGE	DATE	5.50 L X 2.00 W	
INIT.	NO.	DATE	REVISIONS	APPROVED		

<p>Important: Read all product information before using. Keep the box for important information.</p> <p>Active ingredient (in each caplet) Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever</p> <p>Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if ■ you have problems or</p>	<p>NDC 50580-110-09</p> <h1>Motrin[®] IB</h1> <p>Ibuprofen Tablets USP, 200 mg Pain Reliever/ Fever Reducer (NSAID) 165 Coated Caplets* *Capsule-Shaped Tablets</p> <p>serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug</p> <p>When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed</p>	<p>Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor</p> <p>Other information ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "Safety Seal[®]" is broken or missing</p> <p>Questions or comments? Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

150% OF SIZE



07844437






<p>Important: Read all product information before using. Keep the box for important information.</p> <p>Active ingredient (in each caplet) Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever</p> <p>Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if ■ you have problems or</p>	<p>NDC 50580-110-09</p> <h1>Motrin[®] IB</h1> <p>Ibuprofen Tablets USP, 200 mg Pain Reliever/ Fever Reducer (NSAID) 165 Coated Caplets* *Capsule-shaped Tablets</p> <p>serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug</p> <p>When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed</p>	<p>Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor</p> <p>Other information ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "Safety Seal[®]" is broken or missing</p> <p>Questions or comments? Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Description: Lbl Motrin IB Caplet 165-LPBulk	
File #: 3009923	Version: R-1
Date: 1/25/10	Contact: (b) (4)


Labeling Format Information:	
Fonts: McNeilDF, Helvet.Cond., reg., bold, &bold italic	
Drug Facts: NA pt	Leading: 5.25 pt
Header: NA pt	Bullets: 4 pt
Subheader: 5 pt	Barlines: NA pt
Body Text: 5 pt	Hairlines: NA pt
Drug Facts (continued): NA pt	
Horizontal Scale: 80% Average Kerning:-11	

McNeil
Consumer Healthcare

NOTES

- 1) BASED ON NATIONAL LABEL DRAWING #: 6-07-5.
- 2)  NO COPY AREA.
- 3)  CLEAR RE-SEAL AREA.
- 4)  LOT & EXP LASER VARNISH AREA.
- 5)  2D DATA MATRIX CODE AREA.
- 6)  DEADENER AREA, TAKES UP ONE OF THE COLORS ON THE BASE LABEL.

DWS NO.		LB-PS-1152		McNeil Consumer Healthcare Camp Hill Road, Fort Washington, PA 19004	
DRAWN:		(b) (4)		TITLE: BOOKLET LABEL	
PLBT:		NONE		175CC, TAB & SINGLE SHEET DESIGN	
SCALE:		6/12/07			
DATE:		6/12/07			
APPROVED:					
(b) (4)	1	6/12/07	INITIAL DRAFT	REVISIONS	



Drug Facts (continued)
Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding
 ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 ■ do not take more than directed
 ■ the smallest effective dose should be used

Drug Facts (continued)
 adults and children 12 years and older ■ take 1 caplet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
 ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
 children under 12 years ■ ask a doctor

Other information
 ■ store between 20-25°C (68-77°F)
 ■ do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing
 ■ see end panel for lot number and expiration date

Inactive ingredients camauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, polyethylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments?
 Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

50101218

BASE LABEL,
PAGE 3



See New Warnings Inside

NDC 50580-110-62

VALUE SIZE!

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Motrin[®] IB
 Ibuprofen Tablets USP, 200 mg
 Pain Reliever / Fever Reducer (NSAID)
 225 Coated Caplets


OPEN TO READ DIRECTIONS AND ADDITIONAL DRUG FACTS INFORMATION
 Do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing.

Drug Facts

Active ingredient (in each caplet)	Purpose
ibuprofen 200 mg (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches
 ■ minor pain of arthritis ■ toothache
 ■ backache ■ the common cold
 ■ menstrual cramps
 ■ temporarily reduces fever

COVER, PAGE 1



Drug Facts (continued)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
 ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if
 ■ you have problems or serious side effects from taking pain relievers or fever reducers
 ■ the stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you have asthma
 ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 ■ under a doctor's care for any serious condition
 ■ taking any other drug



When using this product
 ■ take with food or milk if stomach upset occurs
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

BACK OF COVER,
PAGE 2

Description: Lbl Booklet Motrin IB EZO Caplet 225		Labeling Format Information:	
File #: 3010062	Version: A-2	Fonts: Helvetica Condensed, Bold, Bold Italic	
Date: 04/15/2010	Contact: (b) (4)	Drug Facts: 10 pt	Leading: 6.5 pt
		Header: 8 pt	Bullets: 5 pt
		Subheader: 6 pt	Barlines: 2.5 pt
		Body Text: 6 pt	Hairlines: .5 pt
		Drug Facts (continued): 8 pt	
		Horizontal Scale: 90-100%	Average Kerning: 0

McNeil
Consumer Healthcare

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) ---- NO PRINT
- 5)  LOT & EXP. DATE AREA
- 6)  NO VARNISH AREA

		DWG. NO.		FC-C-114	
		CONSUMER PRODUCTS CO.		CAMP HILL ROAD, FORT WASHINGTON, PA 19034	
(b) (4)	5	4/2/98	CHANGED DWG TITLE TO MATCH	DRAWN (b) (4)	
	4	10/29/97	IMPORTED INTO MPSS	PLOT SCALE 1:1	
	3	1-31-94	ADDED LOT & EXP. DATE AREA	DATE 1-11-88	
	2	4-3-92	REBORN ON NEW FORMAT	TITLE: CARTON, SEAL END 2.44 L X 2.44 W X 4.50 D	
	1	1E-6-89	FLAP(L) AND RADII CORRECTED		
INIT.	NO.	DATE	REVISIONS		

9737376

8737376



3 0045-0481-375

See New Warnings Information

Motrin^{IB}

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

300 Coated Caplets*

*Capsule-Shaped Tablets

NDC 50580 110 37

8737376



08737376

Drug Facts

Active ingredient
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to

- headache
- muscle aches
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include

- hives
- asthma (wheezing)
- asthma (breathing)
- blisters
- facial swelling
- shock
- rash

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take aspirin, blood thinners, or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers

Drug Facts (continued)

Other information

- store between 20-25°C (68-77°F)
- do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing
- see end panel for lot number and expiration date

Inactive ingredients
 carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments?
 Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

Manufactured by: McNeil Healthcare LLC
 LAS PIEDRAS, PUERTO RICO 00771
 For: McNeil Consumer Healthcare
 DIVISION OF MCNEIL-PPC, INC.
 FORT WASHINGTON, PA 19034 USA
 ©MCN-PPC, INC. 2010
www.motrin.com

Drug Facts (continued)

Warnings

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk, if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed


Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you notice a new or worsening rash or hives in the painful area
- any new symptoms appear



If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last trimester of pregnancy because it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed
 ■ the smallest effective dose should be used
 adults and children under 12 years

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used, do not exceed 6 caplets in 24 hours, unless directed by a doctor

Description: Ctn Motrin IB Caplet 300-LPBulk		Labeling Format Information:	
File #: 3009923	Version: S-1	Fonts: McNeilDF, Helvet.Cond., reg., bold, &bold italic	
Date: 1/15/10	Contact: (b) (4)	Drug Facts: 8.5 pt	Leading: 6.5 pt
		Header: 8 pt	Bullets: 5 pt
		Subheader: 6 pt	Barlines: 2.5 pt
		Body Text: 6 pt	Hairlines: .5 pt
		Drug Facts (continued): 8 pt	
		Horizontal Scale: 90%	Average Kerning: 0
			

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP.
- 2) TOLERANCES UNLESS SPECIFIED ± 1/64.
- 3) BASED ON NATIONAL LABEL DRAWING #: 4-07-16A.
- 4) — — NO PRINT AREA BORDER.
- 5)  LOT & EXP DATE AREA. (NO COLOR, NO COPY, NO VARNISH)
- 6)  VERIFICATION CODE AREA. (NO COLOR, NO COPY)

			DWG. NO.	LB-PS-1149	
(b) (4)	4	10/24/07	ADDED TAMPER EVIDENT SLITS	DRAWN:	(b) (4)
	3	10/15/07	ADDED DIMENSIONS TO LOT/EXP	PLT:	NONE
	2	5/22/07	MOVED BARCODE & LOT/EXP	SCALE:	
	1	5/15/07	INITIAL DRAFT	DATE:	5/15/07
INT.	NO.	DATE	REVISIONS	APPROVED:	

McNeil
Consumer Healthcare
Camp Hill Road, Fort Washington, PA 19004

TITLE:
LABEL, SINGLE PLY
CLUB COUNT, 230cc BOTTLE

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis
 ■ toothache ■ backache ■ the common cold
 ■ menstrual cramps

■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach ulcers or bleeding problems. ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

NDC 50580-110-37



When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing. **Inactive ingredients** camauaba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments? Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
 For: McNeil Consumer Healthcare
 DIVISION OF MCNEIL PPC, INC., FORT WASHINGTON, PA 19034 USA
 GMCN-PPC, INC. 2010 www.motrin.com

125% OF SIZE

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis
 ■ toothache ■ backache ■ the common cold
 ■ menstrual cramps

■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach ulcers or bleeding problems. ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

NDC 50580-110-37



When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing. **Inactive ingredients** camauaba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide


Questions or comments? Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771
 For: McNeil Consumer Healthcare
 DIVISION OF MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA
 GMCN-PPC, INC. 2010 www.motrin.com

Description: <u>Lb1 Motrin IB Caplet 300-LPBulk</u>		Labeling Format Information:	
File #: 3009923	Version: T-1	Fonts: MCNeilDF, Helvet.Cond., reg., bold, & bold italic	
Date: 1/22/10	Contact: (b) (4)	Drug Facts: NA pt	Leading: 6.5-7 pt
		Header: 7 pt	Bullets: 5 pt
		Subheader: 6 pt	Barlines: NA pt
		Body Text: 6 pt	Hairlines: NA pt
		Drug Facts (continued): NA pt	
		Horizontal Scale: 80%	Average Kerning: 0
			

NOTES:

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/32
ANGLES ±1°
- 4) — NO PRINT
- 5)  LDT & EXP. DATE AREA
- 6)  NO VARNISH AREA

				DWG. NO.	FC-C-296	 CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA. 19034
				DRAWN	(b) (4)	
(b) (4)	3	12/1/97	IMPORTED INTO MPSS	PLDT SCALE	1:1	TITLE CARTON, SEAL END
	2	1-31-94	ADDED LDT & EXP. DATE AREA	DATE	12-7-89	2.44 L X 2.44 W X 5.50 D
	1	9-9-91	REDRAWN ON NEW FORMAT			
INT.	NO.	DATE	REVISIONS			

08737457

08737457



For Hospital and Government Use Only

See New Warnings Information

NDC 50580 110 51

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)
500 Coated Caplets*

*Capsule-Shaped Tablets

8737457

EXP



For Hospital and Government Use Only

See New Warnings Information

NDC 50580 110 51

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)
500 Coated Caplets*

*Capsule-Shaped Tablets

Motrin[®] IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)
500 Coated Caplets* *Capsule-Shaped Tablets

McNeil

Manufactured by:
McNeil Healthcare LLC
LAS PIEDRAS, PUERTO RICO 00771
For: McNeil Consumer Healthcare
DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
©MCN-PPC, INC. 2010
www.motrin.com

Drug Facts (continued)

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have black or tarry stools
- have dark or bloody urine
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen if you are pregnant because it can harm your unborn baby. Problems in the unborn child or complications during delivery may occur if you use ibuprofen during pregnancy. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms last
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours unless directed by a doctor

children under 12 years

- ask a doctor

Other information

- store between 20-25 °C (68-77 °F)
- keep container tightly closed
- "Safety Seal" is broken or missing
- see and panel for lot number and expiration date

Inactive ingredients

cambeba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments?
Call 1-877-895-3665 weekdays 8:00 AM to 8:00 PM EST


08737457

Description: Ctn Motrin IB Caplet 500 H/G LP Bulk
File #: 3009923 Version: E-1
Date: 01/07/2010 Contact: (b) (4)

Labeling Format Information:




Fonts: McNeil DF Med, Bold, Bold Italic

Drug Facts:	8.25 pt	Leading:	6.5 pt
Header:	6 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	2.5 pt
Body Text:	6 pt	Hairlines:	0.5 pt
Drug Facts (continued):	8 pt		

Horizontal Scale: 100% Average Kerning: 0



NOTES:

- 1) SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED:
FRACTIONS ±1/64
- 4)  NO PRINT
- 5)  NO COLOR AREA
- 6)  VERIFICATION BAR CODE AREA

				DWG. NO. LBL55	McNEIL	
				CONSUMER PRODUCTS CO. CAMP HILL ROAD, FORT WASHINGTON, PA. 19034		
(b) (4)	4	12/30/09	UPDATED PER ENG. REQUEST	DRAWN :	(b) (4)	TITLE: HA LABEL, MV 5.50 L X 3.25 W
	3	10/26/01	VER. CODE AREA WAS UPC AREA	PLOT :	1:1	
	2	9/5/97	IMPORTED INTO MPSS	DATE :	2-20-90	
	1	7-1-92	REDRAWN ON NEW FORMAT	APPROVED :		
INIT.	NO.	DATE	REVISIONS			

Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches
 ■ minor pain of arthritis ■ toothache
 ■ backache ■ the common cold
 ■ menstrual cramps
 ■ temporarily reduces fever


Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
 ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.
 The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if
 ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug


For Hospital and Government Use Only



NDC 50580-110-51

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)



500 Coated Caplets *Capsule-Shaped Tablets

When using this product
 ■ take with food or milk if stomach upset occurs
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area
 ■ any new symptoms appear
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions ■ do not take more than directed ■ the smallest effective dose should be used
Adults and children 12 years and older:
 ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
Children under 12 years: ■ ask a doctor
Other information ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing
Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide
Questions or comments?
 Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

EXP: _____ LOT: _____

Distributed by: McNeil Consumer Healthcare, DIVISION OF McNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA
 ©McNeil-PPC, INC. 2010 www.motrin.com
 00000000

150% OF ACTUAL SIZE FOR REVIEW ONLY



Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches
 ■ minor pain of arthritis ■ toothache
 ■ backache ■ the common cold
 ■ menstrual cramps
 ■ temporarily reduces fever


Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
 ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.
 The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if
 ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug


For Hospital and Government Use Only



NDC 50580-110-51

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)




500 Coated Caplets *Capsule-Shaped Tablets

When using this product
 ■ take with food or milk if stomach upset occurs
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area
 ■ any new symptoms appear
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions ■ do not take more than directed ■ the smallest effective dose should be used
Adults and children 12 years and older:
 ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
Children under 12 years: ■ ask a doctor
Other information ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing
Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide
Questions or comments?
 Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

EXP: _____ LOT: _____

Distributed by: McNeil Consumer Healthcare, DIVISION OF McNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA
 ©McNeil-PPC, INC. 2010 www.motrin.com
 00000000

Description: <u>Lbi Motrin IB Caplet 500 H/G</u> File #: <u>FDA3010135</u> Version: <u>N-1</u> Date: <u>9/07/2010</u> Contact: <u>(b) (4)</u>	Labeling Format Information: Fonts: McN DF Condensed, Bold, Bold Italic <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td>Drug Facts:</td> <td>NA pt</td> <td>Leading:</td> <td>5.50 pt</td> </tr> <tr> <td>Header:</td> <td>6.00 pt</td> <td>Bullets:</td> <td>4.00 pt</td> </tr> <tr> <td>Subheader:</td> <td>5.00 pt</td> <td>Barlines:</td> <td>NA pt</td> </tr> <tr> <td>Body Text:</td> <td>5.00 pt</td> <td>Hairlines:</td> <td>NA pt</td> </tr> <tr> <td>Drug Facts (continued):</td> <td>NA pt</td> <td></td> <td></td> </tr> </table> Horizontal Scale: 75-80% Average Kerning: 0	Drug Facts:	NA pt	Leading:	5.50 pt	Header:	6.00 pt	Bullets:	4.00 pt	Subheader:	5.00 pt	Barlines:	NA pt	Body Text:	5.00 pt	Hairlines:	NA pt	Drug Facts (continued):	NA pt		
Drug Facts:	NA pt	Leading:	5.50 pt																		
Header:	6.00 pt	Bullets:	4.00 pt																		
Subheader:	5.00 pt	Barlines:	NA pt																		
Body Text:	5.00 pt	Hairlines:	NA pt																		
Drug Facts (continued):	NA pt																				



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
10/21/2010