

00000000

EXP

LOT:

Do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

Motrin[®] IB

ibuprofen Tablets USP, 200 mg

24 Coated Caplets Caplet-shaped Tablets

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet)
 ibuprofen 200 mg (NSAID) — Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to
 ■ headache ■ muscular aches ■ minor pain of arthritis

Purposes

■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ dizziness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery ■ **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling, ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** **Directions** ■ **do not take more than directed** ■ **the smallest effective dose should be used.** **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F)

Inactive ingredients carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**, McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©J&JCI 20XX www.motrin.com

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	4.50 pt
Headers:	4.50 pt	Bulletts:	3.50 pt
Subheaders:	4.50 pt	Barlines:	NA pt
		Drug Facts (continued):	NA pt
		Hairlines:	NA pt
			Average Tracking: -16

Reference ID: 4727725

00000000

EXP

LOT:

Do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

TOLL: 800-898-6849

Motrin^{IB}

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

50 Coated Caplets Caplet-shaped Tablets

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each caplet) Purpose
 Ibuprofen 200 mg (NSAID) — Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to
 ■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shortness of breath ■ skin redness/itching ■ rash ■ dizziness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery ■ **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children. Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F)

Inactive ingredients carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**, McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©J&J,CI 20XX www.motrin.com

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	4.50 pt
Headers:	4.50 pt	Bullets:	3.50 pt
Subheaders:	4.50 pt	Leading:	4.50 pt
		Barlines:	NA pt
		Drug Facts (continued):	NA pt
		Hairlines:	NA pt
			Average Tracking: -16

EXP. DATE

LOT :

Do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

NDC 50580-486-0



Motrin IB
Ibuprofen Tablets
USP 200 mg
Pain Reliever
Fever Reducer (NSAID)
60 Coated Caplets

Keep the box for important information.

Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children. Directions** ■ **do not take more than directed** ■ **the smallest effective dose should be used. Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25 °C (68-77°F). **Inactive ingredients** carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**, McKel Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©J&JCI 20XX www.motrin.com

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets: 3.50 pt
Headers:	4.50 pt	Leading:	4.50 pt	Barlines: NA pt
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines: NA pt
				Horizontal Scale: 75%
				Average Tracking: -10

00000000

FPO

LOT:

EXP:

Active ingredient (in each caplet)
 Ibuprofen 200 mg (NSAID)
 *nonsteroidal anti-inflammatory drug

Purposes
 Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache ■ muscular aches ■ minor pain of arthritis ■ toothache
 - backache ■ the common cold ■ menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you

NDC 50580-486-05



Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
100 Coated Caplets* *Capsule-Shaped Tablets

than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ **do not take more than directed** ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist; ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used; ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor

Other information ■ store between 20-25°C (68-77°F) ■ **do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing** ■ inactive ingredients: carmellose xanth, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-995-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McKel Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©J&J/C 200X www.motrin.com

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic					
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt
Headers:	4.50 pt	Leading:	4.50 pt	Barlines:	NA pt
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt
					Horizontal Scale: 75%
					Average Tracking: -10

00000000

Active ingredient (in each caplet)
 ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems. ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease,

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

VALUE SIZE!

NDC 50580-486-06

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)



225 Coated Caplets*

Capulet-Shaped Tablets

actual size

asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product** ■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing. **Inactive ingredients** carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 © & JCI 20XX www.motrin.com

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	4.50 pt
Headers:	4.50 pt	Leading:	5.00 pt
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt
		Bullets:	3.50 pt
		Barlines:	NA pt
		Hairlines:	NA pt
		Horizontal Scale:	83%
		Average Tracking:	-10

00000000

Active ingredient (in each caplet)
 ibuprofen 200 mg (NSAID)
nonsteroidal anti-inflammatory drug

Purposes
 Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this

NDC 50580-486-07

Motrin[®] IB

ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)



300
Coated Caplets* *Capsule-Shaped Tablets

benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product** ■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better. ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor

Other information ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

Inactive ingredients carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©J&JCI 20XX www.motrin.com

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	4.50 pt
Bullets:	3.50 pt	Horizontal Scale:	90%
Headers:	4.50 pt	Leading:	5.00 pt
Barlines:	NA pt	Average Tracking:	-10
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt
Hairlines:	NA pt		

Active ingredient (in each caplet)
 Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Purposes
 temporarily reduces fever

Uses
 temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis
 ■ toothache ■ backache ■ the common cold ■ menstrual cramps

00000000

EXP LOT

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product**

For Hospital and Government Use Only



NDC 50580-486-02

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)

500
Coated Caplets*

**Capsule-Shaped Tablets*



■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better. ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 ■ do not take more than directed ■ the smallest effective dose should be used
Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor

Other information
 ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

Inactive ingredients
 carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©J&JCI 20XX www.motrin.com

Labeling Format Information:				Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	5.00 pt	Bullets:	4.00 pt	Horizontal Scale: 90%	
Headers:	6.00 pt	Leading:	5.50 pt	Barlines:	NA pt		
Subheaders:	5.00 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt	Average Tracking: -5	



Labeling Format Information					
Font:	Helvetica	Cond	Black	Cond	Black
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullet:	3.50 pt
Warnings:	4.50 pt	Leading:	4.00 pt	Barline:	NA pt
Subheaders:	4.50 pt	Drug Facts container:	NA pt	Headline:	NA pt
				Average Tracking:	25



Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 65%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0

Drug Facts (continued)

Do not use if you are allergic to any of the ingredients listed below. Stop use and ask your doctor if you have ever had an allergic reaction to ibuprofen or if you have a history of stomach problems, such as heartburn, ulcers, or bleeding in the stomach.

Do not use if you are pregnant or breastfeeding. It is especially important to use ibuprofen during the last 3 months of pregnancy because it can harm your unborn child or complicate your pregnancy. Do not use if you are pregnant or breastfeeding unless your doctor says it is safe to use.

Do not use if you are taking other medicines. It is especially important to use ibuprofen if you are taking other medicines. Ask your doctor if you are taking other medicines.

Do not use if you are taking other medicines. It is especially important to use ibuprofen if you are taking other medicines.

Do not use if you are taking other medicines. It is especially important to use ibuprofen if you are taking other medicines.

Drug Facts (continued)

Directions

Take 1 caplet every 6 to 8 hours while symptoms persist. Do not take more than directed. Do not take more than 6 caplets in 24 hours. Do not take more than 3 days unless your doctor says it is safe to use. If symptoms do not improve after 3 days, ask your doctor. If symptoms worsen or last more than 10 days, ask your doctor. If symptoms persist for more than 10 days, ask your doctor. If symptoms persist for more than 10 days, ask your doctor.

Warnings

Do not take more than directed. Do not take more than 6 caplets in 24 hours. Do not take more than 3 days unless your doctor says it is safe to use. If symptoms do not improve after 3 days, ask your doctor. If symptoms worsen or last more than 10 days, ask your doctor. If symptoms persist for more than 10 days, ask your doctor.

Uses

Temporarily relieves minor aches and pains due to the common cold, headache, toothache, muscle aches, and menstrual cramps.

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID) — Pain reliever/fever reducer

Purposes

Temporarily relieves minor aches and pains due to the common cold, headache, toothache, muscle aches, and menstrual cramps.

Other Information

Do not use if printed with "Sealed For Your Protection" or "Do not use if broken or missing."

Do not use if you are allergic to any of the ingredients listed below. Stop use and ask your doctor if you have ever had an allergic reaction to ibuprofen or if you have a history of stomach problems, such as heartburn, ulcers, or bleeding in the stomach.

Do not use if you are pregnant or breastfeeding. It is especially important to use ibuprofen during the last 3 months of pregnancy because it can harm your unborn child or complicate your pregnancy. Do not use if you are pregnant or breastfeeding unless your doctor says it is safe to use.

Do not use if you are taking other medicines. It is especially important to use ibuprofen if you are taking other medicines.

Do not use if you are taking other medicines. It is especially important to use ibuprofen if you are taking other medicines.

Do not use if you are taking other medicines. It is especially important to use ibuprofen if you are taking other medicines.

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 80%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



Labeling Format Information: Fonts: HelveticaDF-Medium, Bold, Bold Italic				
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%	
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt		
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0	

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
01/07/2021 11:09:55 AM