

00000000

EXP

LOT:

Do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

Motrin<sup>IB</sup>

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

50 Coated Caplets      Caplets-shaped Tablets

Important: Read all product information before using. Keep the box for important information.

**Active ingredient (in each caplet)**  
Ibuprofen 200 mg (NSAID) — Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ minor pain of arthritis

Purposes

■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shortness of breath ■ skin reddening ■ rash ■ dizziness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery ■ **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children. Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F)

**Inactive ingredients** carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**, McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA  
©J&J, CI 20XX      [www.motrin.com](http://www.motrin.com)

<b>Labeling Format Information:</b> Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic							
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	80%
Headers:	4.50 pt	Leading:	4.50 pt	Barlines:	NA pt	Average Tracking: -16	
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt		

Reference ID: 4889366

00000000

EXP. DATE

LOT :

**Do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing**

**For Hospital and Government Use Only**

NDC 50580-466-0

Motrin IB

Ibuprofen Tablets

USP 200 mg

Pain Reliever

Fever Reducer (NSAID)

60 Coated Caplets

Caplets shaped Tablets

**Keep the box for important information.**

Active ingredient (in each caplet)      Purposes  
 Ibuprofen 200 mg (NSAID)      Pain reliever/fever reducer  
 nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke. ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children. Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25 °C (68-77°F). **Inactive ingredients** carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**, McKee Consumer Healthcare Division, Fort Washington, PA 19034 USA  
 ©J&JCI 20XX      www.motrin.com

<b>Labeling Format Information:</b> Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	4.50 pt
Headers:	4.50 pt	Bullets:	3.50 pt
Subheaders:	4.50 pt	Leading:	4.50 pt
		Barlines:	NA pt
		Drug Facts (continued):	NA pt
		Hairlines:	NA pt
		Average Tracking:	-10

00000000

FPO  
EPC

LOT:  
EXP:

**Active ingredient (in each caplet)**  
 Ibuprofen 200 mg (NSAID)  
 \*nonsteroidal anti-inflammatory drug

**Purposes**  
 Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you



NDC 50580-486-05  
**Motrin IB**  
**Ibuprofen Tablets USP, 200 mg**  
**Pain Reliever/Fever Reducer (NSAID)**  
**100 Coated Caplets** \*Capulet-Shaped Tablets

than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ **do not take more than directed** ■ **the smallest effective dose should be used.** **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist; ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used; ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor

**Other information** ■ store between 20-25°C (68-77°F) ■ **do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing** **Inactive ingredients** carmellose wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call 1-877-895-3665 (ask-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
 McKel Consumer Healthcare Division, Fort Washington, PA 19034 USA  
 ©J&J 20XX [www.motrin.com](http://www.motrin.com)

**Labeling Format Information:** Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	75%
Headers:	4.50 pt	Leading:	4.50 pt	Barlines:	NA pt		
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt	Average Tracking:	-10

00000000

EXP

LOT:

Do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

Motrin<sup>®</sup> IB

ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

24 Coated Caplets Caplet-shaped Tablets

Important: Read all product information before using. Keep the box for important information.

**Active ingredient (in each caplet)**  
 ibuprofen 200 mg (NSAID) — Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches ■ minor pain of arthritis

Purposes

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ dizziness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery ■ **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children. Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F)

**Inactive ingredients** carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**, McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA  
 ©J&JCI 20XX www.motrin.com

<b>Labeling Format Information:</b> Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic							
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	80%
Headers:	4.50 pt	Leading:	4.50 pt	Barlines:	NA pt	Average Tracking: -16	
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt		

Reference ID: 4889366

00000000

**Active ingredient (in each caplet)**  
 ibuprofen 200 mg (NSAID)\*..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache    ■ muscular aches    ■ minor pain of arthritis
  - toothache    ■ backache    ■ the common cold    ■ menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems. ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease,

**VALUE SIZE!**  
NDC 50580-486-06

Motrin<sup>®</sup> IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

225 Coated Caplets\*

Capsule-Shaped Tablets

asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product** ■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing. **Inactive ingredients** carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, stearic acid, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA  
 ©J&JCI 20XX      [www.motrin.com](http://www.motrin.com)

<b>Labeling Format Information:</b> Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	4.50 pt
Bullets:	3.50 pt	Horizontal Scale:	83%
Headers:	4.50 pt	Leading:	5.00 pt
Barlines:	NA pt	Average Tracking:	-10
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt
Hairlines:	NA pt		

00000000

**Active ingredient (in each caplet)**  
Ibuprofen 200 mg (NSAID)<sup>1</sup>.....Pain reliever/fever reducer  
<sup>1</sup>nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache    ■ muscular aches    ■ minor pain of arthritis
  - toothache   ■ backache        ■ the common cold    ■ menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this

NDC 50580-486-07

# Motrin<sup>®</sup> IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)



300  
Coated Caplets\* \*Capsule-Shaped Tablets

benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product** ■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better. ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used. **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor

**Other information** ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

**Inactive ingredients** carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA  
©J&JCI 20XX [www.motrin.com](http://www.motrin.com)

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic							
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	90%
Headers:	4.50 pt	Leading:	5.00 pt	Barlines:	NA pt		
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt	Average Tracking:	-10

**Active ingredient (in each caplet)**  
 Ibuprofen 200 mg (NSAID)\*.....Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Purposes**  
 temporarily reduces fever

**Uses**  
 temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches ■ minor pain of arthritis  
 ■ toothache ■ backache ■ the common cold ■ menstrual cramps

00000000

EXP LOT

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product**

For Hospital and Government Use Only



NDC 50580-486-02

# Motrin<sup>®</sup> IB

**Ibuprofen Tablets USP, 200 mg**  
**Pain Reliever / Fever Reducer (NSAID)**

**500**  
**Coated Caplets\***

*\*Capsule-Shaped Tablets*



■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better. ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**  
 ■ do not take more than directed ■ the smallest effective dose should be used  
**Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor


**Other information**  
 ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing

**Inactive ingredients**  
 carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA  
 ©J&JCI 20XX [www.motrin.com](http://www.motrin.com)

Labeling Format Information:				Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts:	NA pt	Body Text:	5.00 pt	Bullets:	4.00 pt	Horizontal Scale: 90%	
Headers:	6.00 pt	Leading:	5.50 pt	Barlines:	NA pt		
Subheaders:	5.00 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt	Average Tracking: -5	



Labeling Format Information				Font: HelveticaCond-Bold, Cond Black, Cond Black Italic			
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	71%
Headers:	4.50 pt	Leading:	4.00 pt	Barlines:	NA pt		
Subheaders:	4.50 pt	Drug Facts container:	NA pt	Headlines:	NA pt	Average Tracking:	25



Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 65%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 65%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0

**Drug Facts (continued)**

**Directions**

Take 1 caplet every 6 to 8 hours while symptoms persist. Do not take more than 10 days unless your doctor tells you to do so. If symptoms do not improve after 3 days, stop use and ask your doctor.

**Warnings**

**Do not take more than directed**

Do not take more than directed unless directed by a doctor.

**Other information**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

Take 1 caplet every 6 to 8 hours while symptoms persist. Do not take more than 10 days unless your doctor tells you to do so. If symptoms do not improve after 3 days, stop use and ask your doctor.

**Warnings**

**Do not take more than directed**

Do not take more than directed unless directed by a doctor.

**Other information**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Motrin<sup>IB</sup>**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever, Fever Reducer (NSAID)  
100 Coated Caplets

**LOT:** 00000000  
**EXP:** 00000000

OPEN HERE

9 0045-0481-01 3

**Drug Facts (continued)**

**Directions**

Take 1 caplet every 6 to 8 hours while symptoms persist. Do not take more than 10 days unless your doctor tells you to do so. If symptoms do not improve after 3 days, stop use and ask your doctor.

**Warnings**

**Do not take more than directed**

Do not take more than directed unless directed by a doctor.

**Other information**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

Take 1 caplet every 6 to 8 hours while symptoms persist. Do not take more than 10 days unless your doctor tells you to do so. If symptoms do not improve after 3 days, stop use and ask your doctor.

**Warnings**

**Do not take more than directed**

Do not take more than directed unless directed by a doctor.

**Other information**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 80%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0





**Labeling Format Information:** Fonts: Helvetica Medium, Bold, Bold Italic  
 Drug Facts: 9.00 pt | Body Text: 6.00 pt | Bullets: 5.00 pt | Horizontal Scale: 80%  
 Headers: 8.00 pt | Leading: 6.50 pt | Bullets: 2.50 pt  
 Subheaders: 6.00 pt | Drug Facts (continued): 6.00 pt | Headers: 6.50 pt | Average Tacking: -15



Labeling Format Information: Fonts: Helvetica Medium, Bold, Bold Italic			
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 80%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 6.00 pt	Headlines: 6.50 pt	Average Tracking: -15

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

EXP: \_\_\_\_\_

LOT: \_\_\_\_\_

← OPEN HERE

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

**Motrin<sup>®</sup> IB**  
Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
500 Coated Caplets

EXP: \_\_\_\_\_

LOT: \_\_\_\_\_

**Labeling Format Information:** Fonts: Helvetica®/Medium, Bold, Bold Italic

Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0



-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

VALERIE S PRATT  
11/16/2021 03:01:48 PM