

4-ounce Dyna-Hex 4

Lift Here


NDC 0116-1061-04

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-sterile Solution

Contains: 4% Chlorhexidine Gluconate
Distributed By: Xitrium Laboratories, Inc.
Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

For Single-Use



0 11610 61040 4

Net Contents: 4 fl oz (118 mL)

Drug Facts

Active ingredient	Purpose
Chlorhexidine gluconate 4% solution	Antiseptic

Uses

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
- patient preoperative skin preparation: for the preparation of the patient's skin prior to surgery
- skin wound and general skin cleansing

4-ounce Dyna-Hex 4 (Continued)

Drug Facts (continued)

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area
- as a preoperative skin preparation of the head or face

When using this product

- keep out of the eyes, ears and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation:

- apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel

Drug Facts (continued)

- repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol


Questions or comments?

call 1-800-587-3721
Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

8-ounce Dyna-Hex 4

NDC 0116-1061-08	
DYNA-HEX 4[®] Chlorhexidine Gluconate 4% Solution <i>Antiseptic</i>	
Contains:	4% Chlorhexidine Gluconate
Distributed By:	Xttrium Laboratories, Inc. Mount Prospect, IL 60056
	
FOR EXTERNAL USE ONLY	
Net Contents: 8 fl oz (237 mL)	
4D Y08BTL LBLD	
Drug Facts	Purpose
Active ingredient Chlorhexidine gluconate 4% solution	Antiseptic
Uses	
<ul style="list-style-type: none">■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease■ skin wound and general skin cleansing	
Warnings For external use only	
Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none">■ wheezing/difficulty breathing■ shock	
▲ Lift Here	

8-ounce Dyna-Hex 4 (Continued)

Drug Facts (continued)

- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly
- repeat procedure for an additional 2 minutes and dry with a sterile towel

Drug Facts (continued)

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721

Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

16-ounce Dyna-Hex 4

NDC 0116-1061-16


DYNA-HEX 4[®]

Chlorhexidine Gluconate 4% Solution

Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056



0 11610 61160 9

FOR EXTERNAL USE ONLY

Net Contents: 16 fl oz (473 mL)

Drug Facts

Active ingredient
Chlorhexidine gluconate 4% solution

Purpose
Antiseptic

Uses

- surgical hand scrubs: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- health care personnel hand wash: helps reduce bacteria that potentially can cause disease
- skin wound and general skin cleansing

Warnings
For external use only

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- itch
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

▲ Lift Here

4DY1687LLBLD

16-ounce Dyna-Hex 4 (Continued)

<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none">■ If you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product■ In contact with meninges■ In the genital area <p>When using this product</p> <ul style="list-style-type: none">■ Keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.■ If solution should contact these areas, rinse out promptly and thoroughly with water■ Wounds which involve more than the superficial layers of the skin should not be routinely treated■ repeated general skin cleansing of large body areas should not be done except when advised by a health care provider <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none">■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. <p>Surgical hand scrub:</p> <ul style="list-style-type: none">■ wet hands and forearms with water■ scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces■ a separate nail cleaner may be used■ rinse thoroughly■ wash for an additional 3 minutes with 5 mL of product and rinse under running water■ dry thoroughly <p>Healthcare personnel handwash:</p> <ul style="list-style-type: none">■ wet hands with water■ dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds■ rinse and dry thoroughly <p>Skin wound and general skin cleansing:</p> <ul style="list-style-type: none">■ thoroughly rinse the area to be cleaned with water■ apply the minimum amount of product necessary to cover the skin or wound area and wash gently■ rinse again thoroughly <p>Other information</p> <ul style="list-style-type: none">■ store at 20-25°C (68-77°F)■ avoid excessive heat above 40°C (104°F) <p>Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol.</p>

<p>Drug Facts (continued)</p> <p>Questions or comments? call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST</p> <p>Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.</p>

30-ounce Dyna-Hex 4

NDC 0116-1061-30

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate
 Distributed By: Xttrium Laboratories, Inc.
 Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 30 fl oz (1 qt) (887 mL)

To Install

1. Twist to remove plug from back of dispenser spout.
2. Seat dispenser into bracket, taking care to insert the open air inlet into proper receptacle.
3. Twist to remove plug from end of spout.

To Remove

1. Hold bracket against the wall and pull the empty dispenser straight forward.
2. Discard entire bottle and spout.



Drug Facts

Active ingredient Chlorhexidine gluconate 4% solution **Purpose** Antiseptic

Uses
 ■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
 ■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease

Warnings
 For external use only

Allergy alert:
 This product may cause a severe allergic reaction. Symptoms may include:
 ■ wheezing/difficulty breathing
 ■ shock
 ■ facial swelling
 ■ hives
 ■ rash
 If an allergic reaction occurs, stop use and seek medical help right away.

Do not use
 ■ if you are allergic to chlorhexidine gluconate or any other ingredient in this product

When using this product
 ■ keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
 ■ if solution should contact these areas, rinse out promptly and thoroughly with water
 ■ do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.
 If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
 ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
Surgical hand scrub:
 ■ wet hands and forearms with water
 ■ scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
 ■ a separate nail cleaner may be used
 ■ rinse thoroughly
 ■ wash for an additional 3 minutes with 5 mL of product and rinse under running water
 ■ dry thoroughly
Healthcare personnel handwash:
 ■ wet hands with water
 ■ dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
 ■ rinse and dry thoroughly

Other information
 ■ store at 20-25°C (68-77°F)
 ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol.

Questions or comments?
 call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

4DY30BTLLEL

32-ounce Dyna-Hex 4

NDC 0116-1061-32

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate
 Distributed By: Xtrium Laboratories, Inc.
 Mount Prospect, IL 60056



FOR EXTERNAL USE ONLY

Net Contents: 32 fl oz (1 qt) (946 mL)

4DY32BTLBLB

Drug Facts

Active ingredient	Purpose
Chlorhexidine gluconate 4% solution	Antiseptic

Uses

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
- skin wound and general skin cleansing

Warnings
 For external use only

Allergy alert:
 This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient are allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children.
 If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Skin wound and general skin cleansing:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrances, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, triethyl alcohol.

Questions or comments?
 call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.


128-ounce (1-gallon) Dyna-Hex 4

▼ Lift Here for *Drug Facts* NDC 0116-1061-01

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056


0 11610 61010 7

FOR EXTERNAL USE ONLY

Net Contents: 128 fl oz (1 gal) (3.785 L)

Drug Facts

<i>Active ingredient</i>	<i>Purpose</i>
Chlorhexidine gluconate 4% solution	Antiseptic

Uses

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
- skin wound and general skin cleansing


▼

128-ounce (1-gallon) Dyna-Hex 4 (Continued)

<p>Drug Facts (continued)</p> <hr/> <p>Warnings For external use only</p> <hr/> <p>Allergy alert: This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none">• wheezing/difficulty breathing• shock• facial swelling• hives• rash <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <hr/> <p>Do not use</p> <ul style="list-style-type: none">• if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product• in contact with meninges• in the genital area <hr/> <p>When using this product</p> <ul style="list-style-type: none">• keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.• if solution should contact these areas, rinse out promptly and thoroughly with water• wounds which involve more than the superficial layers of the skin should not be routinely treated• repeated general skin cleansing of large body areas should not be done except when advised by a health care provider <hr/> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <hr/> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <hr/> <p>Directions</p> <ul style="list-style-type: none">• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. <p>Surgical hand scrub:</p> <ul style="list-style-type: none">• wet hands and forearms with water• scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces• a separate nail cleaner may be used• rinse thoroughly• wash for an additional 3 minutes with 5 mL of product and rinse under running water• dry thoroughly
--

<p>Drug Facts (continued)</p> <hr/> <p>Healthcare personnel handwash:</p> <ul style="list-style-type: none">• wet hands with water• dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds• rinse and dry thoroughly <p>Skin wound and general skin cleansing:</p> <ul style="list-style-type: none">• thoroughly rinse the area to be cleaned with water• apply the minimum amount of product necessary to cover the skin or wound area and wash gently• rinse again thoroughly <hr/> <p>Other information</p> <ul style="list-style-type: none">• store at 20-25°C (68-77°F)• avoid excessive heat above 40°C (104°F) <hr/> <p>Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol</p> <hr/> <p>Questions or comments? call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST</p> <hr/> <p>Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.</p>

15-mL Dyna-Hex 4 (carton)-Full view



Contains 4% Chlorhexidine Gluconate
 Distributed By: Xtrium Laboratories, Inc.
 Mount Prospect, IL 60056
 FOR EXTERNAL USE ONLY

Drug Facts
Active ingredient Purpose
 Chlorhexidine gluconate Antiseptic

Uses

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and lowers risk to surgery or patient care
- health care personnel handwash: helps reduce bacteria that primarily on a case closure
- patient preoperative skin preparation: for the preparation of the patient's skin prior to surgery
- skin wound and general skin cleansing

Warnings
 For external use only

Allergy alert
 This product may cause a severe allergic reaction. Symptoms may include:

- severe respiratory breathing
- shock
- local swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with mucous
- in the genital area
- as a preoperative skin preparation of the hand or face

Drug Facts (continued)

When using this product

- keep out of the eyes, ears and mouth. Like all other antiseptics and preservatives eye injury is possible if used in the eye during surgical procedures or any cause contact with mucus in the mouth. Rinse through profusely with water.
- if contact should occur between areas, rinse out completely and thoroughly with water.
- wounds which involve more than the superficial layers of the skin should not be treated with this product.
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider.

Stop use and ask a doctor (178166), pharmacist, or other health care worker.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation of chemical nature.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product as if a wet brush paying close attention to the nails, wrists, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Health care personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Drug Facts (continued)

Patient preoperative skin preparation:

- apply product liberally to surgical site and allow for at least 3 minutes
- dry with a sterile towel
- repeat procedure for an additional 3 minutes and dry with a sterile towel

Wound and general skin cleansing:

- thoroughly rinse the area to be cleaned with water
- apply this minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
 cocaine HCl, butyrate, glucose, beta-alanine, hydroxyethylcellulose, isopropyl alcohol, lauric acid, PEG-30, benzalkonium chloride, benzyl alcohol

Directions or comments?
 Call 1-800-800-3721
 like any through Ffany
 600 Ask to 650 PARCET

ADVERSE REACTIONS

Labeling/Cloning Instructions:
 Chlorhexidine gluconate skin cleansers will cause irritation if used with chlorine-releasing products, such as chlorine bleach. Please completely avoid use only on non-chlorine cleansers.

List number and expiration date printed on side of box.

NDC 0116-1061-15

DYNA-HEX 4[®]
 Chlorhexidine Gluconate 4% Solution
 Antiseptic
 Non-Sterile Solution

Contains: 4% Chlorhexidine Gluconate
 Distributed By: Xtrium Laboratories, Inc.
 Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes



0 11610 61150 0

NDC 0116-1061-15

DYNA-HEX 4[®]
 Chlorhexidine Gluconate 4% Solution
 Antiseptic
 Non-Sterile Solution

Contains: 4% Chlorhexidine Gluconate
 Xtrium Laboratories, Inc.
 Distributed By: Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes

NDC 0116-1061-15

DYNA-HEX 4[®]
 Chlorhexidine Gluconate 4% Solution
 Antiseptic
 Non-Sterile Solution

Contains: 4% Chlorhexidine Gluconate
 Distributed By: Xtrium Laboratories, Inc.
 Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes

LOT# 1061-470B
 EXP 01/2018

15-mL Dyna-Hex 4 (carton)-Enlargement of Drug Facts panel (Continued)

<p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredient</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Chlorhexidine gluconate 4% solution</td> <td>Antiseptic</td> </tr> </tbody> </table>	Active ingredient	Purpose	Chlorhexidine gluconate 4% solution	Antiseptic	<p>Drug Facts (continued)</p> <p>When using this product</p> <ul style="list-style-type: none"> keep out of the eyes, ears and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums. if solution should contact these areas, rinse out promptly and thoroughly with water wounds which involve more than the superficial layers of the skin should not be routinely treated repeated general skin cleansing of large body areas should not be done except when advised by a health care provider <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Patient preoperative skin preparation:</p> <ul style="list-style-type: none"> apply product liberally to surgical site and swab for at least 2 minutes dry with a sterile towel repeat procedure for an additional 2 minutes and dry with a sterile towel <p>Skin wound and general skin cleansing:</p> <ul style="list-style-type: none"> thoroughly rinse the area to be cleaned with water apply the minimum amount of product necessary to cover the skin or wound area and wash gently rinse again thoroughly
Active ingredient	Purpose					
Chlorhexidine gluconate 4% solution	Antiseptic					
<p>Uses</p> <ul style="list-style-type: none"> surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care healthcare personnel handwash: helps reduce bacteria that potentially can cause disease patient preoperative skin preparation: for the preparation of the patient's skin prior to surgery skin wound and general skin cleansing 	<p>Directions</p> <ul style="list-style-type: none"> use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. <p>Surgical hand scrub:</p> <ul style="list-style-type: none"> wet hands and forearms with water scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces a separate nail cleaner may be used rinse thoroughly wash for an additional 3 minutes with 5 mL of product and rinse under running water dry thoroughly <p>Healthcare personnel handwash:</p> <ul style="list-style-type: none"> wet hands with water dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds rinse and dry thoroughly 	<p>Other information</p> <ul style="list-style-type: none"> store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F) 				
<p>Warnings</p> <p>For external use only</p> <p>Allergy alert:</p> <p>This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none"> wheezing/difficulty breathing shock facial swelling hives rash <p>If an allergic reaction occurs, stop use and seek medical help right away.</p>	<p>Do not use</p> <ul style="list-style-type: none"> if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product in contact with meninges in the genital area as a preoperative skin preparation of the head or face 	<p>Inactive ingredients</p> <p>cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol</p>				
		<p>Questions or comments?</p> <p>call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST</p> <p>4DY15CTNLBLA</p> <p>Laundering/Cleaning Instructions:</p> <p>Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.</p> <p>Lot number and expiration date printed on side of box</p>				

15-mL Dyna-Hex 4 (carton)-Enlargement of principal display panel (Continued)

NDC 0116-1061-15

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-Sterile Solution



Contains: 4% Chlorhexidine Gluconate
Xttrium Laboratories, Inc.

Distributed By: Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes

15-mL Dyna-Hex 4 (packet)



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
09/14/2017