

NDC 63323-269-50 260950  
**DIPRIVAN**<sup>®</sup>  
(Propofol) INJECTABLE  
EMULSION, USP

**500 mg per 50 mL**  
(10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION

**SHAKE WELL BEFORE USING**

Twenty 50 mL vials

For Single Patient Use Only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only



FOR INTRAVENOUS ADMINISTRATION  
**SHAKE WELL BEFORE USING**  
Twenty 50 mL vials  
For Single Patient Use Only

NDC 63323-269-50 260950  
**DIPRIVAN**<sup>®</sup>  
(Propofol) INJECTABLE  
EMULSION, USP  
**500 mg per 50 mL**  
(10 mg per mL)

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only



LOT  
EXP

DIPRIVAN is a trademark of  
APP Pharmaceuticals, LLC.  
Manufactured for:  
**APP Pharmaceuticals, LLC**  
Schaumburg, IL 60173  
Made in Austria

Rx only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

**Sterile, nonpyrogenic**  
**SHAKE WELL BEFORE USING**  
Dosage: See package insert.  
In addition to the active component, propofol, the formulation contains: soybean oil (100 mg/mL), glycerol (22.5 mg/mL), egg lecithin (12 mg/mL) and disodium edetate (0.005%); with sodium hydroxide to adjust pH.  
DIPRIVAN<sup>®</sup> should be administered only by persons trained in the administration of general anesthetics and not involved in the conduct of the surgical/diagnostic procedure. Facilities for maintenance of a patent airway, artificial ventilation, and oxygen enrichment and circulatory resuscitation must be immediately available.  
Store between 4° to 25°C (40° to 77°F). Do not freeze.

M027818/01  
62934E

NDC 63323-269-50 260950  
**DIPRIVAN**<sup>®</sup>  
(Propofol) INJECTABLE  
EMULSION, USP  
**500 mg per 50 mL**  
(10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION  
**SHAKE WELL BEFORE USING**  
Twenty 50 mL vials  
For Single Patient Use Only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only



2 03 63323 269 50 4



NDC 63323-269-50 260950

**DIPRIVAN**<sup>®</sup>  
(Propofol) INJECTABLE  
EMULSION, USP

**500 mg per 50 mL**  
(10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION

**SHAKE WELL BEFORE USING**


Twenty 50 mL vials

For Single Patient Use Only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only





30363323269501

Manufactured for:  
**APP Pharmaceuticals, LLC**  
Schaumburg, IL 60173  
Made in Austria

US Pat 5,714,520 5,731,355  
5,731,356 5,908,869

402347C  
M087920/01



- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only

*Sterile, nonpyrogenic*  
**SHAKE WELL BEFORE USING**

**Dosage:** See package insert.  
In addition to the active component, propofol, the formulation contains: soybean oil (100 mg/mL), glycerol (22.5 mg/mL), egg lecithin (12 mg/mL) and disodium edetate (0.005%); with sodium hydroxide to adjust pH. Store between 4° to 25°C (40° to 77°F). Do not freeze.

NDC 63323-269-50 260950

**DIPRIVAN<sup>®</sup>**  
*(Propofol)* INJECTABLE  
EMULSION, USP

**500 mg per 50 mL**  
(10 mg per mL)

**FOR INTRAVENOUS ADMINISTRATION**  
**50 mL - For Single Patient Use Only**

LOT  
EXP

NDC 63323-269-65 260965  
**DIPRIVAN<sup>®</sup>**  
(Propofol) INJECTABLE  
EMULSION, USP

**1000 mg per 100 mL**  
(10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION

**SHAKE WELL BEFORE USING**

Ten 100 mL vials

For Single Patient Use Only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only



NDC 63323-269-65 260965  
**DIPRIVAN<sup>®</sup>**  
(Propofol) INJECTABLE  
EMULSION, USP

**1000 mg per 100 mL**  
(10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION

**SHAKE WELL BEFORE USING**

Ten 100 mL vials

For Single Patient Use Only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only

LOT  
EXP

DIPRIVAN is a trademark of  
APP Pharmaceuticals, LLC.  
Manufactured for:  
APP Pharmaceuticals, LLC.  
Schaumburg, IL 60173  
Made in Austria

Rx only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

**Sterile, nonpyrogenic**  
**SHAKE WELL BEFORE USING**  
Dosage: See package insert.  
In addition to the active component, propofol, the formulation contains: soybean oil (100 mg/mL), glycerol (22.5 mg/mL), egg lecithin (12 mg/mL), and disodium edetate (0.005%); with sodium hydroxide to adjust pH. DIPRIVAN<sup>®</sup> should be administered only by persons trained in the administration of general anesthesia and not involved in the conduct of the surgical/diagnostic procedure. Facilities for maintenance of a patent airway, artificial ventilation, and oxygen enrichment and circulatory resuscitation must be immediately available. Store between 4° to 25°C (40° to 77°F). Do not freeze.



2 03 63323 269 65 8



Rx only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

FOR INTRAVENOUS ADMINISTRATION

**SHAKE WELL BEFORE USING**

Ten 100 mL vials

**1000 mg per 100 mL**  
(10 mg per mL)

NDC 63323-269-65 260965  
**DIPRIVAN<sup>®</sup>**  
(Propofol) INJECTABLE  
EMULSION, USP



NDC 63323-269-65 260965  
**DIPRIVAN<sup>®</sup>**  
(Propofol) INJECTABLE  
EMULSION, USP

**1000 mg per 100 mL**  
(10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION

**SHAKE WELL BEFORE USING**

Ten 100 mL vials

For Single Patient Use Only

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only

M027820/01  
62936E