

NOVARTIS NDC 0078-0180-01 Unit Dose Package

50 mcg/mL; Each mL contains:
octreotide (as acetate) 50 mcg
Inactive ingredients
lactic acid, USP 3.4 mg
mannitol, USP 45 mg
sodium bicarbonate, USP qs to pH 4.2 ± 0.3
water for injection, USP qs to 1 mL

See package insert for dosage and administration information.

Storage: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light. At room temperature, (20°C to 30°C or 70°F to 86°F), Sandostatin is stable for 14 days if protected from light.

Sandostatin®
octreotide acetate INJECTION
50 mcg/mL

10 Ampuls/1 mL size

Instructions for Use

One-point-cut ampul with cut below colored point

To open, hold as shown with thumb above point and snap off backwards


Manufactured by:
Novartis Pharma Stein AG
Schaffhauserstrasse
CH-4332 Stein, Switzerland

Distributed by:
Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936

©Novartis 1256727

NDC 0078-0180-01 Unit Dose Package
10 Ampuls/1 mL size

120 x 105 x 20.5
US
1256727



NOVARTIS NDC 0078-0180-01 Unit Dose Package

Sandostatin®
octreotide acetate
INJECTION

Rx only

50 mcg/mL
For Subcutaneous Injection

10 Ampuls/1 mL size

LOT
EXP.



Please open here  Please open here

NDC 0078-0181-01
Unit Dose Package
10 Ampuls/1 mL size

 **NOVARTIS**

NDC 0078-0181-01
Unit Dose Package

100 mcg/mL; Each mL contains:
octreotide (as acetate) 100 mcg
Inactive ingredients
lactic acid, USP 3.4 mg
mannitol, USP 45 mg
sodium bicarbonate, USP qs to pH 4.2 ± 0.3
water for injection, USP qs to 1 mL

See package insert for dosage and administration information.

Storage: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light. At room temperature, (20°C to 30°C or 70°F to 86°F), Sandostatin is stable for 14 days if protected from light.



Instructions for Use



One-point-cut ampul with cut below colored point



To open, hold as shown with thumb above point and snap off backwards

Manufactured by:
Novartis Pharma Stein AG
Schaffhauserstrasse
CH-4332 Stein, Switzerland

Distributed by:
Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936

©Novartis

1256728

 **Sandostatin**[®]
octreotide acetate INJECTION
100 mcg/mL

120 x 105 x 20.5
US
1256728

NDC 0078-0181-01
Unit Dose Package
10 Ampuls/1 mL size

 **Sandostatin**[®]
octreotide acetate INJECTION
100 mcg/mL

EXP.
LOT

 **NOVARTIS**

NDC 0078-0181-01
Unit Dose Package

 **Sandostatin**[®]
octreotide acetate
INJECTION

Rx only

100 mcg/mL
For Subcutaneous Injection

10 Ampuls/1 mL size



Please open here  Please open here

NDC 0078-0182-01
Unit Dose Package

10 Ampuls/1 mL size

 **NOVARTIS**

NDC 0078-0182-01
Unit Dose Package

500 mcg/mL; Each mL contains:
octreotide (as acetate) 500 mcg
Inactive ingredients
lactic acid, USP 3.4 mg
mannitol, USP 45 mg
sodium bicarbonate, USP qs to pH 4.2 ± 0.3
water for injection, USP qs to 1 mL

See package insert for dosage and administration information.

**Storage: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.
At room temperature, (20°C to 30°C or 70°F to 86°F), Sandostatin is
stable for 14 days if protected from light.**



Instructions for Use



One-point-cut ampul with
cut below colored point



To open, hold as shown
with thumb above point
and snap off backwards

Manufactured by:
Novartis Pharma Stein AG
Schaffhauserstrasse
CH-4332 Stein, Switzerland

Distributed by:
Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936

©Novartis

1256731


Sandostatin[®]
octreotide acetate INJECTION
500 mcg/mL

120 x 105 x 20.5
US

NDC 0078-0182-01
Unit Dose Package
10 Ampuls/1 mL size


Sandostatin[®]
octreotide acetate INJECTION
500 mcg/mL

LOT
EXP.

 **NOVARTIS**

NDC 0078-0182-01
Unit Dose Package

 **Sandostatin**[®]
octreotide acetate
INJECTION

Rx only

500 mcg/mL
For Subcutaneous Injection

10 Ampuls/1 mL size



Label may not be the latest approved by
For more information, please visit <https://www.>

Sandostatin[®]

NDC 0078-0183-25

octreotide acetate

Rx only

INJECTION

1,000 mcg/5 mL (200 mcg/mL)

Total Volume 5 mL Multi-Dose Vial

Each mL of aqueous solution contains: **octreotide (as acetate) 200 mcg**

FOR SUBCUTANEOUS INJECTION

STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.

After initial use, discard within 14 days.

Mfd. by Novartis Pharma Stein AG

©Novartis

1256734B

US

256734A

1256734B



EXP/LOT

Sandostatin[®]

NDC 0078-0183-25

octreotide acetate

Rx only

INJECTION

1,000 mcg/5 mL (200 mcg/mL)

Total Volume 5 mL Multi-Dose Vial

Each mL of aqueous solution contains: **octreotide (as acetate) 200 mcg**

FOR SUBCUTANEOUS INJECTION

STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.

After initial use, discard within 14 days.

Mfd. by Novartis Pharma Stein AG

©Novartis

1256734B

US

256734A

1256734B



EXP/LOT

Sandostatin[®]

NDC 0078-0183-25

octreotide acetate

Rx only

INJECTION

1,000 mcg/5 mL (200 mcg/mL)

Total Volume 5 mL Multi-Dose Vial

Each mL of aqueous solution contains: **octreotide (as acetate) 200 mcg**

FOR SUBCUTANEOUS INJECTION

STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.

After initial use, discard within 14 days.

Mfd. by Novartis Pharma Stein AG

©Novartis

1256734B

US



256734A

1256734B



EXP/LOT

Please open here  Please open here



NDC 0078-0183-25

Sandostatin[®]
octreotide acetate
INJECTION

NOVARTIS

Total Volume **5 mL**
 Multi-Dose Vial

1,000 mcg/5 mL (200 mcg/mL)
 FOR SUBCUTANEOUS INJECTION



NOVARTIS

NDC 0078-0183-25

Sandostatin[®]

octreotide acetate INJECTION

Each mL of aqueous solution contains:

octreotide (as acetate)	200 mcg
Inactive ingredients:	
lactic acid, USP	3.4 mg
mannitol, USP	45 mg
phenol, USP	5.0 mg
sodium bicarbonate, USP	qs to pH 4.2 ± 0.3
water for injection, USP	qs to 1 mL

DOSAGE: See package insert for dosage information.

STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light. After initial use, discard within 14 days. At room temperature, (20°C to 30°C or 70°F to 86°F), Sandostatin is stable for 14 days if protected from light.



0078-0183-25 3

Manufactured by: Novartis Pharma Stein AG
 Schaffhauserstrasse, CH-4332 Stein, Switzerland

Distributed by:
 Novartis Pharmaceuticals Corporation
 East Hanover, New Jersey 07936

©Novartis

1256736

Sandostatin[®]
octreotide acetate
INJECTION



NOVARTIS

1,000 mcg/5 mL (200 mcg/mL)
 FOR SUBCUTANEOUS INJECTION

Total Volume **5 mL**
 Multi-Dose Vial

NDC 0078-0183-25

96 x 73 x 22.5
 1256736
 US



LOT:
 EXP:



NOVARTIS

NDC 0078-0183-25



Sandostatin[®]

octreotide acetate

INJECTION

Rx only

STORE REFRIGERATED AT 2°C to 8°C (36°F to 46°F); PROTECT FROM LIGHT. AFTER INITIAL USE, DISCARD WITHIN 14 DAYS

1,000 mcg/5 mL (200 mcg/mL)

FOR SUBCUTANEOUS INJECTION
 Total Volume **5 mL** Multi-Dose Vial



1256736

Label may not be the latest approved by
For more information, please visit <https://www.>

Sandostatin[®]

octreotide acetate

NDC 0078-0184-25

Rx only

INJECTION

5,000 mcg/5 mL (1,000 mcg/mL)

Total Volume 5 mL Multi-Dose Vial

Each mL of aqueous solution contains: **octreotide (as acetate) 1,000 mcg**

FOR SUBCUTANEOUS INJECTION

STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.

After initial use, discard within 14 days.

Mfd. by Novartis Pharma Stein AG

© Novartis

12567400 US 12567400 12567400



EXP/LOT

Sandostatin[®]

octreotide acetate

NDC 0078-0184-25

Rx only

INJECTION

5,000 mcg/5 mL (1,000 mcg/mL)

Total Volume 5 mL Multi-Dose Vial

Each mL of aqueous solution contains: **octreotide (as acetate) 1,000 mcg**

FOR SUBCUTANEOUS INJECTION

STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.

After initial use, discard within 14 days.

Mfd. by Novartis Pharma Stein AG

© Novartis

12567400 US 12567400 12567400



EXP/LOT

Sandostatin[®]

octreotide acetate

NDC 0078-0184-25

Rx only

INJECTION

5,000 mcg/5 mL (1,000 mcg/mL)

Total Volume 5 mL Multi-Dose Vial

Each mL of aqueous solution contains: **octreotide (as acetate) 1,000 mcg**

FOR SUBCUTANEOUS INJECTION

STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.

After initial use, discard within 14 days.

Mfd. by Novartis Pharma Stein AG

© Novartis

12567400 US 12567400 12567400



EXP/LOT

Please open here  Please open here


NDC 0078-0184-25

Sandostatin[®]
octreotide acetate
INJECTION

NOVARTIS

5,000 mcg/5 mL (1,000 mcg/mL)
FOR SUBCUTANEOUS INJECTION

Total Volume 5 mL
Multi-Dose Vial

 **NOVARTIS** NDC 0078-0184-25

Sandostatin[®]
octreotide acetate INJECTION
Each mL of aqueous solution contains:
octreotide (as acetate) 1,000 mcg
Inactive ingredients:
lactic acid, USP 3.4 mg
mannitol, USP 45 mg
phenol, USP 5.0 mg
sodium bicarbonate, USP qs to pH 4.2 ± 0.3
water for injection, USP qs to 1 mL

DOSAGE: See package insert for dosage information.
STORAGE: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light.
After initial use, discard within 14 days. At room temperature, (20°C to 30°C or 70°F to 86°F), Sandostatin is stable for 14 days if protected from light.



Manufactured by: Novartis Pharma Stein AG
Schaffhauserstrasse, CH-4332 Stein, Switzerland

Distributed by:
Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936 ©Novartis 1256743

 **NOVARTIS**

 **Sandostatin[®]**
octreotide acetate
INJECTION

5,000 mcg/5 mL (1,000 mcg/mL)
FOR SUBCUTANEOUS INJECTION

Total Volume 5 mL
Multi-Dose Vial

NDC 0078-0184-25

96 x 73 x 32.5
1256743
US

LOT
EXP

 **NOVARTIS** NDC 0078-0184-25

 **Sandostatin[®]**
octreotide acetate
INJECTION

Rx only

STORE REFRIGERATED AT 2°C to 8°C (36°F to 46°F); PROTECT FROM LIGHT. AFTER INITIAL USE, DISCARD WITHIN 14 DAYS

5,000 mcg/5 mL (1,000 mcg/mL)
FOR SUBCUTANEOUS INJECTION
Total Volume 5 mL Multi-Dose Vial



This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

NDC 0781-3166-95

OCTREOTIDE Acetate Injection

50 mcg/mL

For Subcutaneous Injection

Rx only 10 x 1 mL Single-Dose Vials

 **SANDOZ**
a Novartis company

WSB107D1R1

UPC CODE
AREA



SOLIDE
BLOCK NOIR

NDC 0781-3166-95
**OCTREOTIDE
Acetate Injection**
50 mcg/mL
Rx only
10 x 1 mL Single-Dose Vials
 **SANDOZ**
a Novartis company

Each mL of aqueous solution contains:
Octreotide (as acetate)..... 50 mcg
Inactive ingredients:
Lactic Acid, USP..... 3.4 mg
Mannitol, USP..... 45 mg
Sodium Bicarbonate..... qs to pH 4.2 ± 0.3
USP
Water for Injection, USP..... qs to 1 mL
Dosage: See package insert for dosage information.
Sandoz Canada Inc. for
Sandoz Inc., Princeton, NJ 08540
01-2015M
THE REACH OF CHILDREN.
KEEP THIS AND ALL DRUGS OUT OF
for 14 days if protected from light.
Lactic Acid, USP..... 3.4 mg
Mannitol, USP..... 45 mg
Sodium Bicarbonate..... qs to pH 4.2 ± 0.3
USP
Water for Injection, USP..... qs to 1 mL
Dosage: See package insert for dosage information.
Sandoz Canada Inc. for
Sandoz Inc., Princeton, NJ 08540
01-2015M
THE REACH OF CHILDREN.
KEEP THIS AND ALL DRUGS OUT OF
for 14 days if protected from light.

NDC 0781-3166-95

OCTREOTIDE Acetate Injection

50 mcg/mL

For Subcutaneous Injection

Rx only 10 x 1 mL Single-Dose Vials


 **SANDOZ**
a Novartis company

VARNISH
FREE AREA (4)

VARNISH
FREE AREA

XXXXXX

XXXX



NDC 0781-3167-95

OCTREOTIDE Acetate Injection

100 mcg/mL

For Subcutaneous Injection
Rx only 10 x 1 mL Single-Dose Vials



WSB107D1R1

UPC CODE
AREA



SOLIDE
BLOCK NOIR

NDC 0781-3167-95
**OCTREOTIDE
Acetate Injection**
100 mcg/mL
Rx only
10 x 1 mL Single-Dose Vials
SANDOZ
a Novartis company

Each mL of aqueous solution contains:
Octreotide (as acetate)..... 100 mcg
Inactive ingredients:
Lactic Acid, USP..... 3.4 mg
Mannitol, USP..... 45 mg
Sodium Bicarbonate..... 45 mg
USP..... qs to pH 4.2 ± 0.3
Water for Injection, USP..... qs to 1 mL
Dosage: See package insert for dosage information.
Sandoz Canada Inc. for
Sandoz Inc., Princeton, NJ 08540
Manufactured in Canada by
01-2015M
REACH OF CHILDREN.
KEEP THIS AND ALL DRUGS OUT OF THE
for 14 days if protected from light.
70°F to 86°F). Octreotide Acetate is stable
to room temperature, (20°C to 30°C or
to 46°F). Protect from light.
Store refrigerated between 2°C to 8°C (36°F

NDC 0781-3167-95

OCTREOTIDE Acetate Injection

100 mcg/mL

For Subcutaneous Injection
Rx only 10 x 1 mL Single-Dose Vials




VARNISH
FREE AREA (4)

VARNISH
FREE AREA

XXXXXX

XXXXX



Label may not be the latest approved by
information, please visit <https://www>.

NDC 10781-3165-75	Each mL of aqueous solution contains: Octreotide (as acetate) 200 mcg	XXXXXX
OCTREOTIDE Acetate Injection	Storage: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light. After initial use, discard within 14 days.	
1,000 mcg/5 mL (200 mcg/mL)	01-2015M	
For Subcutaneous Injection Rx only Total Volume 5 mL Multi-Dose Vial	Manufactured in Canada by Sandoz Canada Inc. for Sandoz Inc. Princeton, NJ 08540	Lot: Exp.:
 SANDOZ		

NON-PRINT MARGIN 1/16

RSS 10307813165757

PRINTED SIDE

XXXXXXX

**OCTREOTIDE
Acetate Injection**
For
Subcutaneous
Injection
1,000 mcg/5 mL
(200 mcg/mL)
Total Volume
5 mL Multi-
Dose Vial
Rx only
SANDOZ
a Novartis company

Each mL of aqueous solution contains:
Octreotide
(as acetate)200 mcg

Inactive Ingredients:
Lactic Acid, USP 3.4 mg
Mannitol, USP 45 mg
Phenol, USP5.0 mg
Sodium Bicarbonate,
USPqs to pH 4.2 ± 0.3
Water for Injection,
USPqs to 1 mL

DOSAGE: See package insert for dosage information.

Storage: Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light. After initial use, discard within 14 days.

At room temperature, (20°C to 30°C or 70°F to 86°F), Octreotide Acetate is stable for 14 days if protected from light.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

NDC 0781-3165-75

OCTREOTIDE Acetate Injection

**1,000 mcg/5 mL
(200 mcg/mL)**

For
Subcutaneous
Injection

Rx only
Total Volume 5 mL
Multi-Dose Vial

SANDOZ
a Novartis company

Manufactured in Canada by Sandoz Canada Inc. for Sandoz Inc. Princeton, NJ 08540

01-2015M

SANDOZ
a Novartis company

UPC CODE
AREA

128 CODE
AREA



(b) (4)

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

