

BACK

FRONT

Advil

COLD & SINUS Non-Drowsy

Ibuprofen 200 mg > Pain Reliever/Fever Reducer (NSAID)
Pseudoephedrine HCl 30 mg > Nasal Decongestant

Relieves Sinus Pressure,
Nasal Congestion and Fever



20 COATED CAPLETS*
*Oval-Shaped Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

<p>Drug Facts (continued)</p> <p>Active ingredients (in each caplet) Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer Pseudoephedrine HCl 30 mg Nasal decongestant</p> <p>Purposes *Nonsteroidal anti-inflammatory drug</p>	<p>Drug Facts (continued)</p> <p>Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing) stomach pain, skin redness, rash, blisters If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach ulcers or bleeding problems. Take a blood thinner (anticoagulant) or steroid drug. Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others). Have 5 or more alcoholic drinks every day while using this product. Take more or for a longer time than directed.</p> <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if stomach bleeding warning applies to you or you have problems or serious side effects from taking pain relievers or fever reducers. You have a history of stomach problems, such as heartburn, asthma, thyroid disease, diabetes, liver cirrhosis, kidney disease, or you have high blood pressure, heart disease, liver cirrhosis, kidney disease, prostate gland, or had a stroke. You are taking a diuretic.</p> <p>Drug Facts (continued) Avoid excessive heat above 40°C (104°F). Read all warnings and directions before use. Keep carton.</p>
<p>Drug Facts (continued)</p> <p>Other information ■ Store at 20-25°C (68-77°F); excursions permitted to 15°-30°C (59°-86°F) (see USP Controlled Room Temperature). ■ Keep this and all other medicines away from children. ■ See the back of the carton for a list of inactive ingredients.</p>	<p>Drug Facts (continued)</p> <p>Directions ■ Do not take more than directed. ■ The smallest effective dose should be used. ■ Adults and children 12 years of age and over: Take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used. ■ Do not use more than 6 caplets in any 24-hour period unless directed by a doctor.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless expressly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: ■ feel faint, ■ vomit blood, ■ have bloody or black stools ■ you have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain, ■ trouble breathing, ■ weakness in one part or side of body, ■ slurred speech, ■ leg swelling</p> <p>When using this product ■ Take with food or milk if stomach upset occurs ■ taking any other drug ■ decrease this benefit of aspirin</p> <p>Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking any other product that contains pseudoephedrine or any other nasal decongestant ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</p>

UNPRINTED AREA FOR LOT & EXP CODING (UNVARNISHED)

Advil

COLD & SINUS Non-Drowsy

Ibuprofen 200 mg > Pain Reliever/Fever Reducer (NSAID)
Pseudoephedrine HCl 30 mg > Nasal Decongestant

Relieves Sinus Pressure,
Nasal Congestion and Fever



40
COATED
CAPLETS*

*Oval-Shaped Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredients (in each caplet) **Purposes**

Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer
Pseudoephedrine HCl 30 mg Nasal decongestant

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives ■ facial swelling ■ asthma (wheezing)
shock ■ skin redness ■ rash ■ blisters

Uses temporarily relieves these symptoms associated with the common cold or flu: ■ headache ■ fever ■ sinus pressure

■ nasal congestion ■ minor body aches and pains

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug; if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

■ you are taking a diuretic
■ you have a history of stomach problems, such as heartburn or fever reducers
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
■ have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ in children under 12 years of age
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug; if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

■ stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have a history of stomach problems, such as heartburn or you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke

When using this product

■ take with food or milk if stomach upset occurs
■ take any other drug
■ decrease this benefit of aspirin

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better

Ask a doctor or pharmacist before use if you are

■ taking any other product that contains pseudoephedrine or any other nasal decongestant
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

When using this product

■ take with food or milk if stomach upset occurs
■ take any other drug
■ decrease this benefit of aspirin

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better

Ask a doctor or pharmacist before use if you are

■ taking any other product that contains pseudoephedrine or any other nasal decongestant
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

Drug Facts (continued)

Other information

■ store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
■ read all warnings and directions before use. Keep carton.

■ children under 12 years of age: do not use by a doctor
■ do not use more than 6 caplets in any 24-hour period unless directed

Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years of age and over:
■ take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
■ do not use more than 6 caplets in any 24-hour period unless directed by a doctor

Drug Facts (continued)

Drug Facts (continued)

Inactive ingredients acetylated monoglycerides, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical grade, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxides, titanium dioxide

Questions or comments?
call toll free 1-800-88-ADVIL

UNPRINTED AREA FOR LOT & EXP CODING (UNVARNISHED)



Distributed by: GSK Consumer Healthcare,
Warren, NJ 07059
For most recent product information,
visit www.advil.com
Trademarks owned or licensed by GSK
©2021 GSK or licensee



Product inside sealed in plastic blister with foil backing. Do Not Use if plastic blister or foil barrier is broken.



 3 0573 0180 214

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
11/19/2021 03:10:02 PM