

Important: Read all product information before using. Keep this card for important information.

Drug Facts (continued)
 Keep out of reach of children. In case of overdose, call your healthcare provider or poison control center. **1-800-438-3273**. If you have had an allergic reaction to any medicine, tell your healthcare provider.

Directions
 Adults and children 12 years of age and older: Take 1 to 2 tablets once daily, depending on the severity of your symptoms. Adults 65 and older: Take only one tablet per day.

Warnings
 Tell your healthcare provider if you are taking any other medicines, especially other allergy medicines, or if you are pregnant, planning to get pregnant, or breastfeeding. Tell your healthcare provider if you have ever had an allergic reaction to any medicine, especially other allergy medicines, or if you have ever had a severe allergic reaction to any medicine, especially other allergy medicines. Tell your healthcare provider if you have ever had a severe allergic reaction to any medicine, especially other allergy medicines. Tell your healthcare provider if you have ever had a severe allergic reaction to any medicine, especially other allergy medicines.

Other information
 Each tablet contains 5 mg of cetirizine HCl. Each tablet also contains 5 mg of antihistamine. Each tablet also contains 5 mg of antihistamine. Each tablet also contains 5 mg of antihistamine.

Questions?
 Call 1-800-438-3273. Visit www.zyrtec.com. © 2011 Johnson & Johnson Consumer Inc. All rights reserved.

Active ingredient made in Belgium.
 Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McKel Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 CUSL0120XX zyrtec.com

00000000





Take 1 to 2 tablets* once daily, depending on the severity of your symptoms. Adults 65 and older: Take only one tablet per day.

24 Hour relief of

- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose
- Sneezing

35 Tablets
5 mg each

Actual Size



Indoor & Outdoor Allergies
 Strength 5 mg Tablets

ZYRTEC
 Cetirizine HCl tablets ALLERGY
 5 mg antihistamine

ZYRTEC
 ALLERGY
 Cetirizine HCl tablets
 5 mg antihistamine


The trade dress of this ZYRTEC® package is subject to trademark protection. Active ingredient made in Belgium.

LOT:
 EXP:



NDC 0000-060-00

LOT: XXXXXXX
EXP: XXXXXXX



ZYRTEC ALLERGY
Cetirizine HCl
antihistamine
5mg tablets

35
Tablets
5mg each

<p>Active ingredient (in each tablet) Cetirizine HCl 5 mg Uses: temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: a runny nose, sneezing, itchy, watery eyes, itching of the nose or throat.</p> <p>Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor if phenothiazin before use if you are taking tranquilizers or sedatives. When using this product, drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away if pregnant or breast-feeding; if breast-feeding, not recommended if pregnant; ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p>	<p>Purpose Antihistamine</p> <p>Directions Adults and children 6 years and over: 1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours. Adults 65 years and over: 1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor. Other information: Store between 20° to 25°C (68° to 77°F) in original container. Do not use after expiration date. Contains inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.</p> <p>Questions? call 1-800-543-7805 (toll-free) or 215-278-8755 (collect).</p> <p>The trade dress of this ZYRTEC® package is subject to trademark protection. Active ingredient made in Belgium. Distributed by: JOHNSON & JOHNSON CONSUMER INC. McKee Consumer Healthcare Division Fort Washington, PA 19034 USA CJA/JCI 200X zyrtec.com</p>
---	---

NDC 00000-000-00

GO PACKS
15 Portable Pouches
One 5 mg tablet per pouch





ZYRTEC
Cetirizine HCl tablets **ALLERGY**
5 mg antihistamine

Indoor & Outdoor Allergies
5 mg Strength Tablets

24
Hour relief of

- Sneezing
- Runny Nose
- Itchy Throat or Nose
- Itchy, Watery Eyes

15
Tablets
5 mg each

Actual Size



15
Tablets
5 mg each

Take 1 to 2 tablets* once daily depending on the severity of your symptoms
*Adults 65 and older take only one tablet per day



000000000



000000000



000000000

Active ingredient made in Belgium
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
Medical Consumer Healthcare Division
Fort Washington, PA 19034 USA
©16JUL 20XX
www.zyrtec.com
Pat. www.jjipats.com
The back flap of the Zyrtec® package is subject to trademark copyright.



PAPER PLASTIC BOX TUBE

Drug Facts

Active ingredient (in each tablet) Cetirizine HCl 5 mg

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before you use if you have liver or kidney disease. Your doctor or should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product:

- do not drink alcohol
- do not drive or operate machinery
- do not drink grapefruit juice
- do not take other sedatives, tranquilizers, or other drugs that may increase drowsiness

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away if you experience any of the following:

- hives or rash
- swelling of the face, lips, or throat
- difficulty breathing
- if it is used for relief of colds, sinusitis, or other respiratory conditions, symptoms may worsen

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-2222)

Directions

adults and children 6 years and over: 1 tablet (5 mg) or 2 tablets (10 mg) once daily

children 6 to 5 years: 1 tablet (5 mg) once daily, do not take more than 1 tablet (5 mg) in 24 hours.

ask a doctor if you are taking other antihistamines.

Other information

- Keep the container tightly closed.
- Do not use if the seal over the top of the container is broken or missing.
- Do not use if the seal over the top of the container is broken or missing.

Inactive ingredients cetirizine hydrochloride, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions? call 1-800-443-7805 (toll-free) or 215-273-8756 (local)



ZYRTEC
Cetirizine HCl tablets **ALLERGY**
5 mg antihistamine



Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product, drowsiness may occur; avoid alcoholic drinks, alcohol, caffeine, and tranquilizers; may increase drowsiness while car driving during a motor vehicle or operating machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast-feeding, or if breast-feeding, not recommended, or if pregnant, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-235-4273). **Directions:** Adults and children 6 years and over: 1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours. Adults 65 years and over: 1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours. Children under 6 years of age: ask a doctor. **Contents:** 10 tablets. **Other information:** Store at controlled room temperature (20° to 25°C (68° to 77°F)); do not use if pouch is torn or damaged. **Contains ingredients:** cetirizine dihydrochloride, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7015 (toll-free) or 215-273-8755 (in PA). Active ingredient made in Belgium. **Distributed by:** JANSSEN & JANSSEN CONSUMER INC., Malvern, Pennsylvania 19354, Fort Washington, PA 19034 USA. © J&J 101100X www.zyrtec.com Pat. www.jjpat.com 00000000