

Labeling Format Information: Fonts: HelveticaNeueLTStd: Cond, Bold Cond, Cond Bold Obl			
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0

Cetirizine
 Cetirizine
 Cetirizine
 Cetirizine



3
Tablets 0.125"

LOT:
 EXP:



00000000



Drug Facts
 Cetirizine HCl 10 mg
 Antihistamine

Active ingredient (in each tablet)
 Cetirizine HCl 10 mg

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
 Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
 Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product
 ■ avoid alcoholic drinks
 ■ drowsiness may occur
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
 ■ If pregnant or breast-feeding:
 ■ if pregnant, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 adults and children 6 years and over: take one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
 children 6 years and over: ask a doctor
 children under 6 years of age: ask a doctor
 consumers with liver or kidney disease: ask a doctor

Other information
 ■ store between 20° to 25°C (68° to 77°F)
 ■ do not use if pouch is torn or damaged
 ■ meets USP Dissolution Test 2

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions? call 1-800-343-7806 (toll-free) or 215-272-8756 (collect)

000000000000000000000000



00000000

Indoor & Outdoor Allergies
10 mg

Actual Size



3
Tablets
10 mg
each

Relief: Watery Eyes



Relief: Throat or Nose



24 Hour relief of



Sneezing



Relief: Runny Nose



Actual Size



ZYRTEC
 Cetirizine HCl tablets
10mg/antihistamine

NDC 0000-000-00

Original Prescription Strength

GO PACKS
 3 Portable Pouches
 One 10mg tablet per pouch

New look
 Same relief



Labeling Format Information: Fonts: HelveticaNeueLTStd: Cond, Bold Cond, Cond Bold Obl			
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0

Cetirizine
 Cetirizine
 Cetirizine
 Cetirizine



5
Tablets 0.125"

LOT:
 EXP:



Drug Facts

Active ingredient (in each tablet)
 Cetirizine HCl 10 mg Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
 Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
 Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product
 ■ drowsiness may occur ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
 ■ If pregnant or breast-feeding:
 ■ if pregnant, ask a health professional before use.
 ■ if breast-feeding, not recommended
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 adults and children 6 years and over **one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.**
 children 6 years and over **ask a doctor**
 children under 6 years of age **ask a doctor**
 consumers with liver or kidney disease **ask a doctor**

Other information
 ■ store between 20° to 25°C (68° to 77°F)
 ■ do not use if pouch is torn or damaged
 ■ meets USP Dissolution Test 2

Inactive ingredients colloid silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions? call 1-800-343-7865 (toll-free) or 215-272-8756 (collect)



0 000000 000000 0

000000000

Active ingredient made in Switzerland
 Distributed by:
JANSSEN & JANSSEN CONSUMER INC.
 1000 Locust Hill Road
 Spring House, PA 15141
 For Washington, PA 15054 USA
 © J&J 2018
www.zyrtec.com
 The Zyrtec logo and the ZYRTEC® package is a registered trademark of Janssen.



 Itchy, Watery Eyes
 Itchy Throat or Nose
 Sneezing
 Runny Nose
 Bumpy Nose

 **24** Hour relief of
 **5** Tablets
10 mg each

 Actual Size
 **5** Tablets
10 mg each



000000000



Original Prescription Strength

Indoor & Outdoor Allergies

10 mg



000000000



Indoor & Outdoor Allergies

10mg Strength

GO PACKS

5 Portable Pouches

One 10mg tablet per pouch

New look Same relief

Labeling Format Information: Fonts: Univers LT Pro Cd, Univers LT Pro Bd Cd, Univers LT Std Bd Cd Ob			
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 75%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 1.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: -.25

Cetirizine HCl tablets
Cetirizine HCl tablets
Cetirizine HCl tablets
Cetirizine HCl tablets

ZYRTEC

40 Tablets 0.125"



**BONUS 33% MORE
10 FREE TABLETS**

24 Hour relief of	

40 Tablets 10 mg each

30 + 10 = 40 Tablets 10 mg each

Active ingredient made in Switzerland.
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 20XX zyrtec.com



00000000


Drug Facts
Keep one of these at all times in case of overdose, get medical help or contact a Poison Control Center right away. (1-800-232-2222)

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing ipratropium.

Directions
Take one 10 mg tablet once daily, do not take more than one 10 mg tablet in 24 hours.

Other important information
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing ipratropium.

Questions? Call 1-800-343-7885 (toll-free) or 215-273-8755 (local).



Labeling Format Information: Form: Unvers LT Std Bold Cond Old, Unvers LT Pro: Cond, Bold Cond					
Drug Facts: 10.00 pt	Body Text: 8.50 pt	Bullets: 8.50 pt	Horizontal Scale: 88%		
Headings: 8.50 pt	Leadings: 7.00 pt	Bottoms: 2.00 pt			
Subheaders: 8.50 pt	Drug Facts (continued): 8.50 pt	Headlines: 0.50 pt	Average Tracking: 0		

Cetirizine
 Cetirizine
 Cetirizine
ZYRTEC

ZYRTEC
 Cetirizine HCl tablets ALLERGY
 10 mg /antihistamine

Indoor & Outdoor Allergies

Important: Read all product information before using. Keep this card for important information.

Drug Facts (continued)

Warnings: Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or tranquilizers. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you have ever had a seizure. Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or tranquilizers. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you have ever had a seizure.

Directions: Take 10 mg (one tablet) once daily with or without food. Do not take more than 10 mg in 24 hours. Do not take more than 10 mg in 24 hours. Do not take more than 10 mg in 24 hours.

Other information: Each bottle contains 70 or 50 tablets. Each bottle contains 70 or 50 tablets. Each bottle contains 70 or 50 tablets.

Keep out of reach of children. See important information about prescription drugs on page 223.

Original Prescription Strength NDC 0000-050-00

ZYRTEC
 Cetirizine HCl tablets ALLERGY
 10 mg /antihistamine

New look Same relief

Indoor & Outdoor Allergies
 Strength Tablets
10 mg

24 Hour relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

120 Tablets
 10 mg each

Actual Size

70 10 mg Tablets

50 10 mg Tablets

120 Total Tablets

70ct + 50ct = 120ct Tablets 10 mg each

Active ingredient made in Switzerland.
 Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 ©J&JCI 20XX
zyrtec.com
 00000000

Actual Size

70 10 mg Tablets

50 10 mg Tablets

Actual Size

The trade dress of this ZYRTEC package is subject to trademark protection. Active ingredient made in Switzerland.

LOT: _____
 EXP: _____

Labeling Format Information: Fonts: HelveticaNeueLTStd Bold Cond, Black Cond, Black Cond Obl			
Drug Facts: 5.00 pt	Body Text: 5.00 pt	Bullets: 4.00 pt	Horizontal Scale: 75%
Headers: 5.00 pt	Leading: 5.00 pt	Barlines: N/A pt	
Subheaders: 5.00 pt	Drug Facts (continued): N/A pt	Hairlines: N/A pt	Average Tracking: -20



Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product, drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. They use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, or if breast-feeding, not recommended. If pregnant, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Directions: Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information: Store between 20° to 25°C (68° to 77°F) in dry place. Do not use if pouch is torn or damaged, or contents are not in original container. Contains 2 tablets. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7809 (ask-3ee) or 215-273-6755 (ask-3ee). Active ingredient made in Switzerland.

This label may not be the latest approved by FDA.
 For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Labeling Format Information: Fonts: Unvers LT Pro Cd, Unvers LT Pro Bd Cd, Unvers LT Std Bd Cd Obl

Drug Facts: N/A pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%
Headers: 4.50 pt	Leading: 4.80 pt	Barlines: N/A pt	
Subheaders: 4.50 pt	Drug Facts (continued): N/A pt	Hairlines: N/A pt	Average Tracking: -20

Active ingredient for each tablet: Cetirizine HCl 10 mg. **Purpose:** Antihistamine

Use: to temporarily relieve these symptoms due to hay fever or other upper respiratory allergies:

- runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings: Do not use if you have ever had a large reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product:

- drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast feeding; ■ if breastfeeding: not recommended ■ if pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



Directions: Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information: Store between 20° to 25°C (68° to 77°F) ■ do not use if foil inner seal imprinted with "ZYRTEC" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Caution:** call 1-800-343-7885 (to-tell) or 215-373-8725 (collect). The trade dress of this ZYRTEC® package is subject to trademark protect. inc.

Active ingredient made in Switzerland.
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Health Care Division
 Fort Washington, PA 19034 USA ©JUL/JCI 20XX

zyrtec.com

NDC 00005-000-00

ZYRTEC®

Cetirizine HCl / ALLERGY
antihistamine
10 mg tablets

30

Tablets
10 mg each

LOT: XXXXXX

EXP: XXXXXX

This label may not be the latest approved by FDA.
 For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Labeling Format Information: Fonts: Unvers LT Pro Cd, Unvers LT Pro Bd Cd, Unvers LT Std Bd Cd Obl

Drug Facts: N/A pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%
Headers: 4.50 pt	Leading: 4.80 pt	Barlines: N/A pt	
Subheaders: 4.50 pt	Drug Facts (continued): N/A pt	Hairlines: N/A pt	Average Tracking: -20

Active ingredient for each tablet: Cetirizine HCl 10 mg.

Other ingredients: croscarmellose sodium, hydroxypropyl methylcellulose, polyethylene glycol, croscarmellose sodium, hydroxypropyl methylcellulose, polyethylene glycol, titanium dioxide, **Opadone®** color 1-800-343-7885 (in-USA) or 215-573-8725 (outside-USA).

Directions: Adults and children 6 years and over: one 10 mg tablet once daily, do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information: Store between 20° to 25°C (68° to 77°F). Do not use if foil inner seal is impaired with "ZYRTEC" is broken or missing or inner USP Dissolve Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Opadone®** color 1-800-343-7885 (in-USA) or 215-573-8725 (outside-USA). The trade dress of this ZYRTEC® package is subject to trademark protection.

Warnings: Do not use if you have ever had a large reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast feeding. If breastfeeding: not recommended. If pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Purpose: Antihistamine. Use to temporarily relieve these symptoms due to hay fever or other upper respiratory allergies: runny nose, sneezing, itchy, watery eyes, itching of the nose or throat.

Other information: Store between 20° to 25°C (68° to 77°F). Do not use if foil inner seal is impaired with "ZYRTEC" is broken or missing or inner USP Dissolve Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Opadone®** color 1-800-343-7885 (in-USA) or 215-573-8725 (outside-USA). The trade dress of this ZYRTEC® package is subject to trademark protection.

Warnings: Do not use if you have ever had a large reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast feeding. If breastfeeding: not recommended. If pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



00000007

NDC 00000-000-00

ZYRTEC®

Cetirizine HCl **ALLERGY**

antihistamine

10 mg tablets

40 Tablets
10 mg each

zyrtec.com

LOT: XXXXXX
EXP: XXXXXX

This label may not be the latest approved by FDA.
 For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Labeling Format Information: Fonts: Unvers LT Pro Cd, Unvers LT Pro Bd Cd, Unvers LT Std Bd Cd Obl

Drug Facts: N/A pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%
Headers: 4.50 pt	Leading: 4.80 pt	Barlines: N/A pt	
Subheaders: 4.50 pt	Drug Facts (continued): N/A pt	Hairlines: N/A pt	Average Tracking: -20

Active ingredient for each tablet: Cetirizine HCl 10 mg. **Purpose:** Antihistamine

Use: to temporarily relieve these symptoms due to hay fever or other upper respiratory allergies:

- runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings: Do not use if you have ever had a large reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product:

- drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast feeding; ■ if breastfeeding: not recommended ■ if pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions: Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information: Store between 20° to 25°C (68° to 77°F) ■ do not use if foil inner seal is impaired with "ZYRTEC" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7885 (1-800-343-7885) or 215-373-8725 (toll-free). The trade dress of this ZYRTEC® package is subject to trademark protect. inc.

Active ingredient made in Switzerland.
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Health Care Division
 Fort Washington, PA 19004 USA ©JUL/JCI 200X

zyrtec.com



ZYRTEC
 Cetirizine HCl **ALLERGY**
 antihistamine
10 mg tablets

50
 Tablets
 10 mg each

LOT: XXXXXX
 EXP: XXXXXX

This label may not be the latest approved by FDA.
 For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Labeling Format Information: Fonts: Unvers LT Pro Cd, Unvers LT Pro Bd Cd, Unvers LT Std Bd Cd Obl

Drug Facts: N/A pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%
Headers: 4.50 pt	Leading: 4.80 pt	Barlines: N/A pt	
Subheaders: 4.50 pt	Drug Facts (continued): N/A pt	Hairlines: N/A pt	Average Tracking: -20

Active ingredient for each tablet: Cetirizine HCl 10 mg.

Other ingredients: croscarmellose sodium, hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyethylene glycol, polyethylene glycol, titanium dioxide, **ALLERGY** (see Warnings).

Directions: Adults and children 6 years and over: one 10 mg tablet once daily, do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information: Store between 20° to 25°C (68° to 77°F). Do not use if foil inner seal is impaired with "ZYRTEC" is broken or missing or inner USP Dissolve Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyethylene glycol, titanium dioxide. **Warnings:** Do not use if you have ever had a large reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast feeding. If breastfeeding: not recommended. If pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Other information: Store between 20° to 25°C (68° to 77°F). Do not use if foil inner seal is impaired with "ZYRTEC" is broken or missing or inner USP Dissolve Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyethylene glycol, titanium dioxide. **Warnings:** Do not use if you have ever had a large reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast feeding. If breastfeeding: not recommended. If pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



NDC 00000-000-00

ZYRTEC

Cetirizine HCl **ALLERGY**
antihistamine
10 mg tablets

70
Tablets
10 mg each

LOT: XXXXXX
EXP: XXXXXX

zyrtec.com

This label may not be the latest approved by FDA.
 For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Labeling Format Information: Fonts: Unvers LT Pro Cd, Unvers LT Pro Bd Cd, Unvers LT Std Bd Cd Obl

Drug Facts: N/A pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%
Headers: 4.50 pt	Leading: 4.80 pt	Barlines: N/A pt	
Subheaders: 4.50 pt	Drug Facts (continued): N/A pt	Hairlines: N/A pt	Average Tracking: -20

Active ingredient for each tablet: Cetirizine HCl 10 mg.

Purpose: Antihistamine

Use: to temporarily relieve these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings: Do not use if you have ever had a large reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tricyclics or sedatives. When using this product:

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast feeding; ■ if breastfeeding: not recommended


- if pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions: Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information: Store between 20° to 25°C (68° to 77°F). Do not use if foil inner seal imprinted with "ZYRTEC" is broken or missing ■ means USP Dissolve in Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7885 (in NJ or 215-373-8725 (collect)). The trade dress of this ZYRTEC® package is subject to trademark protect. inc.

Active ingredient made in Switzerland.
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Health Care Division
 Fort Washington, PA 19004 USA ©JUL/JCI 200X

zyrtec.com



ZYRTEC
 Cetirizine HCl **ALLERGY**
 antihistamine
10 mg tablets

90
 Tablets
 10 mg each

LOT: XXXXXX
 EXP: XXXXXX

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
12/01/2023 03:58:50 PM