



**Active ingredient (in each tablet)**  
Cetirizine HCl 5mg.

**Purpose**  
Antihistamine

**Directions: Adults and children 6 years and over:** 1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10mg) in 24 hours. **Adults 65 years and over:** 1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information:** ■ store between 20° to 25° (68° to 77°) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7005 (toll-free) or 215-273-8755 (collect). The trade dress of this Zyrtec® package is subject to trademark protection. Active ingredient made in Switzerland.

**Warnings Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use** if you have liver or kidney disease. Your doctor should determine if you need a different dose. **Ask a doctor or pharmacist before use** if you are taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery. **Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away if **pregnant or breast-feeding.** ■ if breast-feeding, not recommended. ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Pat. www.jcigpats.com**  
Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA ©J&JCI 2023 **zyrtec.com**



# GO PACKS

15 Portable Pouches  
One 5 mg tablet per pouch





**ZYRTEC**  
Cetirizine HCl Tablets ALLERGY  
5 mg/antihistamine

**24**  
Hour relief of

- Sneezing
- Runny Nose
- Itchy Throat or Nose
- Itchy, Watery Eyes

**15**  
Tablets  
5 mg each

Actual Size



Take 1 to 2 tablets\* once daily depending on the severity of your symptoms  
\*Adults 65 and older take only one tablet per day

**15**  
Tablets  
5 mg each



XXXXXXXXXXXXXXXXXXXX



XXXXXXXXXXXXXXXXXXXX



3 00450 25615 7

Active ingredient made in Switzerland

Manufactured by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
Major Consumer Healthcare Division  
1st Floor, Innovation, 2910 Central Expressway  
Kenilworth, NJ 07033 USA  
© J&J, 2023  
[www.zyrtec.com](http://www.zyrtec.com)  
The Zyrtec packaging is made from 100% recycled materials to protect the planet.  
For more information visit [www.zyrtec.com](http://www.zyrtec.com)  
20230301





**Drug Facts**

**Active ingredient (in each tablet)** Cetirizine HCl 5 mg  
Antihistamine

**Uses**  
Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
runny nose ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.  
**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.  
**Ask a doctor or pharmacist before use if you are taking** tranquilizers or sedatives.

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away if you experience:  
■ drowsiness may occur ■ avoid alcoholic drinks  
■ alcohol, sedatives, and tranquilizers may increase drowsiness  
■ be careful when driving a motor vehicle or operating machinery

**Keep out of reach of children.** In case of overdose, get medical help or contact Poison Control Center right away: (1-800-222-1222)

**Directions**  
adults and children 6 years and over ■ 1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10mg) in 24 hours  
adults 65 years and over ■ 1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours  
children under 6 years of age ■ ask a doctor  
consumers with liver or kidney disease ■ ask a doctor

**Other information**  
■ store between 20° to 25° (68° to 77°)  
■ do not use if cap is opened, do not use if pouch is torn or damaged  
■ meets USP Dissolution Test 2

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

**Questions?** call 1-800-343-7895 (toll-free) or 215-273-8755 (collect)

LOT: XXXXXXXX  
EXP: YY/YY/MM

**Net Content Statement** 11.85 in<sup>2</sup>  
PDP Area: (width x length) 0.125"

**NET CONTENT**  
1.8 IN. MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.  
3/16 IN. MORE THAN 25 SQ. IN. BUT NOT MORE THAN 50 SQ. IN.  
1/4 IN. MORE THAN 50 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.  
1/2 IN. OVER 400 SQ. IN.

**Labeling Format Information:** Front: Helvetica® Cond Bold, Cond Bold Italic

Drug Facts: 9.00 pt | Body Text: 6.00 pt | Bullets: 5.00 pt | Horizontal Scale: 100%  
 Headers: 8.00 pt | Leading: 6.50 pt | Barlines: 1.50 pt  
 Subheaders: 6.00 pt | Drug Facts (continued): 6.00 pt | Hairlines: 0.50 pt | Average Tracking: 0

(b) (4)



-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

MARTHA K LENHART  
03/28/2024 07:57:14 PM