

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	20.58 in ²	1.16 BL	3 SQ. IN. OR LESS
Net Content Size:	0.125"	1.35 BL	MORE THAN 3 SQ. IN. BUT NOT MORE THAN 35 SQ. IN.
		3.16 BL	MORE THAN 35 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1.14 BL	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1.02 BL	OVER 400 SQ. IN.

Drug Facts Format Information							
Drug Facts:	8.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.0 pt	Horizontal Scale:	75%
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt	Tracking:	-25
Headings:	8.00 pt	Leading:	6.50 pt				

The image displays a detailed layout for the Zyrtec Hives Relief 10 mg antihistamine packaging. The design is primarily green and white, with red accents for the brand name and key benefits.

Front Panel: Features the Zyrtec Hives Relief logo at the top, followed by "Original Prescription Strength" and "Zyrtec Cetirizine HCl tablets HIVES 10 mg antihistamine". A central graphic highlights "HIVES RELIEF" with icons for "24 Hour relief", "30 Tablets", and "Reduces Itching Due to Hives". A small image of the product bottle is shown on the right. The bottom right corner includes a barcode and the text "LOT:XXXXXXXX EXP:YYYY/MM/YY".

Back Panel: Contains detailed regulatory information, including "Drug Facts", "Warnings", "Directions", and "Other information". The text is oriented vertically and is smaller than the front panel text.

Side Panel: Shows the "Actual Size" of the blister pack, which is a rectangular box containing 30 tablets.

Bottom Panel: Features a large, stylized graphic of a person's head and neck, with a green background and white lines indicating the shape of the head and neck.

Drug Facts Format Information			
Drug Facts:	N/A pt	Subheadings/Body Text:	4.50 pt
Drug Facts (continued):	N/A pt	Bullets:	3.50 pt
Headings:	4.50 pt	Leading:	4.70 pt
		Barlines:	N/A pt
		Hairlines:	N/A pt
		Horizontal Scale:	80%
		Tracking:	0

This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine**. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use** to prevent hives from an unknown cause such as: **● food** **● insect stings** **● medicines** **● latex** or rubber gloves because the product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. **● If you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have** **● latex** or hives disease. Your doctor should determine if you need a different dose. **● Hives** that are an unusual color, look bruised or blistered **● hives** that do not itch **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product** **● drowsiness** may occur **● avoid alcoholic drinks** **● alcohol**, sedatives, and tranquilizers may increase drowsiness **● be careful** when driving a motor vehicle or operating machinery **Stop use and ask a doctor if** **● an allergic reaction to this product occurs**. Seek medical help right away **● symptoms do not improve after 2 days of treatment** **● the hives have lasted more than 6 weeks** **● if pregnant or breast-feeding** **● if breast-feeding**, not recommended **● if pregnant**, ask a health professional before use. **Keep out of reach of children**. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



Active ingredient (in each tablet) **● Purpose**
 Cetirizine HCl 10mg. Antihistamine.
Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.
Warnings **● Severe Allergy** **Warning**: Get emergency help **immediately** if you have hives along with any of the following symptoms:
 ● trouble swallowing **● dizziness** or loss of consciousness **● swelling of tongue** **● swelling in or around mouth** **● trouble speaking** **● drooling**
 ● wheezing or problems breathing These symptoms may be signs of anaphylactic shock.
● PEEL HERE



Directions **Adults and children 6 years and over**: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. **Adults 65 years and over**: ask a doctor. **Children under 6 years of age**: ask a doctor. **Consumers with liver or kidney disease**: ask a doctor.
Other information **● Store** between 20° to 25°C (68° to 77°F) **● Do not use if foil inner seal imprinted with "Sealed For Your Safety"** **● In hives or itching** **● more** USP-Disinfection Test 2.
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.
Questions? call 1-800-243-7885 (ask for) or 215-272-4755 (collect).
 The trade dress of this ZYRTEC® package is subject to trademark protection.
 Active ingredient made in Switzerland.
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 © J&J, 2024 Pat. www.kanvaqpat.com **zyrtec.com**

01659006

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	9.69 in ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.125"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.30 pt	Horizontal Scale:	80%
				Tracking:	-10

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Cetirizine HCl 5 mg	Antihistamine

Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
 ■ trouble swallowing
 ■ dizziness or loss of consciousness
 ■ swelling of tongue
 ■ trouble speaking
 ■ wheezing or problems breathing
 ■ drooling
 ■ swelling in or around mouth
 ■ wheezing or problems breathing
 ■ trouble speaking
 ■ drooling
 ■ swelling in or around mouth

Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use
 ■ to prevent hives from any known cause such as:
 ■ foods ■ insect stings ■ medicines
 ■ latex or rubber gloves
 because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have
 ■ liver or kidney disease. Your doctor should determine if you need a different dose.
 ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

Drug Facts (continued)

When using this product
 ■ drowsiness may occur ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
 ■ an allergic reaction to this product occurs. Seek medical help right away.
 ■ symptoms do not improve after 3 days of treatment
 ■ the hives have lasted more than 6 weeks

If pregnant or breast-feeding:
 ■ if breast-feeding: not recommended
 ■ if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 ■ use only with enclosed dosing cup
 ■ find right dose on chart below ■ mL = milliliter

adults and children 6 years and over	5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours.
adults 65 years and over	5 mL once daily; do not take more than 5 mL in 24 hours.
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information
 ■ store between 20° to 25°C (68° to 77°F)
 ■ do not use if carton is opened or if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

Inactive ingredients anhydrous citric acid, flavors, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect).

The trade dress of this ZYRTEC® package is subject to trademark protection.
 Active ingredient made in Switzerland
 Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2024
 zyrtec.com Pat. www.kenvuepats.com 30056395

Children's ZYRTEC®
 Cetirizine HCl oral solution **HIVES**
 1mg/mL/antihistamine

OPEN HERE

Grape Flavor

NDC 50580-128-04



Hives Relief
6 yrs & older

HIVES RELIEF

24 Hour relief



Reduces Itching Due to Hives



Reduces Hives



Grape Flavor

4 fl oz (118 ml)

Dosing cup included

LOT XXXXXXX
 EXP YYYY/MMM



3 00450 13904 7

Drug Facts Format Information

Drug Facts:	N/A pt	Subheadings/Body Text:	5.00 pt	Barlines:	N/A pt	Horizontal Scale:	90%
Drug Facts (continued):	N/A pt	Bullets:	4.00 pt	Hairlines:	N/A pt	Tracking:	-20
Headings:	5.00 pt	Leading:	5.25 pt				

Active ingredient (in each 5 mL)
Cetirizine HCl 5 mg


Purpose
Antihistamine

Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings
Severe Allergy Warning: Get emergency help **immediately** if you have hives along with any of the following symptoms: trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ drooling ■ wheezing or problems breathing. These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use** ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may

LOT: XXXXXXXX
EXP: YYYY/MM

NDC 50580-126 04



Children's ZYRTEC[®]
Cetirizine HCl 1 mg/mL HIVES
oral solution antihistamine

6 yrs & older

HIVES RELIEF

24 Hour relief
Reduces Itching Due to Hives
Reduces Hives

Grape Flavor 4 fl oz (118 mL)

be able to help you find a cause. ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have** ■ liver or kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch. **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness; ■ be careful when driving a motor vehicle or operating machinery **Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks. **If pregnant or breast-feeding:** ■ if breast-feeding, not recommended ■ if pregnant, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Directions ■ use only with enclosed dosing cup ■ find right dose on chart below ■ mL = milliliters **Adults and children 6 years and over:** 5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours. **Adults 65 years and over:** 5 mL once daily; do not take more than 5 mL in 24 hours. **Children under 6 years of age:** ask a doctor **Consumers with liver or kidney disease:** ask a doctor **Other information** ■ store between 20° to 25° C (68° to 77° F) ■ **do not use if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing.**

Inactive ingredients anhydrous citric acid, flavors, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose **Questions?** call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

Dosing cup should be washed and left to air dry after each use.

Active ingredient made in Switzerland
The trade dress of this ZYRTEC[®] package is subject to trademark protection.
Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA
©J&J 2024 zyrtec.com Pat. www.kenvuepats.com 30056339

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

MARTHA K LENHART
11/26/2024 03:45:08 PM