

(b) (4)



TYLENOL
ARTHRITIS PAIN
 Contains Acetaminophen Extended-Release Pain Reliever/Fever Reducer
 For The Temporary Relief Of Minor Arthritis Pain
Caplets
 150 CAPLETS 650 mg each
Push & Turn Cap

Contains No Aspirin

Drug Facts (cont. nued)

Directions
 ■ do not take more than directed (see overdose warning)

adults

- take 2 caplets every 8 hours with water
- swallow whole – do not crush, chew or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age

- ask a doctor

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat
- do not use if carton is opened or neck wrap or foil inner seal imprinted with "Safety Seal" is broken
- see end panel for lot number and expiration date

Inactive ingredients
 carnauba wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?
 call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)

What makes Tylenol® Arthritis Pain Extended-Release Caplets different?

- Uses a unique, patented bi-layer caplet. The first layer dissolves quickly to provide prompt relief, while the second layer is time released to provide up to 8 hours of relief.
- For more information or questions, visit our web site www.tylenol.com.
- The makers of Tylenol® do not manufacture store brands.

00000000

XXXXXXX

Drug Facts

Active ingredient (in each caplet)
 Acetaminophen 650 mg Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to
- arthritis
- the common cold
- headache
- toothache
- muscular aches
- backache
- menstrual cramps
- temporarily reduces fever

Warnings

Alcohol warning: if you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage.

Do not use

- if you are allergic to acetaminophen or any of the inactive ingredients in this product
- with any other product containing acetaminophen

Stop use and ask a doctor if

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.


00000000



TYLENOL
ARTHRITIS PAIN
 Contains Acetaminophen Extended-Release Pain Reliever/Fever Reducer
 For The Temporary Relief Of Minor Arthritis Pain
Caplets
 150 CAPLETS 650 mg each
Push & Turn Cap

00000000

00000000



(b) (4)

(b) (4)


(b) (4)

<p>Active ingredient (in each caplet) Purpose Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p>Uses ■ temporarily relieves minor aches and pains due to: ■ muscular aches ■ backache ■ headache ■ toothache ■ the common cold ■ menstrual cramps ■ minor pain of arthritis ■ temporarily reduces fever</p> <p>Warnings Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage. Do not use ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product ■ with any other product containing acetaminophen Stop use and ask a doctor if ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of</p>	<p>III READ THE LABEL III</p> <p>NDC 50580-297-10</p> <p>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</p> <p><i>*Take only as directed.</i></p> <p>TYLENOL®</p> <p>8 Hour</p> <p>Pain Reliever / Fever Reducer</p> <p>Contains <i>Acetaminophen Extended Release Pain Reliever</i></p> <p>For Up to 8 Hour Relief of Minor Muscular Aches & Pain</p> <p>100 CAPLETS 650 mg each</p>	<p>overdose, get medical help or contact a Poison Control Center right away. (1 800 222 1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions do not take more than directed (see overdose warning) Adults and children 12 years and over: ■ take 2 caplets every 8 hours with water ■ swallow whole – do not crush, chew or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor Children under 12 years: ■ do not use</p> <p>Other information ■ store at 20-25°C (68 – 77°F). Avoid excessive heat 40°C (104°F). ■ do not use if neck wrap or foil inner seal imprinted with "Safety Seal®" is broken</p> <p>Inactive ingredients com starch, D&C yellow #10 aluminum lake, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hydroxyethyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, sucralose, talc, titanium dioxide</p> <p>Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)</p>	<p>PREP BY: McNeil Consumer Healthcare, DIVISION OF MCNEIL-PPC, INC. FORT WASHINGTON, PA 19034 USA ©MCNEIL-PPC, INC. 2009 www.tylenol.com U.S. Pat. Nos. 4,368,509 and 5,004,613</p> <p>EXP. DATE: _____</p> <p>LOT: _____</p> <p>XXXXXXXX</p>
---	--	--	--

150% OF SIZE ^a




<p>Active ingredient (in each caplet) Purpose Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p>Uses ■ temporarily relieves minor aches and pains due to: ■ muscular aches ■ backache ■ headache ■ toothache ■ the common cold ■ menstrual cramps ■ minor pain of arthritis ■ temporarily reduces fever</p> <p>Warnings Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage. Do not use ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product ■ with any other product containing acetaminophen Stop use and ask a doctor if ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of</p>	<p>III READ THE LABEL III</p> <p>NDC 50580-297-10</p> <p>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</p> <p><i>*Take only as directed.</i></p> <p>TYLENOL®</p> <p>8 Hour</p> <p>Pain Reliever / Fever Reducer</p> <p>Contains <i>Acetaminophen Extended Release Pain Reliever</i></p> <p>For Up to 8 Hour Relief of Minor Muscular Aches & Pain</p> <p>100 CAPLETS 650 mg each</p>	<p>overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions do not take more than directed (see overdose warning) Adults and children 12 years and over: ■ take 2 caplets every 8 hours with water ■ swallow whole – do not crush, chew or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor Children under 12 years: ■ do not use</p> <p>Other information ■ store at 20-25°C (68 – 77°F). Avoid excessive heat 40°C (104°F). ■ do not use if neck wrap or foil inner seal imprinted with "Safety Seal®" is broken</p> <p>Inactive ingredients com starch, D&C yellow #10 aluminum lake, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hydroxyethyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, sucralose, talc, titanium dioxide</p> <p>Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)</p>	<p>PREP BY: McNeil Consumer Healthcare, DIVISION OF MCNEIL-PPC, INC. FORT WASHINGTON, PA 19034 USA ©MCNEIL-PPC, INC. 2009 www.tylenol.com U.S. Pat. Nos. 4,368,509 and 5,004,613</p> <p>EXP. DATE: _____</p> <p>LOT: _____</p> <p>XXXXXXXX</p>
---	--	--	--



LBL Ty 8 Hour (b) (4)
February 9, 2009

(b) (4)


(b) (4)

<p>Active ingredient (in each caplet) Purpose Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: ■ arthritis ■ the common cold ■ headache ■ toothache ■ muscular aches ■ backache ■ menstrual cramps ■ temporarily reduces fever <p>Warnings</p> <p>Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product ■ with any other product containing acetaminophen <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present <p>These could be signs of a serious condition.</p> <p>If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults</p>	<p>III READ THE LABEL III</p> 	<p>as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <p>do not take more than directed (see overdose warning)</p> <p>Adults:</p> <ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water ■ swallow whole do not crush, chew or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor <p>Under 18 years of age:</p> <ul style="list-style-type: none"> ■ ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F). ■ do not use if neck wrap or foil inner seal imprinted with "Safety Seal®" is broken <p>Inactive ingredients</p> <p>carnauba wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p>Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)</p> <p>Distributed by: McNeil Consumer Healthcare DIVISION OF MCNEIL-PPC, INC. FORT WASHINGTON, PA 19034 USA © MCN-PPC, INC. 2009 www.tylenol.com U.S. Patent Nos. 4,968,509 and 5,004,613</p>
--	--	--

120 % OF SIZE ^a



<p>Active ingredient (in each caplet) Purpose Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: ■ arthritis ■ the common cold ■ headache ■ toothache ■ muscular aches ■ backache ■ menstrual cramps ■ temporarily reduces fever <p>Warnings</p> <p>Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product ■ with any other product containing acetaminophen <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults</p>	<p>III READ THE LABEL III</p> 	<p>as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <p>do not take more than directed (see overdose warning)</p> <p>Adults:</p> <ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water ■ swallow whole – do not crush, chew or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor <p>Under 18 years of age:</p> <ul style="list-style-type: none"> ■ ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F). ■ do not use if neck wrap or foil inner seal imprinted with "Safety Seal®" is broken <p>Inactive ingredients</p> <p>carnauba wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p>Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)</p> <p>Distributed by: McNeil Consumer Healthcare DIVISION OF MCNEIL-PPC, INC. FORT WASHINGTON, PA 19034 USA © MCN-PPC, INC. 2009 www.tylenol.com U.S. Patent Nos. 4,968,509 and 5,004,613</p>
--	---	--



LBL Ty Arth Pain (b) (4)
 February 9, 2009

(b) (4)

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Joel Schiffenbauer
6/17/2009 07:05:30 AM