

(b) (4)

00000000



**Drug Facts**

**Active ingredient (in each caplet)** Acetaminophen 650 mg. **Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - influenza
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcohol drinks every day while using this product

**Do not use**

- if any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if:**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.



How can we help?  
**1-877-895-3665**

00000000

3 0045-0838-500 6



NDC 50580-XXX-XX

# TYLENOL®

**8HR ARTHRITIS PAIN**

**Acetaminophen** extended-release tablets

Pain Reliever / Fever Reducer

For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets

**50 Caplets\* 650 mg each**

**TYLENOL®**

**8HR ARTHRITIS PAIN**

Care To Recycle™

Distributed by:  
 McNeil Consumer Healthcare  
 Division of McNEIL-PPC, Inc.  
 Fort Washington, PA 19034 USA  
 © McNEIL-PPC, INC. 2013

**Contains No Aspirin**

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Other medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

do not take more than directed (see overdose warning)

**adults**

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

**Under 18 years of age**

- ask a doctor

**Other information**

- store at 20-25°C (68-77°F); avoid excessive heat 40°C (104°F)
- do not use if caplets are opened or neck wrap is dented
- inner seal imprinted with "TYLENOL®" is broken

**Inactive ingredients**

canalvaewax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?**  
 Call 1-877-895-3666 (68-HEE) or 215-273-8735 (toll-free)




00000000

<p><b>Active ingredient (in each caplet)</b>                  Acetaminophen 650 mg..... Pain reliever/fever reducer</p> <p><b>Purpose</b>                  Temporarily relieves minor aches and pains due to:</p> <ul style="list-style-type: none"> <li>■ minor pain of arthritis</li> <li>■ muscular aches</li> <li>■ backache</li> <li>■ premenstrual and menstrual cramps</li> <li>■ the common cold</li> <li>■ headache</li> <li>■ toothache</li> </ul> <p>Temporarily reduces fever</p> <p><b>Warnings</b>                  Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount. Do not use with other drugs containing acetaminophen. Do not use with 3 or more alcoholic drinks every day while using this product. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product, ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if you get a rash or have a more than 10 days use fever get two use or use a more than 3 days a new symptoms occur or redness or swelling is present. These could be signs of a serious condition.</p>	<p><b>READ THE LABEL</b></p> <p><b>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</b></p> <p>NDC 50580-XXX-XX</p> <p><b>TYLENOL</b>                  8HR ARTHRITIS PAIN</p> <p>Acetaminophen extended-release tablets                  Pain Reliever / Fever Reducer</p> <p>For The Temporary Relief Of Minor Arthritis Pain</p> <p>50 Caplets*                  650 mg each</p>	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b>                  do not take more than directed (see overdose warning)                  Adults: take 2 caplets every 8 hours with water                  ■ swallow whole; do not crush, chew, split or dissolve                  ■ do not take more than 6 caplets in 24 hours                  ■ do not use for more than 10 days unless directed by a doctor                  Under 18 years of age: ask a doctor</p> <p><b>Other information</b>                  ■ store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F). ■ do not use if neck wrap or foil inner seal imprinted with "TYLENOL" is broken</p> <p><b>Inactive ingredients:</b> carnauba wax, corn starch, hydroxyethyl cellulose, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polydioxanone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p>Questions or comments? call 1-877-495-3465 (ask-tes) or 215-272-0733 (o-let)</p>
---	--	--


(b) (4)



COVER, PAGE 1



BACK OF COVER, PAGE 2



BASE LABEL, PAGE 3



(b) (4)



3 0045-0838-377



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets

NDC 50580-XXX-XX



**Care To Recycle**<sup>™</sup>

Distributed by:  
McNeil Consumer Healthcare  
Division of McNEIL-PPC, Inc.  
Fort Washington, PA 19034 USA  
©McNEIL-PPC, INC. 2013

**Contains No Aspirin**



00000000





00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets







*How can we help?*  
**1-877-895-3665**









00000000





*How can we help?*  
**1-877-895-3665**



EXP



**TYLENOL**<sup>®</sup>  
**8HR ARTHRITIS PAIN**  
Acetaminophen extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets





<p><b>Active ingredient (in each caplet)</b>                  Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p><b>Purpose</b>                  Pain reliever/fever reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>■ temporarily relieves minor aches and pains due to:                         <ul style="list-style-type: none"> <li>■ minor pain of arthritis ■ muscular aches ■ backache</li> <li>■ premenstrual and menstrual cramps</li> <li>■ the common cold ■ headache ■ toothache</li> </ul> </li> <li>■ temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount. With other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product. Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p>	<p style="text-align: center;"><b>READ THE LABEL</b></p> <p style="text-align: center;"><b>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</b></p> <p style="text-align: center;">NDC 50580-XXX-XX</p> <p style="text-align: center;"><b>TYLENOL</b>  <b>8 HR ARTHRITIS PAIN</b>                  Acetaminophen extended-release tablets                  Pain Reliever / Fever Reducer</p> <p style="text-align: center;">For The Temporary Relief                  Of Minor Arthritis Pain</p>  <p style="text-align: center;">*Capsule-Shaped                  Bi-Layer Tablets</p> <p style="text-align: center;">225 Caplets*                  650 mg each</p>	<p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <p>do not take more than directed (see overdose warning)</p> <p><b>Adults:</b></p> <ul style="list-style-type: none"> <li>■ take 2 caplets every 8 hours with water</li> <li>■ swallow whole; do not crush, chew, split or dissolve</li> <li>■ do not take more than 6 caplets in 24 hours</li> <li>■ do not use for more than 10 days unless directed by a doctor</li> </ul> <p><b>Under 18 years of age:</b></p> <ul style="list-style-type: none"> <li>■ ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F).</li> <li>■ do not use if neck wrap or foil inner seal imprinted with "TYLENOL" is broken</li> </ul> <p><b>Inactive ingredients</b>                  camellia wax, croscellose, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p><b>Questions or comments?</b>                  call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p> <p>Distributed by:                  McNeil Consumer Healthcare                  Division of McNEIL-PPC, Inc.                  Fort Washington, PA 19034 USA                  ©McNEIL-PPC, INC. 2013</p>
---	---	--

(b) (4)



How can we help?  
**1-877-895-3665**



NDC 50580-XXX-XX

# TYLENOL®

## 8HR ARTHRITIS PAIN

### Acetaminophen extended-release tablets

Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain \*Capsule-Shaped Bi-Layer Tablets



**290 Caplets\*  
650 mg each**

## TYLENOL®

### 8HR ARTHRITIS PAIN

#### Acetaminophen

extended-release tablets  
Pain Reliever / Fever Reducer  
For The Temporary Relief Of Minor Arthritis Pain

**Contains No Aspirin**

**Directions**  
(see overdose warning)

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age

■ ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F). Avoid excessive heat 40°C (104°F).
- do not use if carton is opened or neck wrap or foil inner seal imprinted with "TYLENOL" is broken

**Inactive ingredients**

camelina wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?**  
call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

**Care To Recycle™**

Distributed by:  
McNeil Consumer Healthcare  
Division of McNEIL-PPC, Inc.  
Fort Washington, PA 19034 USA  
© McNEIL-PPC, INC., 2013



**Drug Facts (continued)**

**Directions**  
(see overdose warning)

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age

■ ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F). Avoid excessive heat 40°C (104°F).
- do not use if carton is opened or neck wrap or foil inner seal imprinted with "TYLENOL" is broken

**Inactive ingredients**

camelina wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?**  
call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

**Care To Recycle™**

Distributed by:  
McNeil Consumer Healthcare  
Division of McNEIL-PPC, Inc.  
Fort Washington, PA 19034 USA  
© McNEIL-PPC, INC., 2013



**Drug Facts**

**Active ingredient (in each caplet)** Purpose  
Acetaminophen 650 mg..... Pain reliever/fever reducer

**Uses**

- temporarily relieve minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



(b) (4)

<p><b>Active ingredient (in each caplet) Purpose</b>                  Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves minor aches and pains due to:                         <ul style="list-style-type: none"> <li>minor pain of arthritis</li> <li>muscular aches</li> <li>backache</li> <li>premenstrual and menstrual cramps</li> <li>the common cold</li> <li>headache</li> <li>toothache</li> </ul> </li> <li>temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> <li>more than 6 caplets in 24 hours, which is the maximum daily amount</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Do not use</b> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <ul style="list-style-type: none"> <li>if you are allergic to acetaminophen or any of the inactive ingredients in this product</li> </ul> <p><b>Ask a doctor before use if you have liver disease</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>pain gets worse or lasts more than 10 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>new symptoms occur</li> <li>redness or swelling is present</li> </ul> <p>These could be signs of a serious condition.</p>	<p style="text-align: center;"><b>READ THE LABEL</b></p> <p style="text-align: center;"><b>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</b></p> <p style="text-align: center;">NDC 50580-XXX-XX</p> <p style="text-align: center;"><b>TYLENOL</b>  <b>8 HR ARTHRITIS PAIN</b></p> <p style="text-align: center;"><b>Acetaminophen</b> extended-release tablets                  Pain Reliever / Fever Reducer</p> <p style="text-align: center;">For The Temporary Relief                  Of Minor Arthritis Pain</p> <p style="text-align: center;">*Capsule-Shaped                  Bi-Layer Tablets</p> <p style="text-align: center;"><b>290 Caplets*</b>  <b>650 mg each</b></p>	<p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed (see overdose warning)</li> </ul> <p><b>Adults:</b></p> <ul style="list-style-type: none"> <li>take 2 caplets every 8 hours with water</li> <li>swallow whole; do not crush, chew, split or dissolve</li> <li>do not take more than 6 caplets in 24 hours</li> <li>do not use for more than 10 days unless directed by a doctor</li> </ul> <p><b>Under 18 years of age:</b></p> <ul style="list-style-type: none"> <li>ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F).</li> <li>do not use if neck wrap or foil inner seal imprinted with "TYLENOL" is broken</li> </ul> <p><b>Inactive ingredients</b>                  camauba wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p><b>Questions or comments?</b>                  call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p> <p>Distributed by:                  McNeil Consumer Healthcare                  Division of McNEIL-PPC, Inc.                  Fort Washington, PA 19034 USA                  ©McNEIL-PPC, INC. 2013                  Visit us at <a href="http://www.tylenol.com">www.tylenol.com</a></p>
--	--	---

(b) (4)

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

THERESA M MICHELE  
12/20/2013