



Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic					
Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%		
Headers: 4.50 pt	Leading: 4.70 pt	Barlines: NA pt	Average Kerning: -25		
Subheaders: 4.50 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt			

Drug Facts

Active ingredient (in each caplet) Acetaminophen 650 mg. **Pain reliever/fever reducer**

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - headache
 - menstrual and menstrual cramps
 - headache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount.

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease or if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Drug Facts (continued)

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

<p>adults and children 12 years and over</p> <ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water ■ swallow whole, do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor 	<p>children under 12 years</p> <ul style="list-style-type: none"> ■ do not use
---	---

Other information

- store between 20-25°C (68-77°F)
- do not use if carton is opened or if lid inner seal imprinted with "SAFETY SEAL" is broken

Inactive ingredients: carmelum xanthan gum, hydroxyethylcellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, polydioxane, polydioxanone, polyethylene glycol, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?
call 1-877-895-3665 (toll-free) or 215-273-8959 (collect)

3 0045-0000-00 2



EXP: LOT:



HERITAGE OF CARE

How can we help?
1-877-895-3665



See New Warning

TYLENOL[®] 8HR

MUSCLE ACHES & PAIN

Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

**Capsule-Shaped 8H-Layer Tablets*



100 Caplets*
650 mg each





00000000


Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 80%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Kerning: 0

McNeil Submission Artwork

(b) (4)

Active ingredient (in each caplet) Acetaminophen 650 mg	Purpose Pain reliever/fever reducer	TO OPEN: PUSH AND TURN CAP READ THE LABEL	Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Uses Temporarily relieves minor aches and pains due to: ■ muscular aches ■ backache ■ minor pain of arthritis ■ toothache ■ premenstrual and menstrual cramps ■ headache ■ the common cold	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.		NDC 50580-309-02 DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
		TYLENOL[®] 8HR MUSCLE ACHES & PAIN Acetaminophen Extended-release tablets Pain Reliever / Fever Reducer For Up to 8 Hours Relief of Minor Muscle Aches & Pain *Caplet-Shaped Bi-Layer Tablets 100 Caplets* 650 mg each	Other information ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted with "SAFETY SEAL" is broken or missing Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect) Distributed by: McNeil Consumer Healthcare Division of McNEIL-PPC, Inc. Fort Washington, PA 19034 USA ©McNEIL-PPC, INC. 2014 00000000

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 75%	
Headers: 6.00 pt	Leading: 6.50 pt	Barlines: NA pt	Average Kerning: 0	
Subheaders: 6.00 pt	Drug Facts (continued): NA pt	Hairlines: NA pt		

TYLENOL® 8HR
MUSCLE ACHES & PAIN
Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

Care To Recycle®

Contains No Aspirin

TYLENOL® 8HR
MUSCLE ACHES & PAIN
Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

Care To Recycle®

Contains No Aspirin

TYLENOL® 8HR
MUSCLE ACHES & PAIN
Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

Care To Recycle®

Contains No Aspirin



How can we help?
1-877-895-3665

See New Warning

TYLENOL® 8HR
MUSCLE ACHES & PAIN
Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

*Capsule-Shaped Bi-Layer Tablets

225 Caplets*
650 mg each



Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses ■ Temporarily relieves minor aches and pains due to:
■ muscular aches ■ backache ■ minor pain of arthritis
■ toothache ■ menstrual and menstrual cramps
■ headache ■ the common cold
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 6 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Aller giv alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding drug, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed (see overdose warning)
adults and ■ take 2 caplets every 8 hours with water
children 12 ■ swallow whole; do not crush, chew, split
years and ■ or dissolve
over ■ do not take more than 6 caplets in 24 hours
■ do not use for more than 10 days
unless directed by a doctor

children under 12 years ■ do not use

Other information
■ store between 20-25°C (68-77°F)
■ do not use if caplet is opened or foil inner seal is imprinted with "SAFE SEAL" is broken

Inactive ingredients: camauaba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?
call 1-877-895-3665 (toll-free) or 215-273-8735 (collect)

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses ■ Temporarily relieves minor aches and pains due to:
■ muscular aches ■ backache ■ minor pain of arthritis
■ toothache ■ menstrual and menstrual cramps
■ headache ■ the common cold
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 6 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Aller giv alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses ■ Temporarily relieves minor aches and pains due to:
■ muscular aches ■ backache ■ minor pain of arthritis
■ toothache ■ menstrual and menstrual cramps
■ headache ■ the common cold
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 6 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Aller giv alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding drug, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed (see overdose warning)
adults and ■ take 2 caplets every 8 hours with water
children 12 ■ swallow whole; do not crush, chew, split
years and ■ or dissolve
over ■ do not take more than 6 caplets in 24 hours
■ do not use for more than 10 days
unless directed by a doctor

children under 12 years ■ do not use

Other information
■ store between 20-25°C (68-77°F)
■ do not use if caplet is opened or foil inner seal is imprinted with "SAFE SEAL" is broken

Inactive ingredients: camauaba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?
call 1-877-895-3665 (toll-free) or 215-273-8735 (collect)

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses ■ Temporarily relieves minor aches and pains due to:
■ muscular aches ■ backache ■ minor pain of arthritis
■ toothache ■ menstrual and menstrual cramps
■ headache ■ the common cold
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 6 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Aller giv alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	Average Kerning: 0
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	


McNeil

<p>Active ingredient (in each caplet) Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p>Purpose Pain reliever/fever reducer</p> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ muscular aches ■ backache ■ minor pain of arthritis ■ toothache ■ premenstrual and menstrual cramps ■ headache ■ the common cold ■ temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount. With other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product. Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, new symptoms occur, redness or swelling is present. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.</p>	<p>TO OPEN: PUSH AND TURN CAP READ THE LABEL</p> <p>NDC 50590-309-03</p> <p>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</p> <p>TYLENOL® 8HR MUSCLE ACHES & PAIN Acetaminophen Extended-release tablets Pain Reliever / Fever Reducer</p> <p>For Up to 8 Hours Relief of Minor Muscle Aches & Pain</p> <p>225 Caplets* 650 mg each</p> <p>*Capsule-Shaped Bi-Layer Tablets</p>	<p>Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <p>do not take more than directed (see overdose warning)</p> <p>Adults and children 12 years and over:</p> <ul style="list-style-type: none"> ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor <p>Children under 12 years:</p> <ul style="list-style-type: none"> ■ do not use <p>Other information</p> <ul style="list-style-type: none"> ■ store between 20-25°C (68-77°F) ■ do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing <p>Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p>Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p> <p>Distributed by: McNeil Consumer Healthcare Division of McNEIL-PPC, Inc. Fort Washington, PA 19034 USA ©McNEIL-PPC, INC. 2014</p>
--	--	--

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: NA pt	
Subheaders: 6.00 pt	Drug Facts (continued): NA pt	Hairlines: NA pt	Average Kerning: 0

McNeil Submission Artwork

(b) (4)



Labeling Format Information:			
Drug Facts:	NA pt	Body Text:	4.50 pt
Headers:	4.50 pt	Leading:	4.70 pt
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt
		Bullets:	3.50 pt
		Barlines:	NA pt
		Hairlines:	NA pt
		Horizontal Scale:	80%
		Average Kerning:	-25

McNeil Submission Artwork

(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY

12/23/2014