






Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic							
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	75%
Headers:	4.50 pt	Leading:	4.50 pt	Barlines:	NA pt	Average Kerning:	-25
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt		

**McNeil** Submission Artwork


(b) (4)



COVER, PAGE 1



BACK OF COVER, PAGE 2



BASE LABEL, PAGE 3

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: 10.00 pt	Body Text: 6.5-7.0pt	Bullets: 5.0-6.0pt	Horizontal Scale: 85%	
Headers: 8.0-9.0 pt	Leading: 6.5-7.5pt	Barlines: 2.50 pt	Average Kerning: -25	
Subheaders: 7.0 pt	Drug Facts (continued): 9.00pt	Hairlines: 0.50 pt		

**McNeil** Submission Artwork

**Drug Facts**  
**Active ingredient (in each caplet)** Acetaminophen 650 mg.....Pain reliever/fever reducer

**Purpose**

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - menstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy** alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away; (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age

- ask a doctor

**Other information**

- store between 20-25°C (68-77°F)
- do not use if carton is opened, or if foil inner seal imprinted with "SAFETYSEAL" is broken

**Inactive ingredients**

cantharidin wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?**  
call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

3 0045-0838-37 7



EXP.



How can we help?  
☎ 1-877-895-3665

NDC 50580-783-25

See New Warning

TYLENOL<sup>®</sup> 8HR

ARTHRITIS PAIN

Acetaminophen  
Extended-release tablets

Pain Reliever / Fever Reducer

For The Temporary Relief Of Minor Arthritis Pain



225 Caplets\*  
650 mg each

\*Capsule-Shaped Bi-Layer Tablets

00000000

**Labeling Format Information:** Fonts: HelveticaDF Medium, Bold, Bold Italic

Drug Facts: 10.00 pt	Body Text: 7.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%
Headers: 9.00 pt	Leading: 7.50 pt	Barlines: 2.50 pt	Average Kerning: -25
Subheaders: 7.00 pt	Drug Facts (continued): 9.00 pt	Hairlines: 0.50 pt	




<p>00000000</p>  <p>010101</p>	<p><b>Active ingredient (in each caplet)</b>                  Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p><b>Purpose</b>                  Pain reliever/fever reducer</p> <p><b>Uses</b>                  ■ temporarily relieves minor aches and pains due to:                  ■ minor pain of arthritis ■ muscular aches ■ backache                  ■ premenstrual and menstrual cramps                  ■ the common cold ■ headache ■ toothache                  ■ temporarily reduces fever</p> <p><b>Warnings</b>  <b>Liver warning</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount. With other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. <b>Allergy alert</b> acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. <b>Do not use</b> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. <b>Ask a doctor before use</b> if you have liver disease. <b>Ask a doctor or pharmacist before use</b> if you are taking the blood thinning drug warfarin. <b>Stop use and ask a doctor</b> if pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.</p>	<p><b>TO OPEN: PUSH AND TURN CAP</b>                  READ THE LABEL</p> <p><b>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</b></p> <p>NDC 50580-783-25</p> <p><b>TYLENOL 8HR</b>  <b>ARTHRITIS PAIN</b></p> <p><b>Acetaminophen Extended-release tablets</b>                  Pain Reliever / Fever Reducer</p> <p>For The Temporary Relief Of Minor Arthritis Pain</p> <p>*Capsule-Shaped Bi-Layer Tablets</p> 	<p>If pregnant or breast-feeding, ask a health professional before use. <b>Keep out of reach of children.</b>  <b>Overdose warning</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b>                  ■ do not take more than directed (see overdose warning)  <b>Adults</b> ■ take 2 caplets every 8 hours with water                  ■ swallow whole; do not crush, chew, split or dissolve                  ■ do not take more than 6 caplets in 24 hours                  ■ do not use for more than 10 days unless directed by a doctor  <b>Under 18 years of age</b> ■ ask a doctor</p> <p><b>Other information</b>                  ■ store between 20-25°C (68-77°F)                  ■ do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing</p> <p><b>Inactive ingredients</b> calcium nauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p><b>Questions or comments?</b> call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)                  Distributed by: McNeil Consumer Healthcare                  Division of McNEIL-PPC, Inc.                  Fort Washington, PA 19034 USA                  ©McNEIL-PPC, INC. 2016</p>
---	---	---	---

<b>Labeling Format Information:</b> Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%
Headers: 8.00 pt	Leading: 6.70 pt	Barlines: NA pt	
Subheaders: 6.00 pt	Drug Facts (continued): NA pt	Hairlines: NA pt	Average Kerning: -25

**McNeil** Submission Artwork

(b) (4)

00000000



**Drug Facts**  
**Active ingredient (in each caplet)** Purpose  
Acetaminophen 650 mg.....Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis  
 ■ muscular aches  
 ■ backache  
 ■ premenstrual and menstrual cramps  
 ■ the common cold  
 ■ headache  
 ■ toothache

**Warnings**  
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 6 caplets in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 1 or more alcoholic drinks every day while using this product  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin redness  
 ■ hives  
 ■ rash  
 ■ if a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ new symptoms occur  
 ■ redness or swelling is present

**These could be signs of a serious condition.**  
 If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)  
 Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



How can we help?  
 1-877-895-3665

NDC 50580-783-29

See New Warning  
**TYLENOL<sup>®</sup> 8HR**  
**ARTHRITIS PAIN**

Acetaminophen  
 Extended-release tablets  
 Pain Reliever / Fever Reducer  
 For The Temporary Relief Of Minor Arthritis Pain

290 Caplets\*  
 650 mg each

\*Caplet-Shaped Bi-Layer Tablets

**TYLENOL<sup>®</sup> 8HR**  
**ARTHRITIS PAIN**

Acetaminophen  
 Extended-release tablets  
 Pain Reliever / Fever Reducer  
 For The Temporary Relief Of Minor Arthritis Pain

Contains No Aspirin



3 005-0838-29 2

**Drug Facts (continued)**

**Directions**  
 ■ do not take more than directed (see overdose warning)  
 adults  
 ■ take 2 caplets every 8 hours with water  
 ■ swallow whole; do not crush, chew, split or dissolve  
 ■ do not take more than 6 caplets in 24 hours  
 ■ do not use for more than 10 days unless directed by a doctor  
 ■ ask a doctor

under 18 years of age

**Other information**  
 ■ store between 20-25°C (68-77°F)  
 ■ do not use if carton is opened, or if foil inner seal imprinted with "SAFETY SEAL" is broken

**Inactive ingredients**  
 carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

**Questions or comments?**  
 call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



Distributed by:  
 McNeil Consumer Healthcare  
 Division of McNEIL-PPC, Inc.  
 Fort Washington, PA 19034 USA  
 © McNEIL-PPC, INC., 2016



00000000


Labeling Format Information: Fonts: HelveticaDF Medium, Bold, Bold Italic

Drug Facts: 10.00 pt	Body Text: 7.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%
Headers: 9.00 pt	Leading: 7.50 pt	Barlines: 2.50 pt	
Subheaders: 7.00 pt	Drug Facts (continued): 9.00 pt	Hairlines: 0.50 pt	Average Kerning: 0

McNeil Submission Artwork

(b) (4)

<p><b>Active ingredient (in each caplet)</b> Purpose                  Acetaminophen 650 mg..... Pain reliever/fever reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves minor aches and pains due to:                         <ul style="list-style-type: none"> <li>minor pain of arthritis</li> <li>muscular aches</li> <li>backache</li> <li>premenstrual and menstrual cramps</li> <li>the common cold</li> <li>headache</li> <li>toothache</li> </ul> </li> <li>temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount</p> <ul style="list-style-type: none"> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Allergy alert:</b> acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <ul style="list-style-type: none"> <li>if you are allergic to acetaminophen or any of the inactive ingredients in this product</li> </ul> <p><b>Ask a doctor before use if you have liver disease</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>pain gets worse or lasts more than 10 days</li> <li>fever gets worse or lasts more than 3 days</li> </ul>	<p><b>TO OPEN: PUSH AND TURN CAP</b>                  READ THE LABEL</p> <p><b>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</b></p> <p>NDC 50580-783-29</p> <p><b>TYLENOL 8HR</b>  <b>ARTHRITIS PAIN</b></p> <p>Acetaminophen                  Extended-release tablets                  Pain Reliever / Fever Reducer</p> <p>For The Temporary Relief                  Of Minor Arthritis Pain</p> <p>*Capsule-Shaped                  Bi-Layer Tablets</p> <p><b>290 Caplets*</b>  <b>650 mg each</b></p>	<ul style="list-style-type: none"> <li>new symptoms occur</li> <li>redness or swelling is present</li> </ul> <p>These could be signs of a serious condition.  <b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children.</b></p> <p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <p><b>do not take more than directed (see overdose warning)</b></p> <p><b>Adults:</b></p> <ul style="list-style-type: none"> <li>take 2 caplets every 8 hours with water</li> <li>swallow whole; do not crush, chew, split or dissolve</li> <li>do not take more than 6 caplets in 24 hours</li> <li>do not use for more than 10 days unless directed by a doctor</li> </ul> <p><b>Under 18 years of age:</b> ask a doctor</p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store between 20-25°C (68-77°F)</li> <li>do not use if foil inner seal imprinted with "SAFETY SEAL" is broken or missing</li> </ul> <p><b>Inactive ingredients</b></p> <p>ca nauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p><b>Questions or comments?</b>                  call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p> <p>Distributed by: McNeil Consumer Healthcare                  Division of McNEIL-PPC, Inc.                  Fort Washington, PA 19034 USA                  ©McNEIL-PPC, INC. 2016</p>
--	---	---

<b>Labeling Format Information:</b> Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%	
Headers: 8.00 pt	Leading: 7.80 pt	Barlines: NA pt	Average Kerning: -25	
Subheaders: 6.00 pt	Drug Facts (continued): NA pt	Hairlines: NA pt		

(b) (4)






Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%	
Headers: 4.50 pt	Leading: 4.70 pt	Barlines: NA pt	Average Kerning: -25	
Subheaders: 4.50 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt		

(b) (4)



<b>Active ingredient (in each caplet)</b>	<b>Purpose</b>	<b>TO OPEN: PUSH AND TURN CAP READ THE LABEL</b>	Keep out of reach of children.
<b>Acetaminophen 650 mg</b>	Pain reliever/fever reducer	NDC 50580-309-02	Overdose warning In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
<b>Uses</b>	temporarily relieves minor aches and pains due to:	<b>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</b>	<b>Directions</b> do not take more than directed (see overdose warning)
<ul style="list-style-type: none"> <li>■ muscular aches</li> <li>■ backache</li> <li>■ minor pain of arthritis</li> <li>■ toothache</li> <li>■ premenstrual and menstrual cramps</li> <li>■ headache</li> <li>■ the common cold</li> </ul>	temporarily reduces fever	<b>TYLENOL<sup>®</sup> 8HR</b>	Adults and children 12 years and over take 2 caplets every 8 hours with water
<b>Warnings</b>	<b>Liver warning</b>	<b>MUSCLE ACHES &amp; PAIN</b>	swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours
This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product.	Do not use with any other drug containing acetaminophen (prescription or nonprescription).	<b>Acetaminophen</b>	Children under 12 years do not use
Allergy alert acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.	Do not use if you are allergic to acetaminophen or any of the inactive ingredients in this product. Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present. These could be signs of a serious condition.	<b>Extended-release tablets</b>	<b>Other information</b> store between 20-25°C (68-77°F) do not use if foil inner seal is imprinted with "SAFETY SEAL" is broken or missing
If pregnant or breast-feeding, ask a health professional before use.	Pain Reliever / Fever Reducer	For Up to 8 Hours Relief of Minor Muscle Aches & Pain	<b>Inactive ingredients</b> carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, polydioxane, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin
		*Caplet-Shaped Bi-Layer Tablets	<b>Questions or comments?</b> call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)
		100 Caplets* 650 mg each	Distributed by: McNeil Consumer Healthcare Division of McNEI LPPC, Inc. Fort Washington, PA 19034 USA ©McNEI LPPC, INC. 2016

<b>Labeling Format Information:</b> Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 75%	
Headers: 6.00 pt	Leading: 6.50 pt	Barlines: NA pt	Average Kerning: -25	
Subheaders: 6.00 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt		

(b) (4)



<p style="font-size: 8pt;">A00000000</p> <p style="font-size: 8pt;">LOT: 101</p> <p style="font-size: 8pt;">EXP: 12/16</p>	<p><b>Active ingredient (in each caplet)</b>                  Acetaminophen 650 mg</p> <p><b>Purpose</b>                  Pain reliever/fever reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>■ temporarily relieves minor aches and pains due to:</li> <li>■ muscular aches    ■ backache    ■ minor pain of arthritis</li> <li>■ toothache        ■ premenstrual and menstrual cramps</li> <li>■ headache         ■ the common cold</li> <li>■ temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount. With other drugs containing acetaminophen. 3 or more alcoholic drinks every day while using this product. <b>Allergy alert</b> acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. <b>Ask a doctor before use</b> if you have liver disease. <b>Ask a doctor or pharmacist before use</b> if you are taking the blood thinning drug warfarin. <b>Stop use and ask a doctor</b> if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.</p>	<p style="text-align: center; font-weight: bold; font-size: 10pt;">TO OPEN: PUSH AND TURN CAP                  READ THE LABEL</p> <p style="font-size: 8pt;">NDC 50580-309-03</p> <p style="text-align: center; font-weight: bold; font-size: 8pt;">DO NOT USE WITH OTHER MEDICINES                  CONTAINING ACETAMINOPHEN</p> <p style="font-size: 12pt; font-weight: bold;">TYLENOL® 8HR</p> <p style="font-size: 10pt; font-weight: bold;">MUSCLE ACHES &amp; PAIN</p> <p style="font-size: 10pt; font-weight: bold;">Acetaminophen                  Extended-release tablets</p> <p style="font-size: 8pt;">Pain Reliever / Fever Reducer</p> <p style="font-size: 8pt;">For Up to 8 Hours Relief                  of Minor Muscle Aches &amp; Pain</p> <div style="text-align: center;">  <p style="font-size: 8pt;">*Capsule-Shaped                      Bi-Layer Tablets</p> <p style="font-size: 10pt; font-weight: bold;">225 Caplets*                      650 mg each</p> </div>	<p>If pregnant or breast-feeding, ask a health professional before use. <b>Keep out of reach of children.</b></p> <p><b>Overdose warning</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <p>■ do not take more than directed (see overdose warning)</p> <p><b>Adults and children 12 years and over</b> ■ take 2 caplets every 8 hours with water</p> <p>■ swallow whole; do not crush, chew, split or dissolve</p> <p>■ do not take more than 6 caplets in 24 hours</p> <p>■ do not use for more than 10 days unless directed by a doctor</p> <p><b>Children under 12 years</b> ■ do not use</p> <p><b>Other information</b></p> <p>■ store between 20-25°C (68-77°F)</p> <p>■ do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing</p> <p><b>Inactive ingredients</b> carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p><b>Questions or comments?</b> call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p> <p>Distributed by: McNeil Consumer Healthcare                  Division of McNEIL-PPC, Inc.                  Fort Washington, PA 19034 USA                  ©McNEIL-PPC, INC. 2016</p>
--	---	---	---

<b>Labeling Format Information:</b> Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: NA pt	
Subheaders: 6.00 pt	Drug Facts (continued): NA pt	Hairlines: NA pt	Average Kerning: 0

McNeil

Submission  
 Artwork

(b) (4)

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/  
-----

KAREN M MAHONEY  
08/26/2016