



0000



10 PIECES, OPEN HERE
2mg EACH



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.

Free Audio CD upon request. See inside.

This package size may not be a full day's supply; it is intended to start or continue a quit attempt.

NDC 0000-0000-00

not for sale to those under 18 years of age
 proof of age required
 not for sale in vending machines or from any source where proof of age cannot be verified

--- PRINT SIDE GLUE ---

No Copy Area-
 This area will contain an extended content label.

Distributed by
GlaxoSmithKline
 Consumer Healthcare, L.P.
 Moon Township, PA 15108
 Made in Sweden

000000XX
 ©2014 GSK

2 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.
 If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicorette 4mg Gum



Coated FOR BOLD FLAVOR
Cinnamon Surge™

10 PIECES, 2mg EACH

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

NICORETTE and the NICORETTE sunburst design are registered trademarks and CINNAMON SURGE is a trademark of the GlaxoSmithKline group of companies.

Theft surveillance tag area



0000

LOT EXP



0 0000-0000-00 0











0000

--- PRINT SIDE GLUE ---

No Copy Area-
This area will contain
an extended content
label.

Distributed by
GlaxoSmithKline
Consumer Healthcare, L.P.
Moon Township, PA 15108
Made in Sweden

000000XX
©2014 GSK

10 PIECES, OPEN HERE
4mg EACH

To remove the gum, tear off single unit. Peel off backing, starting at corner with loose edge. Push gum through foil.

This package size may not be a full day's supply; it is intended to start or continue a quit attempt. NDC 0000-0000-00

not for sale to those under 18 years of age
proof of age required
not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.

Free Audio CD upon request. See inside.

Nicorette
nicotine polacrilex gum, 4mg · stop smoking aid
Gum

Coated FOR BOLD FLAVOR
Fruit Chill™

4 mg FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP. If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicorette 2mg Gum

10 PIECES, 4mg EACH

NICORETTE, NICORETTE FRUIT CHILL and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Theft surveillance tag area

LOT EXP

0000-0000-00

0000



0000

--- PRINT SIDE GLUE ---

No Copy Area-
 This area will contain
 an extended content
 label.

Distributed by
GlaxoSmithKline
 Consumer Healthcare, L.P.
 Moon Township, PA 15108
 Made in Sweden

000000XX
 ©2014 GSK

- TO INCREASE YOUR SUCCESS IN QUITTING:**
1. You must be motivated to quit.
 2. **Use Enough** - Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
 3. **Use Long Enough** - Use Nicorette gum for the full 12 weeks.
 4. **Use with a support program** as directed in the enclosed User's Guide.
- For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.*
- Free Audio CD upon request. See inside.*

10 PIECES, 4mg EACH, OPEN HERE



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

This package size may not be a full day's supply; it is intended to start or continue a quit attempt.

NDC 0000-0000-00

Nicorette
 nicotine polacrilex gum, 4mg · stop smoking aid
Gum



Coated FOR BOLD FLAVOR

White Ice Mint

4 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP. If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicorette 2mg Gum

10 PIECES, 4mg EACH

■ not for sale to those under 18 years of age
 ■ proof of age required
 ■ not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

NICORETTE, WHITE ICE MINT and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.

Theft surveillance tag area



0000

LOT EXP



0 0000-0000-00 0


<p>Drug Facts (continued)</p> <p>tingle returns.</p> <ul style="list-style-type: none"> repeat this process until most of the tingle is gone (about 30 minutes) do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects. 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> do not use more than 24 pieces a day it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider. <p>Other information</p> <ul style="list-style-type: none"> each piece contains: calcium 94mg, sodium 11mg store at 20 - 25°C (68 - 77°F) protect from light <p>Inactive ingredients</p> <p>acacia, acesulfame potassium, carnauba wax, edible ink, gum base, hypromellose, magnesium oxide, menthol, natural</p>	<p>Drug Facts (continued)</p> <p>and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol</p> <p>Questions or comments?</p> <p>call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)</p> <p>©2014 GlaxoSmithKline NICORETTE and the NICORETTE sunburst design are registered trademarks and CINAMMON SURGE is a trademark of the GlaxoSmithKline group of companies.</p>	<p>Nicorette[®] Cinnamon Surge[™] Gum</p> <p>Drug Facts</p> <p>Active ingredient Purpose (in each chewing piece)</p> <p>Nicotine polacrilex (equal to 2mg nicotine) Stop smoking aid</p> <p>Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking</p> <p>Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child.</p> <p>000000XX</p>
---	--	--	---

<p>Drug Facts (continued)</p> <p>Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> a sodium-restricted diet heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. high blood pressure not controlled with medication. Nicotine can increase blood pressure. stomach ulcer or diabetes <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> using a non-nicotine stop smoking drug taking prescription medicine 	<p>Drug Facts (continued)</p> <p>for depression or asthma. Your prescription dose may need to be adjusted.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> mouth, teeth or jaw problems occur irregular heartbeat or palpitations occur you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat oral blistering occurs you have symptoms of an allergic reaction (such as difficulty breathing or rash) <p>Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in</p>	<p>Drug Facts (continued)</p> <p>paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> if you are under 18 years of age, ask a doctor before use before using this product, read the enclosed User's Guide for complete directions and other important information begin using the gum on your quit day if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum if you smoke your first cigarette within 30 	<p>Drug Facts (continued)</p> <p>minutes of waking up, use 4mg nicotine gum according to the following 12 week schedule:</p> <table border="1"> <tr> <td>Weeks 1 to 6</td> <td>1 piece every 1 to 2 hours</td> </tr> <tr> <td>Weeks 7 to 9</td> <td>1 piece every 2 to 4 hours</td> </tr> <tr> <td>Week 10 to 12</td> <td>1 piece every 4 to 8 hours</td> </tr> </table> <ul style="list-style-type: none"> nicotine gum is a medicine and must be used a certain way to get the best results chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the 	Weeks 1 to 6	1 piece every 1 to 2 hours	Weeks 7 to 9	1 piece every 2 to 4 hours	Week 10 to 12	1 piece every 4 to 8 hours
Weeks 1 to 6	1 piece every 1 to 2 hours								
Weeks 7 to 9	1 piece every 2 to 4 hours								
Week 10 to 12	1 piece every 4 to 8 hours								

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • repeat this process until most of the tingle is gone (about 30 minutes) • do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece • to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks • if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects. • do not use more than 	<p>Drug Facts (continued)</p> <p>24 pieces a day</p> <ul style="list-style-type: none"> • it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider. <p>Other information</p> <ul style="list-style-type: none"> • each piece contains: calcium 94mg, sodium 11mg • store at 20 - 25°C (68 - 77°F) • protect from light <p>Inactive ingredients acacia, acesulfame potassium, carnauba wax, edible ink, gum base, magnesium oxide, menthol, peppermint oil, sodium bicarbonate, sodium</p>	<p>Drug Facts (continued)</p> <p>carbonate, titanium dioxide, xylitol</p> <p>Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)</p> <p>©2014 GlaxoSmithKline NICORETTE and the NICORETTE sunburst design are registered trademarks and FRESH MINT is a trademark of the GlaxoSmithKline group of companies.</p>	<p>Nicorette Fresh Mint™ Gum</p> <p>Drug Facts</p> <p>Active ingredient Purpose (in each chewing piece) Nicotine polacrilex (equal to 2mg nicotine) Stop smoking aid</p> <p>Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking</p> <p>Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child.</p> <p>000000XX</p>
--	---	---	--

<p>Drug Facts (continued)</p> <p>Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • a sodium-restricted diet • heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. • high blood pressure not controlled with medication. Nicotine can increase blood pressure. • stomach ulcer or diabetes <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • using a non-nicotine stop smoking drug • taking prescription medicine 	<p>Drug Facts (continued)</p> <p>for depression or asthma. Your prescription dose may need to be adjusted.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • mouth, teeth or jaw problems occur • irregular heartbeat or palpitations occur • you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat • you have symptoms of an allergic reaction (such as difficulty breathing or rash) <p>Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the</p>	<p>Drug Facts (continued)</p> <p>trash. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> • if you are under 18 years of age, ask a doctor before use • before using this product, read the enclosed User's Guide for complete directions and other important information • begin using the gum on your quit day • if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum • if you smoke your first cigarette more than 30 minutes after waking up, 	<p>Drug Facts (continued)</p> <p>use 2mg nicotine gum according to the following 12 week schedule:</p> <table border="1"> <tr> <td>Weeks 1 to 6</td> <td>1 piece every 1 to 2 hours</td> </tr> <tr> <td>Weeks 7 to 9</td> <td>1 piece every 2 to 4 hours</td> </tr> <tr> <td>Week 10 to 12</td> <td>1 piece every 4 to 8 hours</td> </tr> </table> <ul style="list-style-type: none"> • nicotine gum is a medicine and must be used a certain way to get the best results • chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns. 	Weeks 1 to 6	1 piece every 1 to 2 hours	Weeks 7 to 9	1 piece every 2 to 4 hours	Week 10 to 12	1 piece every 4 to 8 hours
Weeks 1 to 6	1 piece every 1 to 2 hours								
Weeks 7 to 9	1 piece every 2 to 4 hours								
Week 10 to 12	1 piece every 4 to 8 hours								

Drug Facts (continued)

- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 


Drug Facts (continued)

- 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- each piece contains: calcium 94mg, sodium 11mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, 

Drug Facts (continued)

sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

Questions or comments?

call toll-free
1-800-419-4766
(English/Spanish)
weekdays (9:00 am - 4:30 pm ET)

©2014 GlaxoSmithKline
NICORETTE, WHITE ICE MINT and the NICORETTE sunburst design are registered trademarks of the GlaxoSmithKline group of companies.

Nicorette[®]
White Ice Mint[®]
Gum

Drug Facts

Active ingredient Purpose (in each chewing piece)

Nicotine polacrilex (equal to 2mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. 

000000XX

Drug Facts (continued)

Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine 


Drug Facts (continued)

for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)


Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the 

Drug Facts (continued)

trash. In case of overdose, get medical help or contact a Poison Control Center right away.


Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, 

Drug Facts (continued)

use 2mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	1 piece every 1 to 2 hours
Weeks 7 to 9	1 piece every 2 to 4 hours
Week 10 to 12	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns. 

Drug Facts (continued)

- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than

Drug Facts (continued)

- 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- **each piece contains:** calcium 94mg, sodium 11mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80,

Drug Facts (continued)

sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

Questions or comments?

call toll-free
1-800-419-4766
(English/Spanish)
weekdays (9:00 am - 4:30 pm ET)

©2014 GlaxoSmithKline

NICORETTE, NICORETTE FRUIT CHILL and the NICORETTE sunburst design are registered trademarks of the GlaxoSmithKline group of companies.

**Nicorette[®]
Fruit Chill[™]
Gum**

Drug Facts

Active ingredient Purpose (in each chewing piece)

Nicotine polacrilex (equal to 2mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child.

000000XX

Drug Facts (continued)

Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine

Drug Facts (continued)

for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the

Drug Facts (continued)

trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum**
- **if you smoke your first cigarette more than 30 minutes after waking up,**

Drug Facts (continued)

use 2mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	1 piece every 1 to 2 hours
Weeks 7 to 9	1 piece every 2 to 4 hours
Week 10 to 12	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.

<p>Drug Facts (continued)</p> <p>tingle returns.</p> <ul style="list-style-type: none"> repeat this process until most of the tingle is gone (about 30 minutes) do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects. 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> do not use more than 24 pieces a day it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider. <p>Other information</p> <ul style="list-style-type: none"> each piece contains: calcium 94mg, sodium 13mg store at 20 - 25°C (68 - 77°F) protect from light <p>Inactive ingredients</p> <p>aacacia, acesulfame potassium, carnauba wax, D&C yellow #10 AL lake, edible ink, gum base, hypromellose, magnesium</p>	<p>Drug Facts (continued)</p> <p>oxide, menthol, natural and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium carbonate, sucralose, titanium dioxide, xylitol</p> <p>Questions or comments?</p> <p>call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)</p> <p>©2014 GlaxoSmithKline NICORETTE and the NICORETTE sunburst design are registered trademarks and CINAMMON SURGE is a trademark of the GlaxoSmithKline group of companies.</p>	<p>Nicorette Cinnamon Surge™ Gum</p> <p>Drug Facts</p> <p>Active ingredient Purpose (in each chewing piece)</p> <p>Nicotine polacrilex (equal to 4mg nicotine) Stop smoking aid</p> <p>Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking</p> <p>Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child.</p> <p>000000XX</p>
---	---	--	--

<p>Drug Facts (continued)</p> <p>Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> a sodium-restricted diet heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. high blood pressure not controlled with medication. Nicotine can increase blood pressure. stomach ulcer or diabetes <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> using a non-nicotine stop smoking drug taking prescription medicine 	<p>Drug Facts (continued)</p> <p>for depression or asthma. Your prescription dose may need to be adjusted.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> mouth, teeth or jaw problems occur irregular heartbeat or palpitations occur you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat oral blistering occurs you have symptoms of an allergic reaction (such as difficulty breathing or rash) <p>Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in</p>	<p>Drug Facts (continued)</p> <p>paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> if you are under 18 years of age, ask a doctor before use before using this product, read the enclosed User's Guide for complete directions and other important information begin using the gum on your quit day if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum if you smoke your first cigarette within 30 	<p>Drug Facts (continued)</p> <p>minutes of waking up, use 4mg nicotine gum according to the following 12 week schedule:</p> <table border="1"> <tr> <td>Weeks 1 to 6</td> <td>1 piece every 1 to 2 hours</td> </tr> <tr> <td>Weeks 7 to 9</td> <td>1 piece every 2 to 4 hours</td> </tr> <tr> <td>Week 10 to 12</td> <td>1 piece every 4 to 8 hours</td> </tr> </table> <ul style="list-style-type: none"> nicotine gum is a medicine and must be used a certain way to get the best results chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the 	Weeks 1 to 6	1 piece every 1 to 2 hours	Weeks 7 to 9	1 piece every 2 to 4 hours	Week 10 to 12	1 piece every 4 to 8 hours
Weeks 1 to 6	1 piece every 1 to 2 hours								
Weeks 7 to 9	1 piece every 2 to 4 hours								
Week 10 to 12	1 piece every 4 to 8 hours								

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> repeat this process until most of the tingle is gone (about 30 minutes) do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects. do not use more than 	<p>Drug Facts (continued)</p> <p>24 pieces a day</p> <ul style="list-style-type: none"> it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider. <p>Other information</p> <ul style="list-style-type: none"> each piece contains: calcium 94mg, sodium 13mg store at 20 - 25°C (68 - 77°F) protect from light <p>Inactive ingredients acacia, acesulfame potassium, carnauba wax, D&C yellow #10 Al. lake, edible ink, gum base, magnesium oxide, menthol, peppermint oil, sodium</p>	<p>Drug Facts (continued)</p> <p>carbonate, titanium dioxide, xylitol</p> <p>Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)</p> <p>©2014 GlaxoSmithKline NICORETTE and the NICORETTE sunburst design are registered trademarks and FRESH MINT is a trademark of the GlaxoSmithKline group of companies.</p>	<p>Nicorette Fresh Mint™ Gum</p> <p>Drug Facts</p> <p>Active ingredient Purpose (in each chewing piece) Nicotine polacrilex (equal to 4mg nicotine) Stop smoking aid</p> <p>Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking</p> <p>Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child.</p> <p>000000XX</p>
--	--	--	--

<p>Drug Facts (continued)</p> <p>Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> a sodium-restricted diet heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. high blood pressure not controlled with medication. Nicotine can increase blood pressure. stomach ulcer or diabetes <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> using a non-nicotine stop smoking drug taking prescription medicine 	<p>Drug Facts (continued)</p> <p>for depression or asthma. Your prescription dose may need to be adjusted.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> mouth, teeth or jaw problems occur irregular heartbeat or palpitations occur you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat you have symptoms of an allergic reaction (such as difficulty breathing or rash) <p>Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the</p>	<p>Drug Facts (continued)</p> <p>trash. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> if you are under 18 years of age, ask a doctor before use before using this product, read the enclosed User's Guide for complete directions and other important information begin using the gum on your quit day if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum if you smoke your first cigarette within 30 minutes of waking up, 	<p>Drug Facts (continued)</p> <p>use 4mg nicotine gum according to the following 12 week schedule:</p> <table border="1"> <tr> <td>Weeks 1 to 6</td> <td>1 piece every 1 to 2 hours</td> </tr> <tr> <td>Weeks 7 to 9</td> <td>1 piece every 2 to 4 hours</td> </tr> <tr> <td>Week 10 to 12</td> <td>1 piece every 4 to 8 hours</td> </tr> </table> <ul style="list-style-type: none"> nicotine gum is a medicine and must be used a certain way to get the best results chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns. 	Weeks 1 to 6	1 piece every 1 to 2 hours	Weeks 7 to 9	1 piece every 2 to 4 hours	Week 10 to 12	1 piece every 4 to 8 hours
Weeks 1 to 6	1 piece every 1 to 2 hours								
Weeks 7 to 9	1 piece every 2 to 4 hours								
Week 10 to 12	1 piece every 4 to 8 hours								

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • repeat this process until most of the tingle is gone (about 30 minutes) • do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece • to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks • if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects. • do not use more than 	<p>Drug Facts (continued)</p> <p>24 pieces a day</p> <ul style="list-style-type: none"> • it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider. <p>Other information</p> <ul style="list-style-type: none"> • each piece contains: calcium 94mg, sodium 13mg • store at 20 - 25°C (68 - 77°F) • protect from light <p>Inactive ingredients</p> <p>acesulfame potassium, carnauba wax, D&C yellow #10 AL lake, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, </p>	<p>Drug Facts (continued)</p> <p>peppermint oil, polysorbate 80, sodium carbonate, starch, sucralose, titanium dioxide, xylitol</p> <p>Questions or comments?</p> <p>call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)</p> <p>©2014 GlaxoSmithKline NICORETTE, WHITE ICE MINT and the NICORETTE sunburst design are registered trademarks of the GlaxoSmithKline group of companies.</p>	<p style="font-size: 24pt; font-weight: bold; margin: 0;">Nicorette®</p> <p style="font-size: 18pt; font-weight: bold; margin: 0;">White Ice Mint®</p> <p style="font-size: 18pt; font-weight: bold; margin: 0;">Gum</p> <hr/> <p>Drug Facts</p> <p>Active ingredient Purpose (in each chewing piece)</p> <p>Nicotine polacrilex (equal to 4mg nicotine)..... Stop smoking aid</p> <p>Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking</p> <p>Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. </p> <p style="text-align: right; font-size: 10pt;">000000XX</p>
---	--	---	---

<p>Drug Facts (continued)</p> <p>Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • a sodium-restricted diet • heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. • high blood pressure not controlled with medication. Nicotine can increase blood pressure. • stomach ulcer or diabetes <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • using a non-nicotine stop smoking drug • taking prescription medicine 	<p>Drug Facts (continued)</p> <p>for depression or asthma. Your prescription dose may need to be adjusted.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • mouth, teeth or jaw problems occur • irregular heartbeat or palpitations occur • you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat • you have symptoms of an allergic reaction (such as difficulty breathing or rash) <p>Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the </p>	<p>Drug Facts (continued)</p> <p>trash. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> • if you are under 18 years of age, ask a doctor before use • before using this product, read the enclosed User's Guide for complete directions and other important information • begin using the gum on your quit day • if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum • if you smoke your first cigarette within 30 minutes of waking up, 	<p>Drug Facts (continued)</p> <p>use 4mg nicotine gum according to the following 12 week schedule:</p> <table border="1" style="width: 100%; border-collapse: collapse; margin: 5px 0;"> <tr> <td style="padding: 2px;">Weeks 1 to 6</td> <td style="padding: 2px;">1 piece every 1 to 2 hours</td> </tr> <tr> <td style="padding: 2px;">Weeks 7 to 9</td> <td style="padding: 2px;">1 piece every 2 to 4 hours</td> </tr> <tr> <td style="padding: 2px;">Week 10 to 12</td> <td style="padding: 2px;">1 piece every 4 to 8 hours</td> </tr> </table> <ul style="list-style-type: none"> • nicotine gum is a medicine and must be used a certain way to get the best results • chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns. 	Weeks 1 to 6	1 piece every 1 to 2 hours	Weeks 7 to 9	1 piece every 2 to 4 hours	Week 10 to 12	1 piece every 4 to 8 hours
Weeks 1 to 6	1 piece every 1 to 2 hours								
Weeks 7 to 9	1 piece every 2 to 4 hours								
Week 10 to 12	1 piece every 4 to 8 hours								




<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • repeat this process until most of the tingle is gone (about 30 minutes) • do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece • to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks • if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects. • do not use more than 	<p>Drug Facts (continued)</p> <p>24 pieces a day</p> <ul style="list-style-type: none"> • it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider. <p>Other information</p> <ul style="list-style-type: none"> • each piece contains: calcium 94mg, sodium 13mg • store at 20 - 25°C (68 - 77°F) • protect from light <p>Inactive ingredients acacia, acesulfame potassium, carnauba wax, D&C yellow #10 AL lake, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint </p>	<p>Drug Facts (continued)</p> <p>oil, polysorbate 80, sodium carbonate, sucralose, titanium dioxide, xylitol</p> <p>Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)</p> <p>©2014 GlaxoSmithKline NICORETTE, NICORETTE FRUIT CHILL and the NICORETTE sunburst design are registered trademarks of the GlaxoSmithKline group of companies.</p>	<p style="text-align: center;">Nicorette[®] Fruit Chill[™] Gum</p> <p>Drug Facts</p> <p>Active ingredient Purpose (in each chewing piece) Nicotine polacrilex (equal to 4mg nicotine) Stop smoking aid</p> <p>Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking</p> <p>Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. </p> <p style="text-align: right;">000000XX</p>
---	---	---	--

<p>Drug Facts (continued)</p> <p>Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • a sodium-restricted diet • heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. • high blood pressure not controlled with medication. Nicotine can increase blood pressure. • stomach ulcer or diabetes <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • using a non-nicotine stop smoking drug • taking prescription medicine 	<p>Drug Facts (continued)</p> <p>for depression or asthma. Your prescription dose may need to be adjusted.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • mouth, teeth or jaw problems occur • irregular heartbeat or palpitations occur • you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat • you have symptoms of an allergic reaction (such as difficulty breathing or rash) <p>Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the </p>	<p>Drug Facts (continued)</p> <p>trash. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> • if you are under 18 years of age, ask a doctor before use • before using this product, read the enclosed User's Guide for complete directions and other important information • begin using the gum on your quit day • if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum • if you smoke your first cigarette within 30 minutes of waking up, 	<p>Drug Facts (continued)</p> <p>use 4mg nicotine gum according to the following 12 week schedule:</p> <table border="1" style="width: 100%; border-collapse: collapse; margin-bottom: 5px;"> <tr> <td style="padding: 2px;">Weeks 1 to 6</td> <td style="padding: 2px;">1 piece every 1 to 2 hours</td> </tr> <tr> <td style="padding: 2px;">Weeks 7 to 9</td> <td style="padding: 2px;">1 piece every 2 to 4 hours</td> </tr> <tr> <td style="padding: 2px;">Week 10 to 12</td> <td style="padding: 2px;">1 piece every 4 to 8 hours</td> </tr> </table> <ul style="list-style-type: none"> • nicotine gum is a medicine and must be used a certain way to get the best results • chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns. 	Weeks 1 to 6	1 piece every 1 to 2 hours	Weeks 7 to 9	1 piece every 2 to 4 hours	Week 10 to 12	1 piece every 4 to 8 hours
Weeks 1 to 6	1 piece every 1 to 2 hours								
Weeks 7 to 9	1 piece every 2 to 4 hours								
Week 10 to 12	1 piece every 4 to 8 hours								


MANUFACTURER'S COUPON **EXPIRES 00/00/0000**


SAVE \$1.00 on any two Nicorette® Gum 10 Pieces


Redeemable at food, drug and mass merchandise stores. CONSUMER: Limit 1 coupon per purchase of indicated product(s). You pay any sales tax. You are not eligible for this offer if reimbursement for the product will be sought from any federal or state healthcare program, including Medicare or Medicaid, or any similar federal or state programs, including any state pharmaceutical assistance program, or under any private insurance, HMO, or other third-party payment arrangement. **RETAILER:** GSK will reimburse you the face value of this coupon plus 12¢ Handling Fee if submitted in compliance with our Redemption Policy. Copy available upon request. Cash value 1/100 of 1¢. Coupon is void if altered, transferred, exchanged, sold or auctioned, copied or if prohibited by law. Good only in USA and at APOs, FPOs. **Send coupons to:** GlaxoSmithKline Consumer Healthcare, P.O. Box 880658, El Paso, Texas 88588-0658. ©2013 GlaxoSmithKline. Read and follow label directions.



Go to www.nicorette.com for additional support and offers








MANUFACTURER'S COUPON EXPIRES 00/00/0000

**SAVE
\$5.00**
on any
Nicorette® Gum
100 Pieces or Larger



Redeemable at food, drug and mass merchandise stores. CONSUMER: Limit 1 coupon per purchase of indicated product(s). You pay any sales tax. You are not eligible for this offer if reimbursement for the product will be sought from any federal or state healthcare program, including Medicare or Medicaid, or any similar federal or state programs, including any state pharmaceutical assistance program, or under any private insurance, HMO, or other third-party payment arrangement. RETAILER: GSK will reimburse you the face value of this coupon plus 12¢ Handling Fee if submitted in compliance with our Redemption Policy. Copy available upon request. Cash value 1/100 of 1¢. Coupon is void if altered, transferred, exchanged, sold or auctioned, copied or if prohibited by law. Good only in USA and at APOs, FPOs. **Send coupons to:** GlaxoSmithKline Consumer Healthcare, P.O. Box 880658, El Paso, Texas 88588-0658. ©2013 GlaxoSmithKline. Read and follow label directions.



Go to www.nicorette.com
for additional support and offers

