



**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	N/A
DRUG FACTS CONTINUED	N/A
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.2 pt.
# OF CHARACTERS PER INCH	N/A
BULLETS	N/A
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	N/A
SPACE BETWEEN HAIRLINES AND BOX END	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	27%

OUTSIDE COPY

**\$5.00**  
Coupon Inside

**\$5.00**  
Coupon Inside

NDC 0135-0241-08 **HALEON**

# NICORETTE

## Gum

**20 PIECES, 2 mg EACH**  
(1 Pocket Pack of 20)

**Drug Facts**

**Active ingredient (in each chewing piece) Purpose**  
Nicotine polacrilex (equal to 2 mg nicotine).....Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Flip open for Directions and additional information**  
**Retain this package for complete product information**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

The gum is contained in a carton packaged inside a clear plastic outer container sealed to a printed card. The inner carton has a clear overwrap.  
**DO NOT USE IF ANY OF THESE TAMPER EVIDENT FEATURES ARE MISSING, TORN, OR BROKEN.**

Lot

Exp

3 07667 84914 1

INSIDE COPY

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

**How To Open the Nicorette Gum Pocket Pack**

**1** Gently push tab on side.

**2** Pull up forcefully, pulling open box.

**3** Dispense Nicorette Gum and follow dosing instructions.

**Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.**

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

204492

**Drug Facts (continued)**

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F) • protect from light and humidity

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

REGULATORY LEGEND

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

OPEN HERE

**Drug Facts**

**Active ingredient (in each chewing piece)** **Purpose**  
Nicotine polacrilex (equal to 2 mg nicotine) Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program

**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

**100 PIECES, 2 mg EACH**

To remove the gum, tear off single unit

Peel off backing, starting at corner with loose edge.

Push gum through foil.

NICORETTE  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Fruit Chill Gum**

**100 PIECES, 2 mg EACH**

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

**100 PIECES, 2 mg EACH**

**Fruit Chill**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Fruit Chill Gum**

**100 PIECES, 2 mg EACH**

**LOT:**

**EXP:**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Fruit Chill Gum**

**100 PIECES, 2 mg EACH**

**LOT:**

**EXP:**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Fruit Chill Gum**

**100 PIECES, 2 mg EACH**

**LOT:**

**EXP:**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Fruit Chill Gum**

**100 PIECES, 2 mg EACH**

**LOT:**

**EXP:**

824741

3 07667 85750 4

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.





**160 PIECES, 2 mg EACH**

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**2 mg**

**Fruit Chill Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

NDC 0135-0241-05

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**Fruit Chill**

**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH**

824742

Includes Quit Support Program



**2 mg**

**Fruit Chill Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

NDC 0135-0241-05

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**Fruit Chill**

**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Fruit Chill Gum**

824742

LOT: [REDACTED]

EXP: [REDACTED]

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Fruit Chill Gum**

**not for sale to those under 18 years of age**

**proof of age required**

**not for sale in vending machines or from any source where proof of age cannot be verified**

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 87701 4

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	N/A
DRUG FACTS CONTINUED	N/A
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.2 pt.
# OF CHARACTERS PER INCH	N/A
BULLETS	N/A
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	N/A
SPACE BETWEEN HAIRLINES AND BOX END	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	27%



**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program



**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NDC 0135-0474-02


HALEON

# NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

## White Ice Mint Gum

**100 PIECES, 2 mg EACH**



**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

ACTUAL SIZE

**White Ice Mint**

COATED FOR BOLD FLAVOR **100 PIECES, 2 mg EACH**



3 0766775000 3

824747

824747

LOT:

EXP:

824747

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**White Ice Mint Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polyorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

2 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**White Ice Mint**

**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH**

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**White Ice Mint Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polyorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

2 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**White Ice Mint**

**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH**

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit



Peel off backing, starting at corner with loose edge.



Push gum through foil.

160 PIECES, 2 mg EACH

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**White Ice Mint Gum**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.

Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 75025 6

824748

LOT:

EXP:

824748

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

OPEN HERE

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- oral blistering occurs
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.

Includes Quit Support Program

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2mg**

**Cinnamon Surge Gum**

**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, gum base, hypromellose, magnesium oxide, menthol, natural and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NOC 0125-0466-02 HALEON

**100 PIECES, 2 mg EACH**

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2mg**

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

**COATED FOR BOLD FLAVOR** **100 PIECES, 2 mg EACH** **Cinnamon Surge**

ACTUAL SIZE

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Cinnamon Surge Gum**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program!*  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.

3 07667 85840 2

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

### Drug Facts

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- oral blistering occurs
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**Cinnamon Surge Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, gum base, hypromellose, magnesium oxide, menthol, natural and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit



Peel off backing, starting at corner with loose edge.



Push gum through foil.

**160 PIECES, 2 mg EACH**

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**Cinnamon Surge Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, gum base, hypromellose, magnesium oxide, menthol, natural and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**Cinnamon Surge**

**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Cinnamon Surge Gum**

824736

LOT: [REDACTED]

EXP: [REDACTED]

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Cinnamon Surge Gum**

■ not for sale to those under 18 years of age  
■ proof of age required  
■ not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.


Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 85820 4

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



OPEN HERE

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.


**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program



**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**


- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavors, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NDC 0135-0532-02

HALEON



**100 PIECES, 2 mg EACH**

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

ACTUAL SIZE  
**Spearmint Burst**  
with a hint of chamomile

**COATED FOR BOLD FLAVOR**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Spearmint Burst Gum**  
with a hint of chamomile

• not for sale to those under 18 years of age

• proof of age required


• not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor. Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program!*  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 77000 1



824758

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Spearmint Burst Gum**  
with a hint of chamomile

824758

LOT:

EXP:

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

**160 PIECES, 2 mg EACH**

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equivalent to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** — reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.


**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavors, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

NDC 0135-0532-03

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.  
If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**Spearmint Burst**  
with a hint of chamomile

COATED FOR BOLD FLAVOR

**160 PIECES, 2 mg EACH**

824759

824759

824759

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Spearmint Burst Gum**  
with a hint of chamomile

824759

LOT:

EXP:

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Spearmint Burst Gum**  
with a hint of chamomile


- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 84360 6

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)** **Purpose**  
Nicotine polacrilex (equal to 2 mg nicotine) Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program



**110 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 117 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
flavors, glycerin, gum base, sodium bicarbonate, sodium carbonate, sorbitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NDC 0135-0157-07

HALEON

# NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

**Original Gum**

not for sale to those under 18 years of age

proof of age required

not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 84508 2

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

110 PIECES, 2 mg EACH

Original



**110 PIECES, 2 mg EACH**

LOT:

EXP:



824752

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.





**170 PIECES, 2 mg EACH**

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**170 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 117 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
flavors, glycerin, gum base, sodium bicarbonate, sodium carbonate, sorbitol

**Questions or comments?** call toll-free 1-888-569-1743

NDC 0135-0157-10

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**170 PIECES, 2 mg EACH** **Original**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Original Gum**

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.*



3 07667 84560 0

824753

824753

LOT:

EXP:

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.





200 PIECES, 2 mg EACH

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**200 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 117 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
flavors, glycerin, gum base, sodium bicarbonate, sodium carbonate, sorbitol

**Questions or comments?** call toll-free 1-888-569-1743

HALEON

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Gum

2 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

200 PIECES, 2 mg EACH Original

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT:
EXP:

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

8247

OUTSIDE COPY

NDC 0135-0474-08 **HALEON**

**\$5.00**  
Coupon Inside

**Drug Facts**

**Active ingredient (in each chewing piece)**    **Purpose**  
Nicotine polacrilex (equal to 2 mg nicotine).....Stop smoking aid

**Use**    • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Flip open for Directions and additional information**  
**Retain this package for complete product information**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

The gum is contained in a carton packaged inside a clear plastic outer container sealed to a printed card. The inner carton has a clear overwrap. **DO NOT USE IF ANY OF THESE TAMPER EVIDENT FEATURES ARE MISSING, TORN, OR BROKEN.**

Lot  
Exp

3 0766775054 6

NICORETTE

Gum

20 PIECES, 2 mg EACH  
(1 Pocket Pack of 20)

INSIDE COPY

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

How To Open the Nicorette Gum Pocket Pack

1

**Gently push tab on side.**

2

**Pull up forcefully, pulling open box.**

3

**Dispense Nicorette Gum and follow dosing instructions.**

Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

204496

**Drug Facts (continued)**

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F) • protect from light and humidity

**Inactive ingredients**  
acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

REGULATORY LEGEND

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.





**170 PIECES, 2 mg EACH**

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**170 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acesulfame potassium, gum base, magnesium oxide, menthol, peppermint oil, sodium bicarbonate, sodium carbonate, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

NICO RETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**Mint Gum**

**170 PIECES, 2 mg EACH**

NOC 0135-0229-05 HALEON

**NICO RETTE**

nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**Mint**

**170 PIECES, 2 mg EACH**

**NICO RETTE**

nicotine polacrilex gum, 2 mg | stop smoking aid

**Mint Gum**

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.*



3 07667 84340 8

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	N/A
DRUG FACTS CONTINUED	N/A
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.2 pt.
# OF CHARACTERS PER INCH	N/A
BULLETS	N/A
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	N/A
SPACE BETWEEN HAIRLINES AND BOX END	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	27%

OUTSIDE COPY

**\$5.00**  
Coupon Inside

**\$5.00**  
Coupon Inside

NDC 0135-0241-08

**HALEON**

## NICORETTE

### Gum

**20 PIECES, 2 mg EACH**  
(1 Pocket Pack of 20)

**Drug Facts**

**Active ingredient (in each chewing piece)**    **Purpose**  
Nicotine polacrilex (equal to 2 mg nicotine).....Stop smoking aid

**Use**    • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Flip open for Directions and additional information**  
**Retain this package for complete product information**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

The gum is contained in a carton packaged inside a clear plastic outer container sealed to a printed card. The inner carton has a clear overwrap.  
**DO NOT USE IF ANY OF THESE TAMPER EVIDENT FEATURES ARE MISSING, TORN, OR BROKEN.**

Lot

Exp

3 07667 84914 1

INSIDE COPY

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

**How To Open the Nicorette Gum Pocket Pack**

**1** Gently push tab on side.

**2** Pull up forcefully, pulling open box.

**3** Dispense Nicorette Gum and follow dosing instructions.

**Nicorette is proud to offer a free individualized quit support program!**  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

204492

**Drug Facts (continued)**

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F) • protect from light and humidity

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

REGULATORY LEGEND

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)** **Purpose**  
Nicotine polacrilex (equal to 2 mg nicotine) Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program



**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NDC 0135-0241-02 HALEON



**100 PIECES, 2 mg EACH**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit



Peel off backing, starting at corner with loose edge.



Push gum through foil.

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

ACTUAL SIZE  
**Fruit Chill**

**COATED FOR BOLD FLAVOR** **100 PIECES, 2 mg EACH**

**824741**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Fruit Chill Gum**

not for sale to those under 18 years of age  
proof of age required  
not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 85750 4

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Fruit Chill Gum**

LOT:

EXP:

824741

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

**160 PIECES, 2 mg EACH**

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**2 mg**

**Fruit Chill Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

NDC 0135-0241-05

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

**Fruit Chill**

**160 PIECES, 2 mg EACH**

COATED FOR BOLD FLAVOR

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Fruit Chill Gum**

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.*



3 07667 87701 4

824742

824742

LOT: XXXXXXXXXX

EXP: XXXXXXXXXX

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	N/A
DRUG FACTS CONTINUED	N/A
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.2 pt.
# OF CHARACTERS PER INCH	N/A
BULLETS	N/A
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	N/A
SPACE BETWEEN HAIRLINES AND BOX END	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	27%

OUTSIDE COPY

**\$5.00**  
Coupon Inside

**\$5.00**  
Coupon Inside

NDC 0135-0474-08

HALEON

# NICORETTE

## Gum

**20 PIECES, 2 mg EACH**  
(1 Pocket Pack of 20)

**Drug Facts**

**Active ingredient (in each chewing piece)** Purpose  
Nicotine polacrilex (equal to 2 mg nicotine).....Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Flip open for Directions and additional information**  
**Retain this package for complete product information**

Lot  
Exp

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

The gum is contained in a carton packaged inside a clear plastic outer container sealed to a printed card. The inner carton has a clear overwrap.

**DO NOT USE IF ANY OF THESE TAMPER EVIDENT FEATURES ARE MISSING, TORN, OR BROKEN.**

3 07667 75054 6

INSIDE COPY

**\$5.00**  
Coupon Inside

**\$5.00**  
Coupon Inside

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

**How To Open the Nicorette Gum Pocket Pack**

1

Gently push tab on side.

2

Pull up forcefully, pulling open box.

3

Dispense Nicorette Gum and follow dosing instructions.

Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

204496

**Drug Facts (continued)**

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F) • protect from light and humidity

**Inactive ingredients**  
acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

REGULATORY LEGEND

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program

**White Ice Mint Gum**

**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NDC 0135-0474-02

HALEON

**NICORETTE**

nicotine polacrilex gum, 2 mg | stop smoking aid

**White Ice Mint Gum**

**100 PIECES, 2 mg EACH**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

ACTUAL SIZE

**White Ice Mint**

COATED FOR BOLD FLAVOR **100 PIECES, 2 mg EACH**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**White Ice Mint Gum**

824747

LOT:

EXP:

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

To remove the gum, tear off single unit

Peel off backing, starting at corner with loose edge.

Push gum through foil.

**White Ice Mint Gum**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program!*  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.

3 0766775000 3

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**White Ice Mint Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polyorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH** **White Ice Mint**

ACTUAL SIZE

824748

LOT: XXXXXXXXXX

EXP: XXXXXXXXXX

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**White Ice Mint Gum**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 75025 6

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

OPEN HERE

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- oral blistering occurs
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour.

However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.

Includes Quit Support Program

**2mg**

**Cinnamon Surge Gum**

**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, gum base, hypromellose, magnesium oxide, menthol, natural and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NDC 0135-0466-02

HALEON

**2mg**

**Cinnamon Surge Gum**

**100 PIECES, 2 mg EACH**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

To remove the gum, tear off single unit

Peel off backing, starting at corner with loose edge.

Push gum through foil.

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

ACTUAL SIZE

**Cinnamon Surge**

**COATED FOR BOLD FLAVOR 100 PIECES, 2 mg EACH**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Cinnamon Surge Gum**

not for sale to those under 18 years of age

proof of age required

not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.

3 07667 85840 2

824735

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Cinnamon Surge Gum**

824735

LOT:

EXP:

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- oral blistering occurs
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**Cinnamon Surge Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, gum base, hypromellose, magnesium oxide, menthol, natural and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit



Peel off backing, starting at corner with loose edge.



Push gum through foil.

**160 PIECES, 2 mg EACH**

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**Cinnamon Surge Gum**

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, gum base, hypromellose, magnesium oxide, menthol, natural and artificial cinnamon flavors, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**



**2 mg**

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



**Cinnamon Surge**

ACTUAL SIZE

**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Cinnamon Surge Gum**

824736

LOT: [REDACTED]

EXP: [REDACTED]

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Cinnamon Surge Gum**

■ not for sale to those under 18 years of age  
■ proof of age required  
■ not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.

Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.*



3 07667 85820 4

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program



**2mg**

**Original Gum**

**110 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- each piece contains: calcium 117 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
flavors, glycerin, gum base, sodium bicarbonate, sodium carbonate, sorbitol

**Questions or comments?**  
call toll-free 1-888-569-1743

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit



Peel off backing, starting at corner with loose edge.



Push gum through foil.

**110 PIECES, 2 mg EACH**

824752

824752

LOT:

EXP:

824752

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

not for sale to those under 18 years of age

proof of age required

not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.

NICORETTE

nicotine polacrilex gum, 2 mg  
stop smoking aid  
Gum



**2mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

110 PIECES, 2 mg EACH

Original



3 07667 84508 2

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.





**170 PIECES, 2 mg EACH**

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**170 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 117 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
flavors, glycerin, gum base, sodium bicarbonate, sodium carbonate, sorbitol

**Questions or comments?** call toll-free 1-888-569-1743

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.  
If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



**170 PIECES, 2 mg EACH** **Original**

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.*



3 07667 84560 0

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Original Gum**

824753

LOT: XXXXXXXXXX

EXP: XXXXXXXXXX

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.





200 PIECES, 2 mg EACH

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**200 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 117 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
flavors, glycerin, gum base, sodium bicarbonate, sodium carbonate, sorbitol

**Questions or comments?** call toll-free 1-888-569-1743

HALEON

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Gum

2 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE

200 PIECES, 2 mg EACH

Original

824754

824754

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

LOT: [REDACTED]

EXP: [REDACTED]

NICORETTE

nicotine polacrilex gum, 2 mg | stop smoking aid

Original Gum

824754

not for sale to those under 18 years of age

proof of age required

not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.

Trademarks are owned by or licensed to the Haleon group of companies.


Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07662 86210 7

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.


**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

Includes Quit Support Program



**100 PIECES, 2 mg EACH**

**Drug Facts (continued)**

- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information**


- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavors, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

NDC 0135-0532-02

HALEON



**100 PIECES, 2 mg EACH**

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum

ACTUAL SIZE  
**Spearmint Burst**  
with a hint of chamomile

**COATED FOR BOLD FLAVOR**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Spearmint Burst Gum**  
with a hint of chamomile

• not for sale to those under 18 years of age

• proof of age required


• not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. **Do not use if individual blisters or printed backings are broken, open, or torn.**

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor. Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program!*  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 77000 1



824758

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Spearmint Burst Gum**  
with a hint of chamomile

824758

LOT:

EXP:

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt.
DRUG FACTS CONTINUED	8 pt.
HEADINGS	8 pt.
SUBHEADINGS/BODY TEXT	6 pt.
LEADING	6.5 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**2 mg**

**Spearmint Burst Gum**  
with a hint of chamomile

**160 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acacia, acesulfame potassium, carnauba wax, edible ink, flavors, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

**VALUE PACK** HALEON

**NICORETTE**  
nicotine polacrilex gum, 2 mg  
stop smoking aid  
**Gum**

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Gum



ACTUAL SIZE  
**Spearmint Burst**  
with a hint of chamomile

**COATED FOR BOLD FLAVOR** **160 PIECES, 2 mg EACH**

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid  
**Spearmint Burst Gum**  
with a hint of chamomile

824759

LOT: [REDACTED]

EXP: [REDACTED]

**NICORETTE**  
nicotine polacrilex gum, 2 mg | stop smoking aid

**Spearmint Burst Gum**  
with a hint of chamomile


not for sale to those under 18 years of age  
proof of age required  
not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

Nicorette is proud to offer a free individualized quit support program!  
Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.



3 07667 84360 6

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

**OPEN HERE**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Chew at least 9 pieces of Nicorette gum per day during the first six weeks.
- Use Long Enough** — Use Nicorette gum for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.





**170 PIECES, 2 mg EACH**

**Drug Facts**

**Active ingredient (in each chewing piece)**  
Nicotine polacrilex (equal to 2 mg nicotine) **Purpose**  
Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Includes Quit Support Program



**170 PIECES, 2 mg EACH**

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 94 mg, sodium 13 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients**  
acesulfame potassium, gum base, magnesium oxide, menthol, peppermint oil, sodium bicarbonate, sodium carbonate, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

NICO 0135-0229-05 HALEON



**170 PIECES, 2 mg EACH**

ACTUAL SIZE  
**Mint**



**170 PIECES, 2 mg EACH**

**Mint Gum**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

This product is protected in sealed blisters. Do not use if individual blisters or printed backings are broken, open, or torn.

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Sweden

©2023 Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the Haleon group of companies.

*Nicorette is proud to offer a free individualized quit support program! Visit [www.quit.com/join-myquit](http://www.quit.com/join-myquit) for details.*



3 07667 84340 8

824762

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	14 pt.
DRUG FACTS CONTINUED	8.5 pt.
HEADINGS	8.5 pt.
SUBHEADINGS/BODY TEXT	6.5 pt.
LEADING	6.6 pt.
# OF CHARACTERS PER INCH	< 39
BULLETS	5 pt.
SPACE BEFORE BULLET	2 spaces
BARLINES, HAIRLINES	1.5 pt / 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	25%
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
01/03/2024 11:30:02 AM